

European Birth Cohorts for Environmental Health Research

Martine Vrijheid,^{1,2,3} Maribel Casas,^{1,2,3} Anna Bergström,⁴ Amanda Carmichael,⁵ Sylvaine Cordier,⁶ Merete Eggesbø,⁷ Esben Eller,⁸ Maria P. Fantini,⁹ Mariana F. Fernández,^{3,10} Ana Fernández-Somoano,^{3,11} Ulrike Gehring,¹² Regina Grazuleviciene,¹³ Cynthia Hohmann,¹⁴ Anne M. Karvonen,¹⁵ Thomas Keil,¹⁴ Manolis Kogevinas,^{1,2,3,16} Gudrun Koppen,¹⁷ Ursula Krämer,¹⁸ Claudia E. Kuehni,^{19,20} Per Magnus,⁷ Renata Majewska,²¹ Anne-Marie Nybo Andersen,²² Evridiki Patelarou,²³ Maria Skaalum Petersen,²⁴ Frank H. Pierik,^{25,26} Kinga Polanska,²⁷ Daniela Porta,²⁸ Lorenzo Richiardi,²⁹ Ana Cristina Santos,³⁰ Rémy Slama,³¹ Radim J. Sram,³² Carel Thijs,³³ Christina Tischer,³⁴ Gunnar Toft,³⁵ Tomáš Trnovec,³⁶ Stephanie Vandentorren,³⁷ Tanja G.M. Vrijkotte,³⁸ Michael Wilhelm,³⁹ John Wright,⁴⁰ and Mark Nieuwenhuijsen^{1,2,3}

¹Centre for Research in Environmental Epidemiology (CREAL), Barcelona, Spain; ²Hospital del Mar Research Institute (IMIM), Barcelona, Spain; ³Spanish Consortium for Research on Epidemiology and Public Health (CIBERESP), Spain; ⁴Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden; ⁵School of Social and Community Medicine, University of Bristol, United Kingdom; ⁶French National Institute of Health and Medical Research (INSERM), U625; University of Rennes I; IFR140; Rennes, France; ⁷Division of Epidemiology, Norwegian Institute of Public Health, Oslo, Norway; ⁸Department of Dermatology and Allergy Centre, Odense University Hospital, Denmark; ⁹Department of Public Health, University of Bologna, Bologna, Italy; ¹⁰Laboratory of Medical Investigations, San Cecilio University Hospital, University of Granada, Granada, Spain; ¹¹University of Oviedo, Asturias, Spain; ¹²Institute for Risk Assessment Sciences, Utrecht University, Utrecht, the Netherlands; ¹³Department of Environmental Sciences, Vytautas Magnus University, Kaunas, Lithuania; ¹⁴Institute of Social Medicine, Epidemiology and Health Economics, Charité University Medical Center, Berlin, Germany; ¹⁵Department of Environmental Health, National Institute for Health and Welfare, Kuopio, Finland; ¹⁶National School of Public Health, Athens, Greece; ¹⁷Environmental Risk and Health Unit, Flemish Institute of Technological Research (VITO), Mol, Belgium; ¹⁸IUF - Leibniz Research Institute for Environmental Medicine (IUF), Düsseldorf, Germany; ¹⁹Institute of Social and Preventive Medicine (ISPM), University of Bern, Bern, Switzerland; ²⁰Department of Infection, Immunity and Inflammation, University of Leicester, Leicester, United Kingdom; ²¹Epidemiology and Preventive Medicine, Jagiellonian University Medical College, Kraków, Poland; ²²Section for Social Medicine, Department of Public Health, University of Copenhagen, Copenhagen, Denmark; ²³Department of Social Medicine, Medical School, University of Crete, Heraklion, Greece; ²⁴Department of Occupational Medicine and Public Health, The Faroese Hospital System, Tórshavn, Faroe Islands; ²⁵Department of Urban Environment, Netherlands Organisation for Applied Scientific Research (TNO), Utrecht, the Netherlands; ²⁶The Generation R Study Group, Erasmus Medical Center Rotterdam, Rotterdam, the Netherlands; ²⁷Nofer Institute of Occupational Medicine, Department of Environmental Epidemiology, Lodz, Poland; ²⁸Department of Epidemiology Regional Health Service, Lazio, Rome, Italy; ²⁹Cancer Epidemiology Unit, CPO-Piemonte and University of Turin, Turin, Italy; ³⁰Department of Hygiene and Epidemiology, University of Porto Medical School and Institute of Public Health, University of Porto, Porto, Portugal; ³¹INSERM and Grenoble University, Team of Environmental Epidemiology Applied to Reproduction and Respiratory Health, Institut Albert Bonniot (U823), Grenoble, France; ³²Institute of Experimental Medicine AS CR, Prague, Czech Republic; ³³Department of Epidemiology, School for Public Health and Primary Care (CAPHRI), Maastricht University, Maastricht, the Netherlands; ³⁴Helmholtz Zentrum, München, German Research Centre for Environmental Health, Institute of Epidemiology, Neuherberg, Germany; ³⁵Department of Occupational Medicine, Aarhus University Hospital, Aarhus, Denmark; ³⁶Slovak Medical University, Bratislava, Slovakia; ³⁷National Institute of Public Health Surveillance (InVS), Saint Maurice, France; ³⁸Department of Public Health, Academic Medical Centre, Amsterdam, The Netherlands; ³⁹Hygiene, Social and Environmental Medicine, Ruhr-University, Bochum, Germany; ⁴⁰Bradford Institute for Health Research, Bradford, United Kingdom

BACKGROUND: Many pregnancy and birth cohort studies investigate the health effects of early-life environmental contaminant exposure. An overview of existing studies and their data is needed to improve collaboration, harmonization, and future project planning.

OBJECTIVES: Our goal was to create a comprehensive overview of European birth cohorts with environmental exposure data.

METHODS: Birth cohort studies were included if they *a*) collected data on at least one environmental exposure, *b*) started enrollment during pregnancy or at birth, *c*) included at least one follow-up point after birth, *d*) included at least 200 mother–child pairs, and *e*) were based in a European country. A questionnaire collected information on basic protocol details and exposure and health outcome assessments, including specific contaminants, methods and samples, timing, and number of subjects. A full inventory can be searched on www.birthcohortsnetico.net.

RESULTS: Questionnaires were completed by 37 cohort studies of > 350,000 mother–child pairs in 19 European countries. Only three cohorts did not participate. All cohorts collected biological specimens of children or parents. Many cohorts collected information on passive smoking ($n = 36$), maternal occupation ($n = 33$), outdoor air pollution ($n = 27$), and allergens/biological organisms ($n = 27$). Fewer cohorts ($n = 12$ – 19) collected information on water contamination, ionizing or nonionizing radiation exposures, noise, metals, persistent organic pollutants, or other pollutants. All cohorts have information on birth outcomes; nearly all on asthma, allergies, childhood growth and obesity; and 26 collected information on child neurodevelopment.

CONCLUSION: Combining forces in this field will yield more efficient and conclusive studies and ultimately improve causal inference. This impressive resource of existing birth cohort data could form the basis for longer-term and worldwide coordination of research on environment and child health.

KEY WORDS: birth cohorts, child health, environmental exposures, Europe, review. *Environ Health Perspect* 120:29–37 (2012). <http://dx.doi.org/10.1289/ehp.1103823> [Online 29 August 2011]

It is well recognized that the fetus and infant are especially vulnerable to the effects of environmental risk factors that disrupt developmental processes, due to *a*) critical windows of vulnerability that occur during the rapid growth and development of organs and

systems, *b*) immaturities in metabolism, and *c*) greater intake and absorption of noxious agents in children relative to their body weight (Grandjean et al. 2008). Chemical, physical, and biological hazards in the environment may lead to serious health problems at birth and

during childhood, ranging from premature birth, low birth weight, and congenital anomalies to respiratory diseases, childhood cancer, learning disabilities, behavioral problems, and possibly even obesity (Trasande et al. 2009; Wigle et al. 2007). The economic and societal costs associated with nonoptimal child health and development are substantial. Furthermore, the effects of exposure to environmental risk factors may manifest themselves throughout a lifetime and even over generations. Common physical and mental diseases in adult life appear to have part of their origin in early life (Gillman et al. 2007; Kuh et al. 2003).

Epidemiological studies worldwide have investigated the role of early environmental

Address correspondence to M. Vrijheid, Centre for Research in Environmental Epidemiology, Doctor Aiguader, 88, 08003 Barcelona, Spain. Telephone: 34 932 147 346. Fax: 34 932 147 301. E-mail: [mvrijheid@creal.cat](mailto:m.vrijheid@creal.cat)

Supplemental Material is available online (<http://dx.doi.org/10.1289/ehp.1103823>).

We thank all other cohort researchers and technical staff who have helped with the completion of the questionnaires.

This work was supported by Environmental Health Risks in European Birth Cohorts (ENRIECO), a project conducted within the European Union's Seventh Framework Programme (Theme 6, Environment, including climate change), grant agreement 226285.

The authors declare they have no actual or potential competing financial interests.

Received 14 April 2011; accepted 29 August 2011.

contaminant exposures on pregnancy outcomes and child health (Wigle et al. 2007). Many have suffered from biases due to retrospective study designs, surrogate exposure and outcome measures, and incomplete confounder data.

Pregnancy and birth cohort studies are ideally suited to improve causal inference in this field, because they are designed to study the impacts of early exposures prospectively and at multiple time points during development

of the child. Furthermore, birth cohort studies usually collect biological material from mothers and children, enabling the measurement of biomarkers of exposure, early effect, or susceptibility. In the last two decades, new pregnancy

Table 1. General description of European birth cohorts with environmental contaminant data.

Birth cohort	Full name and key reference	Country	Regions covered	Enrollment period	No. of children at birth
Eligible cohorts that completed the ENRIECO inventory questionnaires:					
ABCD	Amsterdam Born Children and their Development study (van Eijsden et al. 2010)	Netherlands	Amsterdam	2003–2004	7,863
ALSPAC	The Avon Longitudinal Study of Parents and Children (Golding et al. 2001)	United Kingdom	Bristol	1991–1992	14,062
ArcRisk-Norway	Impacts on health in the Arctic and Europe owing to climate-induced changes in contaminant cycling	Norway	Troms, Finnmark, and Nordland	2007–2009	430
BAMSE	The Stockholm Children Allergy and Environmental Prospective Birth Cohort Study (Wickman et al. 2002)	Sweden	Stockholm	1994–1996	4,089
BiB	Born in Bradford (Raynor 2008)	United Kingdom	Bradford	2007–2010	13,000
Co.N.ER	Cohort of newborns in Emilia Romagna (Porta et al. 2006)	Italy	Bologna	2004–2005	654
Czech	Czech Republic Early Childhood Health (Dejmek et al. 2000)	Czech Republic	Teplice and Prachatice	1994–1999	7,577
DARC	The Danish Allergy Research Centre cohort (Johnke et al. 2005)	Denmark	Odense	1998–1999	562
DNBC	Danish National Birth Cohort (Olsen et al. 2001)	Denmark	Denmark	1996–2002	96,986
Duisburg	Duisburg cohort (Wilhelm et al. 2008)	Germany	Duisburg	2000–2003	234
EDEN	Study of determinants of pre and postnatal developmental, psychomotor development and child health (Drouillet et al. 2009)	France	Nancy, Poitiers	2003–2006	1,899
ELFE	French longitudinal study of children (Vandentorren et al. 2009)	France	France	2011–2012	20,000
Faroes ^a	Children's Health and the Environment in the Faroes (Grandjean et al. 1992, 1997)	Faroe Islands	Faroe Islands	1986–2009	2,351
FLEHS I	Flemish Environment and Health Survey (Koppen et al. 2009)	Belgium	Flanders	2002–2004	1,196
GASPII	Gene and Environment: Prospective Study on Infancy in Italy (Porta et al. 2007)	Italy	Rome	2003–2004	708
Generation R	Generation R (Jaddoe et al. 2008)	Netherlands	Rotterdam	2001–2006	9,778
Generation XXI	Generation XXI (Pinto et al. 2009)	Portugal	Porto	2005–2006	8,647
GINplus	German Infant Nutritional Intervention Study – plus influence of pollution and genetics on allergy development (Zirngibl et al. 2002)	Germany	Munich, Wesel	1995–1998	5,991
HUMIS	Norwegian Human Milk Study (Eggesbo et al. 2009)	Norway	Norway	2002–2011	2,500
INMA old ^b	Environment and Childhood (Guxens et al. 2011)	Spain	Granada, Menorca, Ribera d'Ebre	1997–2002	1,252
INMA new ^c	Environment and Childhood (Guxens et al. 2011)	Spain	Asturias, Gipuzkoa, Sabadell, Valencia	2003–2008	2,505
INUENDO ^d	Biopersistent organochlorines in diet and human fertility (Toft et al. 2005)	Greenland, Sweden, Poland, Ukraine	Greenland, Sweden (east and west coast), Warsaw (Poland), Kharkiv (Ukraine)	2002–2004	1,322
KANC	Kaunas cohort (Grazuleviciene et al. 2009)	Lithuania	Kaunas	2007–2009	4,000
KOALA	Child, parents and health: lifestyle and genetic constitution (Kummeling et al. 2005)	Netherlands	Southern Netherlands	2000–2003	2,834
Kraków	Kraków cohort (Jedrychowski et al. 2003)	Poland	Kraków	2000–2003	505
Leicester ^e	The Leicester Respiratory Cohorts (Kuehni et al. 2007)	United Kingdom	Leicestershire and Rutland	1985–1997	10,350
LISAplus	Influences of life-style related factors on the immune system and the development of allergies in childhood – plus the influence of traffic emissions and genetic (Heinrich et al. 2002)	Germany	Munich, Wesel, Leipzig, Bad Honneff	1997–1998	3,097
LUKAS	LUKAS (Karvonen et al. 2009)	Finland	Kuopio, Jyväskylä, Joensuu, and Iisalmi	2002–2005	442
MAS	Multicentre Allergy Study (Bergmann et al. 1994)	Germany	Berlin, Duesseldorf, Freiburg, Mainz, Munich	1990	1,314
MoBa	The Norwegian Mother and Child Cohort Study (Magnus et al. 2006)	Norway	Norway	1999–2008	107,000
NINFEA	Birth and Infancy: Effects of the Environment (Richiardi et al. 2007)	Italy	Italy	2005–	7,500
PARIS	Pollution and Asthma Risk: An Infant Study (Clarisse et al. 2007)	France	Paris	2003–2006	3,840
PCB cohort	Early Childhood Development and PCB exposures in Slovakia (Sonneborn et al. 2008)	Slovakia	Michalovce, Stropkov, Svidnik	2001–2003	1,134
PÉLAGIE	Endocrine disruptors: longitudinal study on pregnancy abnormalities, infertility, and childhood (Chevrier et al. 2011)	France	Brittany	2002–2006	3,421
PIAMA	Prevention and Incidence of Asthma and Mite Allergy (Brunekreef et al. 2002)	Netherlands	Northern, western and central parts	1996–1997	3,963
REPRO_PL	Polish Mother and Child Cohort (Polanska et al. 2009)	Poland	Lodz, Wroclaw, Lask, Kielce, Katowice, Legnica, Lublin, Szczecin, Piekary Slaskie	2007–2011	1,800
RHEA	Mother Child Cohort in Crete (Vardavas et al. 2010)	Greece	Heraklion, Crete	2007–2008	1,500
Potentially eligible cohorts that did not complete the ENRIECO inventory questionnaire:					
Ukraine	Children of Ukraine (Hryhorczuk et al. 2009)	Ukraine	Kyiv, Dniprodzerzhynsk, Mariupol	1992–1996	4,510
MAAS	The National Asthma Campaign Manchester Asthma and Allergy Study (Custovic et al. 2002)	United Kingdom	Manchester	1995–1997	1,211
Northern Adriatic Cohort	Northern Adriatic Cohort (Barbone et al. 2004)	Italy	Northeastern Italy	1999–2001	243

^aThe Faroes cohorts consist of four subcohorts: cohort 1 (enrollment: 1986–1987; *n* = 1022), cohort 2 (enrollment: 1994–1995; *n* = 182), cohort 3 (enrollment: 1997–2000, *n* = 656), cohort 5 (enrollment: 2007–2009; *n* = 491); ^bINMA old consists of three subcohorts: Granada (enrollment: 2000–2002; *n* = 668), Menorca (enrollment: 1997–1998; *n* = 482), Ribera Ebre (enrollment: 1997–1999; *n* = 102); ^cINMA new consists of four subcohorts: Asturias (enrollment: 2004–2007; *n* = 485), Gipuzkoa (enrollment: 2006–2008; *n* = 611), Sabadell (enrollment: 2004–2007; *n* = 622), Valencia (enrollment: 2003–2005; *n* = 787); ^dINUENDO consists of four subcohorts: Greenland, Sweden (east and west coast), Warsaw (Poland), Kharkiv (Ukraine); ^eThe Leicester Respiratory Cohorts consist of two subcohorts: the Leicester I cohort (enrollment: 1985–1989; *n* = 1,650) and the Leicester II cohorts (enrollment: 1993–1997; *n* = 8,700).

and birth cohorts have been set up across Europe (e.g., Golding et al. 2001; Hofman et al. 2004; Magnus et al. 2006; Olsen et al. 2001; Raynor 2008; Richiardi et al. 2007), in North America, and elsewhere (Brown et al. 2007; Herbstman et al. 2010; Marks et al. 2010; Richter et al. 2011), and more are being planned (Hirschfeld et al. 2011). Many cohort studies include assessments of environmental pollutant exposure, but not all have large expertise in this area. Although it is clear that individual cohorts are able to make, and have made, important contributions to understanding environmental causes of childhood disease and ill health, it is becoming increasingly clear that their full potential can be realized only with collaboration across large regions (Kogevinas et al. 2004).

The European Union (EU) has funded the Environmental Health Risks in European Birth Cohorts (ENRIECO) project to coordinate birth cohort research in Europe in the area of environmental contaminant exposures (ENRIECO 2009). ENRIECO recognized that a comprehensive overview of birth cohorts and their data was needed as a first step toward improved collaboration, future project planning, and effective use of existing data. Currently, no inventory of environmental exposure data exists. The aim of this review is to provide an overview of European birth cohorts with data on environmental contaminant exposures and health outcomes.

Methods

Definition and identification of cohorts. Birth cohort studies were included in the project if they *a*) collected data on at least one environmental contaminant exposure topic (air pollution, water contamination, pesticides, metals, persistent organic pollutants, other chemical pollutants, noise, radiations, and allergens and biological organisms); *b*) started enrollment of mothers into the cohort during pregnancy or at birth; *c*) included in their protocol at least one follow-up point after birth with direct contact with mothers and children; *d*) included at least 200 mother–child pairs (a low sample size limit was chosen so small studies with relevant exposure data or based in less well-covered regions were not excluded); and *e*) were based in a European country. Collection of data on an environmental exposure topic included data collected through questionnaires, environmental sampling, spatial maps and models, personal monitoring, biomarker measurements, or a combination of these methods. The mere availability of residential location (addresses and geocodes) was not sufficient for inclusion. All cohorts fitting the criteria were considered regardless of when they started, when they conducted their most recent follow-up, whether data had been published, or whether recruitment of subjects was ongoing.

Cohorts were identified through a variety of sources: www.birthcohorts.net (Birthcohorts.net 2010), a searchable Web site that contains basic protocol information for a number of birth cohorts; European projects using birth cohort data; and personal contacts. Publications and Web sites of identified cohorts were checked to see whether the cohort appeared to match the criteria of the inventory. All cohorts identified as potentially eligible between 1 March 2009 (start of the project) and 31 December 2010 were approached.

Inventory questionnaire. An inventory questionnaire was designed to collect detailed information from the cohorts on biological sample collection, assessment of environmental exposures, assessment of child health outcomes of interest, and basic information on collection of other variables. The full questionnaire can be found in Supplemental Material, Annex 1 (<http://dx.doi.org/10.1289/ehp.1103823>). The questionnaire was divided into four sections:

- Basic protocol description: protocol description, basic data, and biological sample collection scheme.
- Exposure assessment: air pollution (outdoor and indoor), water contamination, allergens and biological organisms, metals, pesticides, radiations (ionizing, ultraviolet, and non-ionizing), tobacco smoke (active and passive), noise, persistent organic pollutants,

occupation (maternal and paternal), and other chemical compounds. Each section asked for the exposure assessment methods used, numbers of subjects assessed, and time point of assessment.

- Health outcome assessment: reproduction and birth outcomes, neurodevelopment, allergies and asthma, cancer, child growth, and metabolic syndrome indicators such as blood pressure, cholesterol, triglycerides, insulin, or glucose. Each section asked which outcomes were assessed, by which methods (if relevant), how many subjects were assessed, and the time point of assessment.
- Other information: mainly yes/no questions on collection of genotype data, residential history, time–activity patterns, sociodemographic variables, breast-feeding, diet, physical exercise, medical history, and anthropometry.

The questionnaire was sent to principal investigators of each identified cohort, who were asked to distribute specific sections to the appropriate researchers and data managers within their cohort. The completed questionnaire was checked and, if necessary, the cohort study teams were contacted again to correct errors and complete information. Cohorts that did not reply to the first e-mail were reminded repeatedly by e-mail or phone. Cohorts were invited to two workshops during the project, in which the aims and objectives were presented.




Figure 1. European birth cohort studies collaborating in ENRIECO.

A Microsoft Access database was constructed to contain the responses from the cohorts. All cohorts were asked to check this database for errors before making it public. The last corrections and updates were made in February 2011. A Web-based searchable inventory is now publicly available on www.birthcohortsenrieco.net (ENRIECO 2011) and also accessible through www.birthcohorts.net (Birthcohorts.net 2010).

Results

A total of 46 birth cohort studies were identified as potentially eligible based on publications, Web sites, and word of mouth. Six of these were excluded after completion of

questionnaires, because they did not match the eligibility criteria: Three did not assess environmental exposures, and three did not have a follow-up point after birth. Another three birth cohorts did not return the questionnaires (Table 1). The 37 included cohorts, totaling > 350,000 study subjects, are situated in 19 European countries, principally in Northern and Western Europe (Figure 1). Five of the cohorts are in Eastern Europe and seven in Southern Europe (based on the European regions classified in the United Nations geoscheme). Twenty-five of the 37 cohorts defined themselves as regionally based studies, six as nationally based, four as hospital

based, and two as studying selected populations [farmers in the LUKAS cohort and high-risk allergic children in the Multicentre Allergy Study (MAS)]. Two studies, the Danish National Birth Cohort (DNBC) and the Norwegian Mother and Child Study (MoBa) have recruited around 100,000 mother-child pairs each (Table 1). Ten cohorts have recruited between 5,000 and 20,000 pairs, 18 between 1,000 and 5,000 pairs, and seven cohorts concern < 1,000 pairs.

At our reference date (February 2011), children in 19 cohorts were between 5 and 10 years of age, children in 12 cohorts were > 10 years of age, and in five cohorts they were < 5 years of age. One cohort in France, Etude Longitudinale Française depuis l'Enfance (ELFE), started data collection in April 2011. Most of the cohorts started recruitment of mothers during pregnancy ($n = 25$); the remaining started at birth. Most cohorts have multiple follow-up points after birth, and the majority have follow-up points in each of the child age periods specified in the questionnaire (1–6 months, 6–18 months, 18 months–5 years, 5–10 years, > 10 years) (Figure 2).

Biological samples were collected in all cohorts from the mother and/or the child and, more rarely, from the father (Table 2). Sampling covered prenatal and postnatal periods and many types of tissue, including blood, urine, hair, nails, breast milk, placentas, and saliva. All but six cohorts [The Stockholm Children Allergy and Environmental Prospective Birth Cohort Study (BAMSE); German Infant Nutritional Intervention Study—plus influence of pollution and genetics on allergy development (GINIplus); The Leicester Respiratory Cohorts (Leicester); Birth and Infancy: Effects of the Environment (NINFEA); Pollution and Asthma Risk: An Infant Study (PARIS); Prevention and Incidence of Asthma and Mite Allergy (PIAMA)] collected, or plan to collect, blood samples from the mother during pregnancy or cord blood at birth.

Many cohort studies collected information on passive smoking ($n = 36$), maternal occupation ($n = 33$), outdoor air pollution ($n = 27$), and allergens/biological organisms ($n = 27$) (Table 3). Fewer cohorts have collected information on water contamination ($n = 13$), ionizing or nonionizing radiation exposures ($n = 12$), noise ($n = 14$), or emerging chemical exposures such as phthalates and bisphenol A ($n = 17$). Details of each assessment (methods, numbers of subjects, and time period of assessment) can be found in the Web inventory (ENRIECO 2011). Twenty-seven cohorts have measured or are measuring contaminants or their metabolites as markers of internal dose, mostly in the categories of metals, persistent organic pollutants, and tobacco smoke (Table 4).




Figure 2. Start of enrollment and time points of follow-up^a (vertical bars) in European birth cohorts collaborating in ENRIECO. T1, T2, and T3 are first, second, and third trimesters. mo, months. See Table 1 for full names and locations of cohorts.

^aThese are points in predefined periods. Some cohorts have many more (e.g., yearly) follow-up points not reflected in this figure. The four Faroës cohorts and the two Leicester cohorts are shown separately because they have different time lines.

All cohorts have information on birth outcomes, mostly birth weight, gestational age, and mode of delivery [see Supplemental Material, Annex 2 (<http://dx.doi.org/10.1289/ehp.1103823>)]. Fewer asked questions about time to pregnancy or collected data on congenital anomalies. Fifteen cohorts collected information on ultrasound measurements. Neurodevelopment and behavioral development of the children was assessed in 26 cohorts, using a large variety of methods including questionnaires, neuropsychological tests, and school results. Nineteen cohorts assessed cognitive function of the children (e.g., global IQ, executive function, memory, etc.), 18 cohorts assessed behavior or ADHD (attention deficit/hyperactivity disorder) symptoms, and few have data on the clinical diagnosis of autism ($n = 13$) or mental health problems ($n = 3$). Names of the questionnaires and tests used to assess neurodevelopment, as well as the developmental periods at which

they were used are available in the Web inventory. Growth and obesity and allergies ($n = 34$), and asthma and respiratory infections ($n = 33$) were assessed in nearly all cohorts. Metabolic syndrome indicators, such as blood pressure or cholesterol levels, were collected in around half of the cohorts. Childhood cancer ($n = 12$) and sexual maturation ($n = 14$) have been assessed in few cohorts to date [see Supplemental Material, Annex 2 (<http://dx.doi.org/10.1289/ehp.1103823>)].

Discussion

More than 35 birth cohorts in Europe, studying more than 350,000 mother-child pairs, have data that can be used to examine associations between chemical, physical, and biological environmental exposures and child health. The searchable Web-based inventory is the first place where this information is made publicly available. The inventory intends to facilitate collaborations between birth cohort researchers

in this field and can be searched to identify cohorts relevant for comparison, replication, and combined studies. Moreover, it may be used by policy makers and other stakeholders to identify birth cohorts that can provide specific information on environmental exposures or related outcomes. This initiative may quite easily be expanded beyond Europe to stimulate collaboration between cohorts worldwide.

The cohorts we evaluated include a few very large general cohorts with multiple aims as well as many smaller cohorts, which are generally more focused on specific environmental risk factors or specific health outcomes. The distribution of the identified birth cohorts is not homogeneous across Europe, with more and larger cohorts mostly located in the north and west of Europe and fewer and smaller cohorts in the south and east. Indeed, more than half of all mother-child pairs come from two very large cohorts in Norway and Denmark. However, studies of

Table 2. Biological samples collected in the European birth cohorts collaborating in ENRIECO.^a

Cohort	Mother					Child						Father	
	T1	T2	T3	Birth	Postnatal	Birth	1-6 months	7-18 months	19 months-5 years	5-10 years	≥ 10 years	Prenatal	Postnatal
ABCD	B									B			
ALSPAC	B, U	B, U	B, U	P	H, N	CB		H, N	B, H, N	B, U, T	B, U, S		H, N
ArcRisk-Norway		B, U		B, U, H	B, U, BM	CB, B, M							
BAMSE									B, U	B			
BiB			B, U			CB		B	B			S	
Co.N.ER				B		CB							
Czech				B, P		CB			U	U			
DARC						CB	B	B	B	B			
DNBC	B	B				CB	B						
Duisburg			B		B, U, BM	CB			B, U	B, U			
EDEN		B, U, S		H, BM, P		CB, H, M		H	H	B, U, H		B	
ELFE				B, U, BM, H		CB, CT, M			U, H, S	U			
Faroese			B	H, BM	B, U, H, BM	CB	B	B, H	B, H	B, H	B, H		U, H, Se
FLEHS I					S	CB	F	F	U	U			
GASPII				B		CB							
Generation R	B, U	B, U	B, U			CB			S	B, U, S		B	
Generation XXI	B	B	B, AT		B	CB			B				B
GINplus									B	B	B		
HUMIS ^b		B, U		B	BM	CB				S		B	
INMA old	B	B	B	B, BM, P	B, BM	CB, U	B	B	B, U, H, S	B, U		B	
INMA new	B, U		U, N	BM, P	U, S	CB, H		U, N, S	H, S	B, U, H, S			B, S
INUENDO	B	B	B							S		B, Se	
KANC			B			B							
KOALA			B		BM, S		F	S	B	B, F			S
Kraków				B, P	U	CB, M			U	B, U			U
Leicester											S		
LISAplus						CB, B			B	B	B		
LUKAS				B, P	BM	CB	F	B, F	B	B, U			B
MAS				B		CB, B		B, U	B, U	B, U	B		B
MoBa		B, U		B		CB						B	
NINFEA					S		S						
PARIS									B	B, U			
PCB cohort				B, BM		CB	B	B	B	B, U			
PELAGIE	U			H, P		CB				S, U			
PIAMA					B, BM, S		B	B	B, S	B	B, S		B, S
REPRO_PL	B, S	B, U, S	B, U, H, S	B	BM	CB		B, U	B, U				
RHEA	B, U	U		B		CB		U, H	B				B, Se

Abbreviations: AT, adipose tissue; B, blood; BM, breast milk; CB, cord blood; CT, cord tissue; F, feces; H, hair; M, meconium; N, nails; P, placenta; S, saliva or buccal swabs; Se, semen; T, teeth; U, urine. T1, T2, and T3 are first, second, and third trimesters. See Table 1 for full names and locations of cohorts.

^aMany but not all ENRIECO birth cohorts are also included in the BBMRI Catalogue of European Biobanks (BBMRI 2011), which can be queried by specimen type. ^bBlood, urine, and cord blood are available through overlap with MoBa in approximately 800 participants.

environmental contaminant exposures, especially those measuring exposure biomarkers, usually cannot cover large numbers of subjects. Smaller, specialized cohorts, often located in high-exposure areas, have made very important contributions in this regard.

Passive smoking, maternal (and paternal, less often) occupation, air pollution, and allergen exposures have been assessed by many birth cohorts. Efforts to combine data analyses on these topics are already ongoing within international projects such as European Study of Cohorts for Air Pollution Effects (ESCAPE) (2009), Global Allergy and Asthma European Network (GA2LEN) (2009), and International Collaboration on Air Pollution and Pregnancy Outcomes (ICAPPO) (Woodruff et al. 2010). Biomarker data on some metals (lead and mercury) and some persistent organic pollutants [polychlorinated biphenyls (PCBs) and dichlorodiphenyldichloroethylene (DDE)] are also available in a relatively large number of

cohorts, but usually for a small number of subjects in each cohort and with considerable variation in the type and timing of the biological samples. Studies combining data from different cohorts have been conducted for few such pollutants so far; examples are the 7-cohort lead meta-analysis (Lanphear et al. 2005) and a recent meta-analysis of 12 European cohorts evaluating PCB and DDE exposure and fetal growth (Govarts et al. 2011). Exposure areas that are less well covered in the cohorts include noise, ionizing and nonionizing radiations, and chemical exposures such as brominated and fluorinated compounds, phthalates, and phenols (including bisphenol A). These are areas of emerging concern for child health and high on international research agendas [e.g., the World Health Organization (WHO) Research Agenda for Radiofrequency Fields] (WHO 2010). Although actual measurements are scarce, future assessments of these exposures are planned in many cohorts, making

this an optimal time to work toward a high level of data comparability.

Birth cohorts are most suitable to examine child health outcomes that are common or can be measured on a continuous scale. Rare outcomes and pathologies such as extremely preterm birth, childhood cancers, congenital anomalies, and autism can be studied only in the very large cohorts or by combining cohorts, as is being done, for example, in the International Childhood Cancer Cohort Consortium (Brown et al. 2007). All cohorts collected information on fetal growth and gestational length and on childhood anthropometric measures; combined studies of these outcomes appear feasible. A number of cohorts specialize in the early causes of asthma and allergies, and combined studies of these outcomes are underway (Keil et al. 2006a, 2006b). Neurodevelopment is a wide area that encompasses many cognitive and behavioral outcomes and includes even more assessment

Table 3. Assessment of environmental exposures (prenatal or postnatal) in European birth cohorts participating in ENRIECO.^a

Cohort	Air pollution outdoor	Air pollution indoor	Water contamination	Allergens and biological organisms	Metals ^b	Pesticides ^c	Persistent organic pollutants	Other chemical exposures ^d	Radiations	Passive smoking ^e	Noise	Occupation
ABCD	X					X			X ^g	X		X
ALSPAC		X		X	X	X	X ^g		X	X	X	X
ArcRisk-Norway					X		X	X ^g		X		X
BAMSE	X	X		X						X		X
BiB	X		X									X
Co.N.ER	X ^g	X		X						X	X	X
Czech	X									X		X
DARC	X	X		X						X		X
DNBC	X ^g	X		X		X	X		X	X		X
Duisburg	X	X	X ^g	X	X		X	X		X		X
EDEN	X	X	X ^g		X			X	X	X		X
ELFE	X ^g	X ^g	X ^g	X ^g	X ^g	X ^g	X ^g	X ^g	X ^g	X ^g		X ^g
Faroes					X	X ^g	X	X ^g		X		X
FLEHS I	X	X		X	X	X	X			X		X
GASPII	X	X		X						X	X	
Generation R	X			X		X	X ^g	X		X	X	X
Generation XXI				X						X		X
GINIplus	X	X		X						X		
HUMIS ^f	X	X	X	X	X ^g	X	X	X	X	X		X
INMA old	X	X	X	X	X ^g	X	X	X	X	X	X	X
INMA new	X	X	X	X	X	X	X	X	X	X	X	X
INUENDO					X ^g	X ^g	X ^g	X ^g		X		X
KANC	X		X							X	X ^g	X
KOALA		X	X	X					X ^g	X	X	X
Kraków	X	X		X	X	X	X	X		X		X
Leicester	X	X		X						X		X
LISAplus	X	X		X						X	X ^g	
LUKAS		X		X	X	X	X	X ^g		X		X
MAS		X		X						X	X	X
MoBa	X ^g	X	X	X	X ^g	X	X ^g	X	X	X	X	X
NINFEA	X ^g	X ^g		X ^g		X ^g		X ^g	X ^g	X ^g	X ^g	X ^g
PARIS	X	X	X	X						X		X
PCB cohort		X			X	X	X	X ^g		X		X
PELAGIE		X	X		X	X	X	X		X		X
PIAMA	X	X		X						X		
REPRO_PL	X ^g			X	X		X			X	X	X
RHEA	X	X	X	X	X	X	X ^g	X ^g	X	X	X	X

See Table 1 for full names and locations of cohorts.

^aDetails of each assessment (method, time period, number of subjects) are available at www.birthcohortsenrieco.net (ENRIECO 2011). ^bBiomonitoring data only, not questionnaires.

^cOrganochlorine pesticides are included under persistent organic pollutants. ^dPhthalates and phenols (including bisphenol A). ^ePassive smoking exposure of the mother during pregnancy or of the child postnatally. ^fData on occupation and passive smoking are available through overlap with MoBa in approximately 800 participants. ^gMeasurements are planned or ongoing but not completed.

instruments, ranging from parental questionnaires on developmental milestones to detailed neuropsychological assessments. Combined analyses may therefore prove difficult and may need to be restricted to cohorts that use similar tests in similar age groups.

This inventory inevitably simplifies the data available in the cohorts. For example, we summarized time periods over which data were collected, meaning that not each follow-up point in each cohort is represented. Furthermore, we restricted information about exposure and health outcome assessment methods to relatively basic questions [see Supplemental Material, Annex 1 (<http://dx.doi.org/10.1289/ehp.1103823>)]. It is also inevitable that the inventory information

will become out of date quickly. The Web-based database will be maintained by sending requests for updates to the included and newly identified cohorts at approximately yearly intervals for as long as resources allow. The inventory is meant to give researchers and other professionals looking for specific data the first overview of relevant cohorts; only direct contact with the cohorts can then establish the suitability of data for a specific purpose. Beyond our overview here, the ENRIECO project has made an extensive effort to compare available data and methods across cohorts and to develop recommendations for the use of existing data and for the collection of data in the future. These reports are available on request (ENRIECO 2009).

The 37 European birth cohorts described in this review fulfilled all our inclusion criteria. We also identified other studies that did not strictly match our criteria of a birth cohort study or did not collect data on the environmental exposures. We may have missed other birth cohorts that have not yet been published or are still in planning stages. In addition, many important birth cohorts outside Europe were not included. The Web-based inventory will be open for the inclusion of additional birth cohorts for the foreseeable future. Overall, we have achieved an excellent response from the European cohorts; only three potentially eligible cohorts did not return questionnaires. Involvement of the cohorts in workshops, working groups, and case studies was instrumental in these results.

Table 4. Biomarkers of selected environmental exposures measured by European birth cohorts collaborating in ENRIECO.^a

Cohort	Metals	Persistent organic pollutants	Other pesticides	Tobacco smoking	Other chemicals
ALSPAC	As, Cd, Hg, Mn, Pb, Se, ^b TMS	PFCs ^b	—	Cotinine	—
ArcRisk-Norway	As, Cd, Co, Hg, Mb, Mn, Pb	Chlordane, DDT/DDE, HCB, PCBs	—	—	Planned
BAMSE	—	—	—	Cotinine	—
Czech	—	—	—	Cotinine	PAHs
DNBC	—	PFCs	—	—	—
Duisburg	Cd, Hg, Pb, Se	DDT/DDE, HCB, ^b HCH, PCBs, PCDDs, PCDFs, PFCs	—	Cotinine ^b	BPA, phthalates
EDEN	B, Cd, Mn, Hg, Pb	—	—	Cotinine	Phthalates, phenols (including BPA)
ELFE	Al, ^b As, ^b Cd, ^b Hg, ^b Pb ^b	BFRs, ^b organochlorines, ^b PFCs, ^b PCBs, ^b PCDDs, ^b PCDFs ^b	Organophosphates, ^b pyrethroids ^b	Cotinine ^b	BPA, ^b phthalates ^b
Faroese	Hg, Pb, Se	BFRs, ^b chlordane, DDT/DDE, dieldrin/endrin, heptachlor, HCB, β -HCH, mirex, organometallic compounds, PCBs, PFCs, ^b toxaphene	—	—	BPA, ^b phthalates ^b
FLEHS I	Cd, Pb	DDT/DDE, dioxin-like compounds, HCB, PCBs	—	—	—
Generation R	—	Organochlorines ^b	Organophosphates	—	BPA, phthalates
GINplus	—	—	—	Cotinine	—
HUMIS	Hg, ^b Pb ^b	BFRs, DDT/DDE, HCB, HCH, mirex, PCBs, PCDDs, PCDFs, PFCs, toxaphene	—	—	Phthalates
INMA old	Hg, Pb, ^b TMS ^b	Aldrin/dieldrin/endrin, BFRs, DDT/DDE, endosulfan, lindane, methoxychlor, mirex, HCB, HCH, PCBs	—	Cotinine	Phthalates, phenols (including BPA)
INMA new	Hg, Pb, TMS ^b	BFRs, DDT/DDE, HCB, HCH, PCBs	—	Cotinine	Phthalates, phenols (including BPA)
INUENDO	Cd, ^b Hg, ^b Pb ^b	BFRs, ^b DDT/DDE, HCB, ^b PCBs, PFCs ^b	—	—	BPA, ^b phthalates ^b
Kraków	Cd, Hg, Pb	—	—	Cotinine	Phthalates, PAHs, benzo[a]pyrene-adducts
LISAplus	—	—	—	Cotinine	—
LUKAS	As, Cd, Hg, Pb, Se	BFRs, DDT/DDE, organometallic compounds, PCBs, PCDDs, PCDFs, polychlorinated naphthalene	—	—	Phthalates ^b
MAS	—	—	—	Cotinine	—
MoBa	Planned	BFRs, ^b DDT/DDE, ^b PCBs ^b	Organophosphates	—	BPA, phthalates
NINFEA	—	—	—	—	—
PCB cohort	Hg, Pb	DDT/DDE, HCB, HCH, PCBs, PFCs ^b	—	—	Phthalates ^b
PÉLAGIE	Hg	Aldrin, BFRs, DDT/DDE, dieldrin/endrin, heptachlor, HCB, PCBs	Acetochlor, alachlor, metolachlor, organophosphorus, propoxur, triazines	—	Phthalates
REPRO_PL	Cd, Hg, Pb, Se, Zn, Cu	PCBs, PCDDs, PCDFs	—	Cotinine	PAHs (1-hydroxypyrene)
RHEA	As, Cd, Hg, Mn, Pb	DDT/DDE, ^b HCB, ^b PCBs, ^b PCDDs, ^b PFCs ^b	—	NNAL, cotinine	Phthalates ^b

Abbreviations: As, arsenic; B, boron; BFRs, brominated flame retardants; BPA, bisphenol A; Cd, cadmium; Co, cobalt; Cu, copper; DDE, dichlorodiphenyldichloroethylene; DDT, dichlorodiphenyltrichloroethane; HCB, hexachlorobenzene; HCH, hexachlorocyclohexane; Hg, mercury; Mb, molybdenum; Mn, manganese; Ni, nickel; NNAL, 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; PAHs, polycyclic aromatic hydrocarbons; Pb, lead; PCBs, polychlorinated biphenyls; PCDDs, polychlorinated dibenzo-*p*-dioxins; PCDFs, polychlorinated dibenzofurans; PFCs, perfluorinated compounds; Se, selenium; Tl, thallium; TMS, total metals spectrum; Zn, zinc. See Table 1 for full names and locations of cohorts.

^aIncludes all exposures biomarkers measured as part of the exposure topics included the inventory questionnaire [see Supplemental Material, Annex 1 (<http://dx.doi.org/10.1289/ehp.1103823>)]. Details of each assessment (method, time period, number of subjects) are available at www.birthcohortsenrieco.net (ENRIECO 2011). ^bMeasurements are planned or ongoing but not completed.

Future collaboration. The 350,000 mother–child pairs covered by European birth cohorts form an impressive number, but of course this is only a small percentage of all births in Europe [(> 5 million births annually in 25 EU countries (Eurostat 2011)]. For the investigation of any specific exposure–outcome relationship, the number of available subjects will be much smaller than the total number of children followed by the cohorts. The cohorts are in very different stages of follow-up and thus have not all reached the ages at which some outcomes, for example, asthma or puberty onset, can be assessed. Moreover, not all exposure and outcome topics included in ENRIECO are immediately suitable for combined analyses of individual data. However, there are many reasons why continued and tightly coordinated collaboration between existing cohorts is desirable (Kogevinas et al. 2004; Willett et al. 2007): *a*) combined analyses of cohort data to improve statistical power and reduce publication bias; *b*) improved contrast and diversity in exposure and outcome; *c*) faster response to concerns about new environmental exposures; *d*) facilitation of replication and comparison studies in different settings; *e*) facilitation of linkage to large-scale, routinely collected environmental and health data; and *f*) widening expertise to improve methodological approaches, including protocols of biological and environmental sample collection and analysis.

In conclusion, the impressive resource of existing data in birth cohorts should form the basis for a long-term infrastructure that brings together birth cohort research on early-life environmental contaminant exposure and child health, with the ultimate aim to improve causal inference in this field. Further expansion to countries outside Europe should be one important aim for the future. This will be possible only with continued support for coordination.

REFERENCES

- Barbone F, Valent F, Pisa F, Daris F, Fajon V, Gibicari D, et al. 2004. Prenatal low-level methyl mercury exposure and child development in an Italian coastal area. *Seychelles Med Dent J, Special Issue* 7.
- BBMRI (Biobanking and Biomolecular Resources Research Infrastructure). 2011. Homepage. Available: <http://www.bbMRI.eu/> [accessed 6 July 2011].
- Bergmann RL, Bergmann KE, Lau-Schadensdorf S, Luck W, Danneberg A, Bauer CP, et al. 1994. Atopic diseases in infancy. The German multicenter atopy study (MAS-90). *Pediatr Allergy Immunol* 5:19–25.
- Birthcohorts.net. 2010. Homepage. Available: <http://www.birthcohorts.net/> [accessed 6 July 2011].
- Brown RC, Dwyer T, Kasten C, Krotoski D, Li Z, Linet MS, et al. 2007. Cohort profile: the International Childhood Cancer Cohort Consortium (I4C). *Int J Epidemiol* 36:724–730.
- Brunekeef B, Smit J, de Jongste J, Neijens H, Gerritsen J, Postma D, et al. 2002. The prevention and incidence of asthma and mite allergy (PIAMA) birth cohort study: design and first results. *Pediatr Allergy Immunol* 13(suppl 15):55–60.
- Chevrier C, Limon G, Monfort C, Rouget F, Garlandezec R, Petit C, et al. 2011. Urinary biomarkers of prenatal atrazine exposure and adverse birth outcomes in the PELAGIE birth cohort. *Environ Health Perspect* 119:1034–1041.
- Clarisse B, Nikasinovic L, Poinard R, Just J, Momas I. 2007. The Paris prospective birth cohort study: which design and who participates? *Eur J Epidemiol* 22:203–210.
- Custovic A, Simpson BM, Murray CS, Lowe L, Woodcock A. 2002. The National Asthma Campaign Manchester Asthma and Allergy Study. *Pediatr Allergy Immunol* 13(suppl 15):32–37.
- Dejmek J, Solansky I, Benes I, Lenček J, Šrám RJ. 2000. The impact of polycyclic aromatic hydrocarbons and fine particles on pregnancy outcome. *Environ Health Perspect* 108:1159–1164.
- Drouillet P, Kaminski M, Lauzon-Guillain B, Forhan A, Ducimetiere P, Schweitzer M, et al. 2009. Association between maternal seafood consumption before pregnancy and fetal growth: evidence for an association in overweight women. The EDEN mother-child cohort. *Paediatr Perinat Epidemiol* 23:76–86.
- Eggesbo M, Stigum H, Longnecker MP, Polder A, Aldrin M, Basso O, et al. 2009. Levels of hexachlorobenzene (HCB) in breast milk in relation to birth weight in a Norwegian cohort. *Environ Res* 109:559–566.
- ENRIECO (Environmental Health Risks in European Birth Cohorts). 2009. Enrieco Project: Environmental Health Risks in European Birth Cohorts. Available: <http://www.enrieco.org/> [accessed 6 July 2011].
- ENRIECO (Environmental Health Risks in European Birth Cohorts). 2011. Inventory of ENRIECO Cohorts. 2011. Available: <http://www.birthcohortsenrieco.net/> [accessed 6 July 2011].
- ESCAPE (European Study of Cohorts for Air Pollution Effects). 2009. Homepage. Available: <http://www.escapeproject.eu/> [accessed 6 July 2011].
- Eurostat 2011. Homepage. Available: <http://epp.eurostat.ec.europa.eu/> [accessed 3 March 2011].
- GAZLEN (Global Allergy and Asthma European Network). 2009. Homepage. Available: <http://www.gazlen.net/> [accessed 6 July 2011].
- Gillman MW, Barker D, Bier D, Gagampang F, Challis J, Fall C, et al. 2007. Meeting report on the 3rd International Congress on Developmental Origins of Health and Disease (DOHaD). *Pediatr Res* 61:625–629.
- Golding J, Pembrey M, Jones R. 2001. ALSPAC—the Avon Longitudinal Study of Parents and Children. I. Study methodology. *Paediatr Perinat Epidemiol* 15:74–87.
- Govarts E, Nieuwenhuijse M, Schoeters G, Ballester F, Bloemen K, Weihe P, et al. 2011. Prenatal exposure to polychlorinated biphenyls (PCB) and dichlorodiphenyl-dichloroethylene (DDE) and fetal growth: a meta-analysis within 12 European birth cohorts. *Environ Health Perspect*; doi: 10.1289/ehp.1103823 [Online 13 October 2011].
- Grandjean P, Bellinger D, Bergman A, Cordier S, Davey-Smith G, Eskenazi B, et al. 2008. The Faroes statement: human health effects of developmental exposure to chemicals in our environment. *Basic Clin Pharmacol Toxicol* 102:73–75.
- Grandjean P, Weihe P, Jorgensen PJ, Clarkson T, Cernichiari E, Videro T. 1992. Impact of maternal seafood diet on fetal exposure to mercury, selenium, and lead. *Arch Environ Health* 47:185–195.
- Grandjean P, Weihe P, White RF, Debes F, Araki S, Yokoyama K, et al. 1997. Cognitive deficit in 7-year-old children with prenatal exposure to methylmercury. *Neurotoxicol Teratol* 19:417–428.
- Grazuleviciene R, Danileviciute A, Nadisauskiene R, Vencloviene J. 2009. Maternal smoking, GSTM1 and GSTT1 polymorphism and susceptibility to adverse pregnancy outcomes. *Int J Environ Res Public Health* 6:1282–1297.
- Guxens M, Ballester F, Espada M, Fernandez MF, Grimalt JO, Ibarluzea J, et al. 2011. Cohort Profile: The INMA—Infancia y Medio Ambiente—(Environment and Childhood) Project. *Int J Epidemiol*; doi: 10.1093/ije/dyr054 [Online 5 April 2011].
- Heinrich J, Bolte G, Holscher B, Douwes J, Lehmann I, Fahlbusch B, et al. 2002. Allergens and endotoxin on mothers' mattresses and total immunoglobulin E in cord blood of neonates. *Eur Respir J* 20:617–623.
- Herbstman JB, Sjödin A, Kurzon M, Lederman SA, Jones RS, Rauh V, et al. 2010. Prenatal exposure to PBDEs and neurodevelopment. *Environ Health Perspect* 118:712–719.
- Hirschfeld S, Songco D, Kramer BS, Guttmacher AE. 2011. National Children's Study: update in 2010. *Mt Sinai J Med* 78:119–125.
- Hofman A, Jaddoe VW, Mackenbach JP, Moll HA, Snijders RF, Steegers EA, et al. 2004. Growth, development and health from early fetal life until young adulthood: the Generation R Study. *Paediatr Perinat Epidemiol* 18:61–72.
- Hryhorczuk D, Dardynska I, Lukyanova E, Matwyszyn-Fuoco M, Friedman L, Shkiryak-Nizhnyk Z, et al. 2009. Risk factors for wheezing in Ukrainian children: Ukraine European Longitudinal Study of Pregnancy and Childhood Group. *Paediatr Perinat Epidemiol* 23:346–351.
- Jaddoe VW, van Duijn CM, van der Heijden AJ, Mackenbach JP, Moll HA, Steegers EA, et al. 2008. The Generation R Study: design and cohort update until the age of 4 years. *Eur J Epidemiol* 23:801–811.
- Jedrychowski W, Whyatt RM, Camann DE, Bawle UV, Peki K, Spengler JD, et al. 2003. Effect of prenatal PAH exposure on birth outcomes and neurocognitive development in a cohort of newborns in Poland. Study design and preliminary ambient data. *Int J Occup Med Environ Health* 16:21–29.
- Johnke H, Vach W, Norberg LA, Bindlev-Jensen C, Host A, Andersen KE. 2005. A comparison between criteria for diagnosing atopic eczema in infants. *Br J Dermatol* 153:352–358.
- Karvonen AM, Hyvarinen A, Roponen M, Hoffmann M, Korppi M, Remes S, et al. 2009. Confirmed moisture damage at home, respiratory symptoms and atopy in early life: a birth-cohort study. *Pediatrics* 124:e329–e338.
- Keil T, Kulig M, Simpson A, Custovic A, Wickman M, Kull I, et al. 2006a. European birth cohort studies on asthma and atopic diseases: I. Comparison of study designs—a GALEN initiative. *Allergy* 61:221–228.
- Keil T, Kulig M, Simpson A, Custovic A, Wickman M, Kull I, et al. 2006b. European birth cohort studies on asthma and atopic diseases: II. Comparison of outcomes and exposures—a GAZLEN initiative. *Allergy* 61:1104–1111.
- Kogevinas M, Andersen AM, Olsen J. 2004. Collaboration is needed to co-ordinate European birth cohort studies. *Int J Epidemiol* 33:1172–1173.
- Koppen G, Den Hond E, Nelen V, Van De ME, Bruckers L, Bilau M, et al. 2009. Organochlorine and heavy metals in newborns: results from the Flemish Environment and Health Survey (FLEHS 2002–2006). *Environ Int* 35:1015–1022.
- Kuehni CE, Brooke AM, Strippoli MP, Spycher BD, Davis A, Silverman M. 2007. Cohort profile: the Leicester respiratory cohorts. *Int J Epidemiol* 36:977–985.
- Kuh D, Ben Shlomo Y, Lynch J, Hallqvist J, Power C. 2003. Life course epidemiology. *J Epidemiol Community Health* 57:778–783.
- Kummeling I, Thijs C, Penders J, Snijders BE, Stelma F, Reimerink J, et al. 2005. Etiology of atopy in infancy: the KOALA Birth Cohort Study. *Pediatr Allergy Immunol* 16:679–684.
- Lanphear BP, Hornung R, Khoury J, Yolton K, Baghurst P, Bellinger DC, et al. 2005. Low-level environmental lead exposure and children's intellectual function: an international pooled analysis. *Environ Health Perspect* 113:894–899.
- Magnus P, Irgens LM, Haug K, Nystad W, Skjaerven R, Stoltenberg C. 2006. Cohort profile: the Norwegian Mother and Child Cohort Study (MoBa). *Int J Epidemiol* 35:1146–1150.
- Marks AR, Harley K, Bradman A, Kogut K, Barr DB, Johnson C, et al. 2010. Organophosphate pesticide exposure and attention in young Mexican-American children: the CHAMACOS study. *Environ Health Perspect* 118:1768–1774.
- Olsen J, Melbye M, Olsen SF, Sorensen TI, Aaby P, Andersen AM, et al. 2001. The Danish National Birth Cohort—its background, structure and aim. *Scand J Public Health* 29:300–307.
- Pinto E, Barros H, dos Santos Silva I. 2009. Dietary intake and nutritional adequacy prior to conception and during pregnancy: a follow-up study in the north of Portugal. *Public Health Nutr* 12:922–931.
- Polanska K, Hanke W, Gromadzinska J, Ligocka D, Gulczynska E, Sobala W, et al. 2009. Polish mother and child cohort study—defining the problem, the aim of the study and methodological assumption. *Int J Occup Med Environ Health* 22:383–391.
- Porta D, Fantini MP, on behalf of the GASPIII and Co.NER Study Groups. 2006. Prospective cohort studies of newborns in Italy to evaluate the role of environmental and genetic characteristics on common childhood disorders. *Ital J Pediatr* 32:350–357.
- Porta D, Forastiere F, Di Lallo D, Perucci CA. 2007. Enrolment and follow-up of a birth cohort in Rome [in Italian]. *Epidemiol Prev* 31:303–308.
- Raynor P. 2008. Born in Bradford, a cohort study of babies born in Bradford, and their parents: protocol for the recruitment phase. *BMC Public Health* 8:327; doi:10.1186/1471-2458-8-327 Sept 21 2008 [Online 23 September 2008].
- Richiardi L, Baussano I, Vizzini L, Douwes J, Pearce N, Merletti F. 2007. Feasibility of recruiting a birth cohort

- through the Internet: the experience of the NINFEA cohort. *Eur J Epidemiol* 22:831–837.
- Richter LM, Victora CG, Hallal PC, Adair LS, Bhargava SK, Fall CH, et al. 2011. Cohort profile: the Consortium of Health-Orientated Research in Transitioning Societies. *Int J Epidemiol*; doi:10.1093/ije/dyq251 [Online 21 April 2011].
- Sonneborn D, Park HY, Petrik J, Kocan A, Palkovicova L, Trnovec T, et al. 2008. Prenatal polychlorinated biphenyl exposures in eastern Slovakia modify effects of social factors on birthweight. *Paediatr Perinat Epidemiol* 22:202–213.
- Toft G, Axmon A, Giwercman A, Thulstrup AM, Rignell-Hydbom A, Pedersen HS, et al. 2005. Fertility in four regions spanning large contrasts in serum levels of widespread persistent organochlorines: a cross-sectional study. *Environ Health* 4:26; doi: 10.1186/1476-069X-4-26 [Online 9 November 2005].
- Trasande L, Cronk C, Durkin M, Weiss M, Schoeller DA, Gall EA, et al. 2009. Environment and obesity in the National Children's Study. *Environ Health Perspect* 117:159–166.
- Vandentorren S, Bois C, Pirus C, Sarter H, Salines G, Leridon H. 2009. Rationales, design and recruitment for the Elfe longitudinal study. *BMC Pediatr* 9:58; doi:10.1186/1471-2431-9-58 [Online 21 September 2009].
- van Eijsden M, Vrijkotte TG, Gemke RJ, van der Wal MF. 2010. Cohort profile: the Amsterdam Born Children and their Development (ABCD) Study. *Int J Epidemiol* 40(5):1176–1186.
- Vardavas CI, Patelarou E, Chatzi L, Roumeliotaki T, Sarri K, Murphy S, et al. 2010. Factors associated with active smoking, quitting, and secondhand smoke exposure among pregnant women in Greece. *J Epidemiol* 20:355–362.
- WHO (World Health Organization). 2010. WHO Research Agenda for Radiofrequency Fields. Available: <http://www.who.int/peh-emf/research/agenda/en/index.html> [accessed 6 July 2011].
- Wickman M, Kull I, Pershagen G, Nordvall SL. 2002. The BAMSE project: presentation of a prospective longitudinal birth cohort study. *Pediatr Allergy Immunol* 13(suppl 15):11–13.
- Wigle DT, Arbuckle TE, Walker M, Wade MG, Liu S, Krewski D. 2007. Environmental hazards: evidence for effects on child health. *J Toxicol Environ Health B Crit Rev* 10:3–39.
- Wilhelm M, Wittsiepe J, Lemm F, Ranft U, Kramer U, Furst P, et al. 2008. The Duisburg birth cohort study: influence of the prenatal exposure to PCDD/Fs and dioxin-like PCBs on thyroid hormone status in newborns and neurodevelopment of infants until the age of 24 months. *Mutat Res* 659:83–92.
- Willett WC, Blot WJ, Colditz GA, Folsom AR, Henderson BE, Stampfer MJ. 2007. Merging and emerging cohorts: not worth the wait. *Nature* 445:257–258.
- Woodruff TJ, Parker JD, Adams K, Bell ML, Gehring U, Glinianaia S, et al. 2010. International Collaboration on Air Pollution and Pregnancy Outcomes (ICAPPO). *Int J Environ Res Public Health* 7:2638–2652.
- Zirngibl A, Franke K, Gehring U, von Berg A, Berdel D, Bauer CP, et al. 2002. Exposure to pets and atopic dermatitis during the first two years of life. A cohort study. *Pediatr Allergy Immunol* 13:394–401.