# Peer

## Reliability of the velocity achieved during the last repetition of sets to failure and its association with the velocity of the 1-repetition maximum

Amador García-Ramos<sup>1,2</sup>, Danica Janicijevic<sup>3</sup>, Jorge M. González-Hernández<sup>4</sup>, Justin W.L. Keogh<sup>5,6,7,8</sup> and Jonathon Weakley<sup>9,10</sup>

- <sup>1</sup> Department of Sports Sciences and Physical Conditioning, Universidad Católica de la Santísima Concepción, Concepción, Chile
- <sup>2</sup> Department of Physical Education and Sport, Faculty of Sport Sciences, University of Granada, Granada, Spain
- <sup>3</sup> Faculty of Sport and Physical Education, The Research Centre, University of Belgrade, Belgrade, Serbia
- <sup>4</sup> Faculty of Health Science, Universidad Europea de Canarias, La Orotava, Tenerife, Spain
- <sup>5</sup> Faculty of Health Sciences and Medicine, Bond University, Gold Coast, QLD, Australia
- <sup>6</sup> Sports Performance Research Centre New Zealand, AUT University, Auckland, New Zealand
- <sup>7</sup> Cluster for Health Improvement, Faculty of Science, Health, Education and Engineering, University of the Sunshine Coast, Gold Coast, QLD, Australia
- <sup>8</sup> Kasturba Medical College, Manipal Academy of Higher Education, Manipal, Karnataka, India
- <sup>9</sup> School of Behavioural and Health Sciences, Australian Campus University, Brisbane, QLD, Australia
- <sup>10</sup> Carnegie Applied Rugby Research (CARR) centre, Institute for Sport, Physical Activity and Leisure, Leeds Beckett University, Leeds, UK

## ABSTRACT

**Background:** This study aimed to determine the reliability of the velocity achieved during the last repetition of sets to failure ( $V_{\text{last}}$ ) and the association of  $V_{\text{last}}$  with the velocity of the 1-repetition maximum ( $V_{1\text{RM}}$ ) during the paused and touch-and-go bench press (BP) exercises performed in a Smith machine.

**Methods:** A total of 96 healthy men participated in this study that consisted of two testing sessions. A single BP variant (paused BP or touch-and-go BP) was evaluated on each session in a randomized order. Each session consisted of an incremental loading test until reaching the 1RM, followed by two sets of repetitions to failure against a load ranging from 75% to 90% of 1RM.

**Results:** The reliability of  $V_{\text{last}}$  was unacceptable for both BP variants (CV > 18.3%, ICC < 0.60). The correlations between  $V_{1\text{RM}}$  and  $V_{\text{last}}$  were small for the paused BP (r = 0.18) and moderate for the touch-and-go BP (r = 0.37).

**Conclusions:** Although these results suggest that  $V_{\text{last}}$  could be a better indicator of the minimal velocity threshold than  $V_{1\text{RM}}$ , the low reliability of  $V_{\text{last}}$  and the similar values of  $V_{\text{last}}$  for both BP variants suggest that a standard  $V_{1\text{RM}}$  should be used to estimate the 1RM from the individualized load-velocity relationship.

#### Subjects Kinesiology, Public Health

**Keywords** Bench press, Linear position transducer, Minimal velocity threshold, Strength testing, Velocity-based training

Submitted 3 December 2019 Accepted 17 February 2020 Published 11 March 2020

Corresponding author Amador García-Ramos, amagr@ugr.es

Academic editor Scotty Butcher

Additional Information and Declarations can be found on page 11

DOI 10.7717/peerj.8760

Copyright 2020 García-Ramos et al.

Distributed under Creative Commons CC-BY 4.0

#### **OPEN ACCESS**

How to cite this article García-Ramos A, Janicijevic D, González-Hernández JM, Keogh JWL, Weakley J. 2020. Reliability of the velocity achieved during the last repetition of sets to failure and its association with the velocity of the 1-repetition maximum. PeerJ 8:e8760 DOI 10.7717/peerJ.8760

## INTRODUCTION

Movement velocity has been proposed as an accurate variable for estimating the 1-repetition maximum (1RM) during a number of resistance training exercises (*García-Ramos & Jaric, 2018; McBurnie et al., 2019*). Early studies proposed generalized load-velocity (L-V) relationship equations to estimate the percentage of 1RM from the velocity value recorded against a submaximal load lifted with maximal effort (*González-Badillo & Sánchez-Medina, 2010*). The basic premise of generalized L-V relationship equations is that a given velocity represents the same percentage of 1RM for all individuals. However, this premise has been refuted in subsequent studies which have shown a greater accuracy in the estimation of the percentage of 1RM using individualized L-V relationships (*García-Ramos & Jaric, 2018; García-Ramos et al., 2018a; McBurnie et al., 2019*). This is also demonstrated by the between-subject variability in the velocity associated with a given percentage of 1RM being higher than the within-subject variability (*Pestaña-Melero et al., 2018*). On the basis of these results, an increasing number of studies have recently been conducted to refine the testing procedure of the individualized L-V relationship.

Assessment of individualized L-V relationships requires the recording of movement velocity against at least two submaximal loads and subsequently, the 1RM can be estimated through a linear regression as the load associated with the velocity of the 1RM (V<sub>1RM</sub> or minimal velocity threshold) (García-Ramos & Jaric, 2018; García-Ramos et al., 2018a; McBurnie et al., 2019). However, one of the challenges associated with the use of individualized L-V relationships for predicting the 1RM is how to select the minimal velocity threshold. Previous studies have selected the minimal velocity threshold as either the individualized V<sub>1RM</sub> (Banyard, Nosaka & Haff, 2017; Ruf, Chery & Taylor, 2018) or a general  $V_{1RM}$  for all subjects (*García-Ramos et al., 2018a, 2019a*). The assessment of the individualized  $V_{1RM}$  is associated with at least two problems: (I) the individual is required to perform a lift against the 1RM load and (II) the individual  $V_{1RM}$  has been demonstrated to be an unreliable metric for a number of exercises such as the back squat (coefficient of variation (CV) = 22.5%, intraclass correlation coefficient (ICC) = 0.42) (Banyard, Nosaka & Haff, 2017), deadlift (CV = 15.7%, ICC = 0.63) (Ruf, Chery & *Taylor*, 2018), Smith machine bench press (BP) (CV = 13.9-15.7%, ICC = 0.54-0.64) (Pestaña-Melero et al., 2018), or bench pull (CV = 6.36%, ICC = 0.18) (García-Ramos et al., 2019b). Therefore, it would be of interest to investigate whether the minimal velocity threshold (i.e., velocity value used to estimate the 1RM from the individualized L-V relationship) can be obtained with a higher reliability using other approaches that do not require the individual to perform a lift against the 1RM load.

An alternative approach for determining the minimal velocity threshold could be the assessment of the velocity of the last repetition performed during a set to failure ( $V_{last}$ ) (*Lake et al., 2017*). This approach is supported by the results of *Izquierdo et al. (2006)* who did not find significant differences between the individual  $V_{1RM}$  and the  $V_{last}$  collected against four submaximal loads (60%, 65%, 70% and 75% of 1RM) during the BP and parallel back squat exercises performed in a Smith machine. A recent review has suggested

that the accuracy of the individualized L-V relationship for predicting the 1RM is higher for upper-body (e.g., BP or bench pull) compared to lower-body exercises (e.g., squat or deadlift), while the BP is the exercise most explored in velocity-based training research (*McBurnie et al., 2019*). The two main variants of the BP exercise examined in the scientific literature are the paused BP (a pause is introduced between the eccentric and concentric phases) and the touch-and-go BP (the concentric phase is performed immediately after the eccentric phase) (*García-Ramos et al., 2018b*; *Pallarés et al., 2014*; *Pérez-Castilla et al., 2018*). *García-Ramos et al. (2018a)* revealed a comparable  $V_{1RM}$  for the paused BP (0.168 ± 0.026 m/s) and touch-and-go BP (0.178 ± 0.030 m/s) (P = 0.232) performed in a Smith machine, while the  $V_{1RM}$  was poorly related between the BP variants (r = -0.11, P = 0.554). However, no study has examined the reliability of  $V_{last}$  or the association between  $V_{last}$  and  $V_{1RM}$ . Therefore, a comprehensive examination of the behavior of  $V_{last}$  during the BP exercise is needed to clarify whether  $V_{last}$  may provide useful information for increasing the accuracy in the estimation of the 1RM through the individualized L-V relationship.

To address the aforementioned gaps in the literature, the main aim of the present study was to assess the reliability of  $V_{\text{last}}$  and the association between  $V_{\text{last}}$  and  $V_{1\text{RM}}$  during the paused and touch-and-go BP exercises performed in a Smith machine. In addition, we aimed to determine the effect of the number of repetitions performed to failure on  $V_{\text{last}}$  and the effect of 1RM strength on  $V_{1\text{RM}}$  and  $V_{\text{last}}$ . Our main hypothesis was that, regardless of the BP variant,  $V_{\text{last}}$  would present a low level of reliability and it would be poorly correlated with  $V_{1\text{RM}}$ . We also hypothesized that no significant correlations would be observed between  $V_{\text{last}}$  and the number of repetitions performed to failure, while the 1RM strength would be negatively associated with  $V_{1\text{RM}}$  and  $V_{\text{last}}$ .

## **MATERIALS AND METHODS**

## **Participants**

Ninety-six healthy men volunteered to participate in this study (age =  $20.8 \pm 3.4$  years (range = 18-38 years); body height =  $1.73 \pm 0.06$  m; body mass =  $75.3 \pm 15.7$  kg; paused BP 1RM =  $62.3 \pm 17.6$  kg; touch-and-go BP 1RM =  $66.5 \pm 18.4$  kg; resistance training experience =  $1.3 \pm 2.4$  years (range = 0-10 years)). Eighty-six participants completed both testing sessions, seven participants only performed the paused BP and three participants only performed the touch-and-go BP. A total of 10 participants only performed one BP variant because they reported not to be interested in attending a second testing session. All participants without resistance training experience were first year sport science students and they were familiarized with the BP exercise during several sessions before the onset of the study. Prior to study initiation, participants were informed of the study procedures and provided written informed consent. Additionally, participants were instructed to avoid any strenuous exercise for the 48 h before each testing session. The study protocol adhered to the tenets of the University of Granada (491/CEIH/ 2018).

#### Study design

A randomized crossover design was used to comprehensively examine the relationship between  $V_{1RM}$  and  $V_{last}$  during the BP exercise performed in a Smith machine. Participants came to the laboratory on two occasions separated by 72-96 h. A single BP variant (paused BP or touch-and-go BP) was evaluated on each session in a randomized order. Each session consisted of an incremental loading test until reaching the 1RM, followed by two sets of repetitions to failure against a load ranging from the 75% of 1RM to the 90% of 1RM. The same participant performed the two sets against the same load for reliability purposes, while the prescribed relative load (% of 1RM) differed between participants to explore the effect of the number of repetitions performed on  $V_{last}$ . Participants were instructed to perform all repetitions at the maximum possible velocity and the mean concentric velocity (MCV, average velocity value from the start of the concentric phase until the velocity of the barbell was  $0 \text{ m} \cdot \text{s}^{-1}$ ) of the barbell was recorded with a linear velocity transducer (T-Force System; Ergotech, Murcia, Spain). The two sessions for the same participant were held at the same time of the day  $(\pm 1 h)$  to minimize the influence of the circadian rhythm on physical performance.

#### Procedures

Both testing sessions began with a standardized warm-up which consisted of 5 min of jogging, dynamic stretching, arm and shoulder mobilization and one set of 10 repetitions with an external load of 20 kg (mass of the unloaded Smith machine barbell) in the tested BP variant. The initial load of the incremental loading test was 20 kg and it was increased in 10 kg until the MCV of the barbell was lower than 0.50 ms<sup>-1</sup>. From that moment, the load was increased in increments of 1–5 kg until the 1RM load was achieved. During the incremental loading test, two repetitions were performed when the MCV was higher than 0.50 ms<sup>-1</sup> and only one repetition when the MCV was lower than 0.50 ms<sup>-1</sup>. The rest between sets was 3 min when the MCV was higher than 0.50 ms<sup>-1</sup> and 5 min when the MCV was lower than 0.50 ms<sup>-1</sup>.

Ten minutes after the 1RM assessment, participants performed two sets of repetitions to failure with a load ranging between 75% and 90% of the previously determined 1RM. The two sets were separated by 10 min. Participants performed the two sets with the same load for reliability purposes, but the relative load differed between participants to explore the effect of the number of repetitions performed on  $V_{\text{last}}$ . Participants were instructed to perform all repetitions at the maximum possible velocity. Two trained spotters were present during the test for safety reasons and to encourage participants to lift the maximum possible load during the 1RM assessment and to perform the maximum possible number of repetitions during the sets of repetitions to failure. The BP was performed in a Smith machine (Ffittech, Taiwan, China), while a linear velocity transducer, which sampled the velocity of the barbell at 1,000 Hz, was used to collect the MCV of all repetitions. Note that the MCV has been reported to be the most accurate velocity variable for determining the load-velocity relationship during the BP exercise

(*García-Ramos et al., 2018c*). The two BP variants evaluated in the present study are described below:

*Paused BP*: Participants initiated the task holding the barbell with their arms fully extended. They lowered the barbell at a self-selected velocity until the barbell made contact with their chest, waited with the barbell on the chest for 2 s and on the word "Go!" performed a purely concentric action at maximum possible velocity until their arms were fully extended.

*Touch-and-go BP*: Participants initiated the task holding the barbell with their arms fully extended. They were instructed to lower the barbell until it touched the chest and then immediately perform the concentric phase at the maximum possible velocity.

## Statistical analysis

Descriptive data are presented as means and standard deviations. The normal distribution of the data was confirmed by the Shapiro–Wilk test (P > 0.05). The reliability of  $V_{\text{last}}$ was assessed through the standard error of measurement (SEM) the CV and the ICC (model 3.1) with the corresponding 95% confidence interval. Acceptable reliability was determined as a CV < 10% and a ICC > 0.70 (*Cormack et al.*, 2008). The association between the variables was quantified by the Pearson's correlation coefficient (r) and the following scale was used to quantify the magnitude of the r coefficient: trivial (0.00–0.09) small (0.10-0.29) moderate (0.30-0.49) large (0.50-0.69) very large (0.70-0.89) nearly perfect (0.90-0.99) and perfect (1.00) (Hopkins et al., 2009). Paired sample's t tests, the Cohen's d effect size (ES) and Bland–Altman plots were used to compare the magnitude of the variables. The following scale was used to interpret the magnitude of the ES: trivial (<0.20) small (0.20–0.59) moderate (0.60–1.19) large (1.20–1.99) and very large (≥2.00) (Hopkins et al., 2009). The reliability assessment was performed by means of a custom spreadsheet (*Hopkins*, 2000), while all other statistical analyses were performed using the software SPSS version 22.0 (SPSS, Chicago, IL, USA). Statistical significance was set at an alpha level of 0.05.

## RESULTS

The reliability of  $V_{\text{last}}$  was unacceptable for both the paused BP (CV = 18.3%, ICC = 0.58) and the touch-and-go BP (CV = 20.4%, ICC = 0.48) (Table 1). The  $V_{1\text{RM}}$  was significantly higher than  $V_{\text{last}}$  during the paused BP (P = 0.028, ES = 0.30) and the touch-and-go BP (P = 0.020, ES = 0.29), while the magnitude of the correlations between  $V_{1\text{RM}}$  and  $V_{\text{last}}$ was small for the paused BP (r = 0.18, P = 0.083) and moderate for the touch-and-go BP (r = 0.37, P < 0.001) (Fig. 1). No significant differences between the BP variants were observed for  $V_{1\text{RM}}$  or  $V_{\text{last}}$  (P > 0.80), while the correlations between the BP variants was trivial for  $V_{1\text{RM}}$  or  $V_{\text{last}}$  (P > 0.80), while the correlations between the BP variants was trivial for  $V_{1\text{RM}}$  (r = -0.01, P = 0.937) and moderate for  $V_{\text{last}}$  (r = 0.46, P < 0.001) (Fig. 2). Bland–Altman plots revealed comparable systematic bias (0.015–0.016 m·s<sup>-1</sup>) and random errors (0.054–0.057 m·s<sup>-1</sup>) between  $V_{1\text{RM}}$  and  $V_{\text{last}}$  for both BP variants (Fig. 3). A small and inverse relationship was observed between  $V_{\text{last}}$  and the number of repetitions performed to failure (r = -0.23, P < 0.001) and the 1RM load (r = -0.17, P = 0.001), while moderate correlations were observed between  $V_{1\text{RM}}$  and the 1RM load

Table 1      Reliability of mean concentric velocity values achieved during the last repetition of sets to
failure during the paused and touch-and-go bench press exercises.

Exercise	Set 1 (m·s <sup>−1</sup> )	Set 2 (m·s <sup>−1</sup> )	Þ	CV (95% CI)	ICC (95% CI)	SEM (m·s <sup>−1</sup> )
Paused BP $(n = 93)$	0.161 (0.045)	0.157 (0.045)	0.303	18.3% [16.0%, 21.4%]	0.58 [0.43, 0.70]	0.029
Touch-and-go BP $(n = 89)$	0.162 (0.046)	0.156 (0.043)	0.266	20.4% [17.8%, 24.0%]	0.48 [0.30, 0.62]	0.032

Note:

BP, bench press; p, p-value; CV, coefficient of variation; ICC, intraclass correlation coefficient; SEM, standard error of measurement; 95% CI, 95% confidence interval.



Figure 1 Association and comparison between the velocity of the 1-repetition maximum and the velocity achieved during the last repetition of sets to failure. Association (A and C) and comparison (B and D) between the velocity of the 1-repetition maximum ( $V_{1RM}$ ) and the velocity achieved during the last repetition of sets to failure ( $V_{last}$ ) during the paused (A and B) and touch-and-go (C and D) bench press (BP) exercises. *r*, Pearson's correlation coefficient; *p*, *p*-value obtained from paired sample's *t* tests; ES, Cohen's *d* effect size (( $V_{1RM}$  mean- $V_{last}$  mean)/SD both). Full-size  $\square$  DOI: 10.7717/peerj.8760/fig-1

(r = -0.33, P < 0.001) (Fig. 4). Participants performed the sets of repetitions to failure against an average load of 82.1 ± 3.7% of 1RM and completed 6.9 ± 2.3 repetitions.

## DISCUSSION

This study is the first to investigate the reliability of  $V_{\text{last}}$  during the paused and touch-andgo BP exercises and explore the association between  $V_{1\text{RM}}$  and  $V_{\text{last}}$  within and between



Figure 2 Association and comparison between the paused bench press and the touch-and-go bench press for the velocity of the 1-repetition maximum and the velocity achieved during the last repetition of sets to failure. Association (A and C) and comparison (B and D) between the paused bench press (BP) and the touch-and-go BP for the velocity of the 1-repetition maximum ( $V_{1RM}$ ; A and B) and the velocity achieved during the last repetition of sets to failure ( $V_{last}$ ; C and D). *r*, Pearson's correlation coefficient; *p*, *p*-value obtained from paired sample's *t* tests; ES, Cohen's *d* effect size ((paused BP mean-touch-and-go BP mean)/SD both). Full-size  $\Box$  DOI: 10.7717/peerj.8760/fig-2

each BP variant. Additionally, this study determined the effect of the number of repetitions performed to failure on  $V_{\text{last}}$  and the effect of 1RM strength on  $V_{1\text{RM}}$  and  $V_{\text{last}}$ . The main findings of the study revealed (I) low reliability for  $V_{\text{last}}$  (II)  $V_{\text{last}}$  was significantly lower than  $V_{1\text{RM}}$  (III) no significant differences between the BP variants for  $V_{1\text{RM}}$  or  $V_{\text{last}}$  (IV) larger associations between the BP variants for  $V_{1\text{RM}}$  and (V) a negative, albeit weak, association of  $V_{\text{last}}$  with the number of repetitions performed to failure and the 1RM strength. These results suggest that  $V_{\text{last}}$  could be a better indicator of the minimal velocity threshold during the BP exercise than  $V_{1\text{RM}}$  because  $V_{\text{last}}$  was lower than  $V_{1\text{RM}}$  and it demonstrated a stronger relationship between the BP variants. However, the low reliability of  $V_{\text{last}}$ , moderate association between the BP variants for  $V_{\text{last}}$  and the lack of differences between the BP variants for the  $V_{1\text{RM}}$  or  $V_{\text{last}}$  suggest that a general  $V_{1\text{RM}}$  could be more appropriate.

Previous studies have found that the magnitude of  $V_{\text{last}}$  is not affected by the load applied during sets of repetitions to failure performed with the BP and back squat exercises (*Izquierdo et al., 2006*). However, no previous study had examined the reliability of  $V_{\text{last}}$  in



Figure 3 Differences between the velocity of the 1-repetition maximum and the velocity achieved during the last repetition of sets to failure during the paused bench press (BP) and the touch-andgo BP. Bland–Altman plots showing the differences between the velocity of the 1-repetition maximum ( $V_{1RM}$ ) and the velocity achieved during the last repetition of sets to failure ( $V_{last}$ ) during the paused bench press (BP) (A) and the touch-and-go BP (B). Each plot depicts the systematic bias and 95% limits of agreement (± 1.96 SD; dashed lines), along with the regression line (solid line). The systematic bias ± random error together with the strength of the relationship ( $r^2$ ) are depicted in each plot. Full-size DOI: 10.7717/peerj.8760/fig-3

any resistance training exercise. Our results confirmed that  $V_{\text{last}}$  is an unreliable metric for the two variants of the BP exercise examined in the present study. The reliability of  $V_{\text{last}}$ (CV = 18.3% and 20.4% for the paused and touch-and-go BP, respectively) was somewhat comparable to the reliability of the MCV associated with the 1RM load reported by **Peer** 



Figure 4 Relationship between the number of repetitions performed to failure and velocity of the last repetition performed during sets of repetitions to failure  $(V_{last})$ , 1-repetition maximum and  $V_{last}$  and 1-repetition maximum and velocity. Relationship between the number of repetitions performed to failure and  $V_{last}$  (A) 1-repetition maximum and  $V_{last}$  (B) and 1-repetition maximum and  $V_{1RM}$  (C).  $V_{last}$ , velocity of the last repetition performed during sets of repetitions to failure;  $V_{1RM}$ , velocity achieved during the 1-repetition maximum trial. The regression equation and the Pearson's correlation coefficient (r) are depicted. Full-size DOI: 10.7717/peerj.8760/fig-4

*Pestaña-Melero et al. (2018)* (CV =13.9% and 15.7% for the paused and touch-and-go BP, respectively). Therefore, it seems that neither the  $V_{1RM}$  nor the  $V_{last}$  should be used on an individual basis since the within-subject variability of these variables is not meaningfully lower than the variability existing between subjects (*Pestaña-Melero et al., 2018*).

The generally low correlations between  $V_{1RM}$  and  $V_{last}$  for both BP variants ( $r \le 0.37$ ) indicates that the value of one variable cannot be inferred from the other. Similarly, as it was previously shown by García-Ramos et al. (2018a) no significant correlations were observed between the two BP variants for  $V_{1RM}$  (r = -0.01). Only the value of  $V_{last}$  was moderately correlated between the two BP variants (r = 0.46), revealing that participants with a higher  $V_{\text{last}}$  in one BP variant also tended to have a higher  $V_{\text{last}}$  in the other BP variant. These results provide additional support for the use of the same minimal velocity threshold (i.e.,  $V_{1RM}$ ) for all participants when predicting the 1RM from the individualized L-V relationship during the BP exercise performed in a Smith machine. The lack of differences between the BP variants for the magnitudes of  $V_{1\text{BM}}$  and  $V_{\text{last}}$  simplifies this approach because the same minimal velocity threshold could be used for both BP variants. According to the results of this and previous studies, the minimal velocity threshold should be set at 0.17 m·s<sup>-1</sup> (García-Ramos et al., 2018a; González-Badillo & Sánchez-Medina, 2010; Sánchez-Medina et al., 2014). However, Helms et al. (2017) reported in 15 powerlifters (12 men and 3 women) a lower  $V_{1RM}$  during the free-weight BP exercise that was performed using a "press" command to simulate a powerlifting competition. Therefore, additional research should be conducted with different variants of the free-weight BP and more trained populations to elucidate whether the minimal velocity threshold recommended in the present study can also be applicable to these conditions.

In the present study we also explored the influence of 1RM strength on the values of  $V_{1RM}$  and  $V_{last}$ . Based on the lower  $V_{1RM}$  reported by Helms et al. (2017) in powerlifters during the free-weight BP (0.10  $\pm$  0.04 m·s<sup>-1</sup>) compared to the V<sub>1RM</sub> reported by *Loturco et al. (2017)* in rugby players and combat athletes (0.17 m $\cdot$ s<sup>-1</sup>), we hypothesized that the 1RM strength would be negatively associated with  $V_{1RM}$  and  $V_{last}$ . This hypothesis was somewhat supported with a significant, albeit weak, negative correlation between the 1RM load and  $V_{1RM}$  (r = -0.33) and  $V_{last}$  (r = -0.17). This result is also in line with the findings of Ormsbee et al. (2019) who reported a slower  $V_{1RM}$  for experienced benchers  $(0.14 \pm 0.04 \text{ m} \cdot \text{s}^{-1})$  compared to novice benchers  $(0.20 \pm 0.05 \text{ m} \cdot \text{s}^{-1})$ . Therefore, although based on these results a slightly lower minimal velocity threshold could be obtained by stronger subjects, the low reliability and low correlations found in the present study for  $V_{\rm IRM}$  and  $V_{\rm last}$  suggest that it may not be necessary to modify the minimal velocity threshold based on the 1RM strength values when the BP is performed in a Smith machine. Finally, we also explored the influence of the number of repetitions performed during the sets to failure on  $V_{\text{last}}$  and we observed a relatively weak negative association (r = -0.23), indicating that a slightly lower  $V_{\text{last}}$  can be obtained when higher number of repetitions are performed. Therefore, in the case that coaches are interested in determining  $V_{\text{last}}$  we recommended that  $V_{\text{last}}$  is derived from a set to failure of at least five repetitions.

A limitation of the present study is that the 1RM was directly assessed only once for each BP variant and therefore, it was not possible to compare the reliability between  $V_{1RM}$ and  $V_{\text{last.}}$  However, the lower correlations between the BP variants observed in this study for  $V_{1RM}$  compared to  $V_{last}$  suggest that the  $V_{1RM}$  could also be less reliable than  $V_{last}$ . This would corroborate the results of previous studies that have reported a poor reliability for the  $V_{1RM}$  during the BP (*Pestaña-Melero et al., 2018*) and other resistance training exercises such as the back squat (Banyard, Nosaka & Haff, 2017), deadlift (Ruf, Chery & Taylor, 2018), or bench pull (García-Ramos et al., 2019b). Another limitation is that the BP was performed in a Smith machine, while the vast majority of athletes use the free-weight BP during their resistance training programs. In the current study, the Smith machine was used to simplify the testing procedure and to improve the reproducibility of velocity readings. Thus, it is plausible that the reliability of  $V_{\text{last}}$  and  $V_{\text{IRM}}$  may be even lower if the BP is performed with free-weights due to the additional control required and horizontal movements characterizing the free weight BP. Finally, the sample of this study was composed of males with a mean of 1.3 years of experience with the BP exercise and therefore, it is possible that the accuracy of  $V_{1RM}$  and  $V_{last}$  could increase when testing participants with more experience. Future studies should compare the magnitude of  $V_{1RM}$  and  $V_{last}$  as well as the reliability of these variables between individuals with different training backgrounds to shed light on this topic.

## **CONCLUSIONS**

The reliability of  $V_{\text{last}}$  was below the threshold of acceptable reliability for both BP variants.  $V_{\text{last}}$  was always significantly lower than  $V_{1\text{RM}}$ , while no significant differences between the BP variants were observed for  $V_{1\text{RM}}$  or  $V_{\text{last}}$ .  $V_{\text{last}}$  was more correlated between the BP variants than  $V_{1\text{RM}}$ . The correlations between  $V_{1\text{RM}}$  and  $V_{\text{last}}$  ranged from small (paused BP) to moderate (touch-and-go BP). An inverse, but generally weak, association was observed between  $V_{\text{last}}$  and the number of repetitions performed to failure and the 1RM load, as well as between the  $V_{1\text{RM}}$  and the 1RM load. Therefore, even though  $V_{\text{last}}$  could be a more appropriate indicator of the minimal velocity threshold than  $V_{1\text{RM}}$ , our results (i.e., low reliability, lack of differences in magnitude and only moderate association between the BP variants) also suggest that a general  $V_{1\text{RM}}$  could be more appropriate to estimate the 1RM during the BP exercise performed in a Smith machine.

## ACKNOWLEDGEMENTS

We would like to thank all the participants who volunteered to participate in this study.

## **ADDITIONAL INFORMATION AND DECLARATIONS**

## Funding

This research was funded by the Universidad Católica de la Santísima Concepción (DINREG 09/2019). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

## **Grant Disclosures**

The following grant information was disclosed by the authors: Universidad Católica de la Santísima Concepción: DINREG 09/2019.

#### **Competing Interests**

Amador García-Ramos and Justin W.L. Keogh are Academic Editors for PeerJ.

#### **Author Contributions**

- Amador García-Ramos conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Danica Janicijevic performed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Jorge M. González-Hernández performed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Justin W.L. Keogh conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Jonathon Weakley analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.

#### **Human Ethics**

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

The Institutional Review Board of the University of Granada approved this study (491/CEIH/2018).

#### **Data Availability**

The following information was supplied regarding data availability:

The data is available in the Supplemental Files.

#### **Supplemental Information**

Supplemental information for this article can be found online at http://dx.doi.org/10.7717/ peerj.8760#supplemental-information.

## REFERENCES

- Banyard HG, Nosaka K, Haff GG. 2017. Reliability and validity of the load-velocity relationship to predict the 1RM back squat. *Journal of Strength and Conditioning Research* 31:1897–1904 DOI 10.1519/JSC.000000000001657.
- **Cormack SJ, Newton RU, McGuigan MR, Doyle TLA. 2008.** Reliability of measures obtained during single and repeated countermovement jumps. *International Journal of Sports Physiology and Performance* **3**:131–144 DOI 10.1123/ijspp.3.2.131.
- García-Ramos A, Barboza-Gonzalez P, Ulloa-Diaz D, Rodriguez-Perea A, Martinez-Garcia D, Guede-Rojas F, Hinojosa-Riveros H, Chirosa-Rios LJ, Cuevas-Aburto J, Janicijevic D, Weakley J. 2019a. Reliability and validity of different methods of estimating the one-repetition maximum during the free-weight prone bench pull exercise. *Journal of Sports Sciences* 37:2205–2212 DOI 10.1080/02640414.2019.1626071.

- García-Ramos A, Haff GG, Pestaña-Melero FL, Pérez-Castilla A, Rojas FJ, Balsalobre-Fernández C, Jaric S. 2018a. Feasibility of the 2-point method for determining the 1-repetition maximum in the bench press exercise. *International Journal of Sports Physiology and Performance* 13:474–481 DOI 10.1123/ijspp.2017-0374.
- **García-Ramos A, Jaric S. 2018.** Two-point method: a quick and fatigue-free procedure for assessment of muscle mechanical capacities and the one-repetition maximum. *Strength and Conditioning Journal* **40**:54–66 DOI 10.1519/SSC.00000000000359.
- García-Ramos A, Pestaña-Melero FL, Pérez-Castilla A, Rojas FJ, Haff GG. 2018b. Differences in the load-velocity profile between 4 bench press variants. *International Journal of Sports Physiology and Performance* 13:326–331 DOI 10.1123/ijspp.2017-0158.
- García-Ramos A, Pestaña-Melero FL, Pérez-Castilla A, Rojas FJ, Haff GG. 2018c. Mean velocity vs. mean propulsive velocity vs. peak velocity: which variable determines bench press relative load with higher reliability? *Journal of Strength and Conditioning Research* 32:1273–1279 DOI 10.1519/JSC.000000000001998.
- García-Ramos A, Ulloa-Díaz D, Barboza-González P, Rodríguez-Perea Á, Martínez-García D, Quidel-Catrilelbún M, Guede-Rojas F, Cuevas-Aburto J, Janicijevic D, Weakley J. 2019b. Assessment of the load-velocity profile in the free-weight prone bench pull exercise through different velocity variables and regression models. *PLOS ONE* 14:e0212085–e0212085 DOI 10.1371/journal.pone.0212085.
- **González-Badillo J, Sánchez-Medina L. 2010.** Movement velocity as a measure of loading intensity in resistance training. *International Journal of Sports Medicine* **31**:347–352 DOI 10.1055/s-0030-1248333.
- Helms ER, Storey A, Cross MR, Brown SR, Lenetsky S, Ramsay H, Dillen C, Zourdos MC. 2017.
  RPE and velocity relationships for the back squat, bench press, and deadlift in powerlifters.
  *Journal of Strength and Conditioning Research* 31:292–297
  DOI 10.1519/JSC.000000000001517.
- Hopkins W. 2000. Calculations for reliability (Excel spreedsheet). Available at http://www.sportsci. org/resource/stats/relycalc.html%7B#%7Dexcel.
- Hopkins WG, Marshall SW, Batterham AM, Hanin J. 2009. Progressive statistics for studies in sports medicine and exercise science. *Medicine and Science in Sports and Exercise* 41:3–13 DOI 10.1249/MSS.0b013e31818cb278.
- Izquierdo M, González-Badillo JJ, Häkkinen K, Ibáñez J, Kraemer WJ, Altadill A, Eslava J, Gorostiaga EM. 2006. Effect of loading on unintentional lifting velocity declines during single sets of repetitions to failure during upper and lower extremity muscle actions. *International Journal of Sports Medicine* 27:718–724 DOI 10.1055/s-2005-872825.
- Lake J, Naworynsky D, Duncan F, Jackson M. 2017. Comparison of different minimal velocity thresholds to establish deadlift one repetition maximum. *Sports* 5:70 DOI 10.3390/sports5030070.
- Loturco I, Kobal R, Moraes JE, Kitamura K, Cal Abad CC, Pereira LA, Nakamura FY. 2017. Predicting the maximum dynamic strength in bench press: the high precision of the bar velocity approach. *Journal of Strength and Conditioning Research* **31**:1127–1131 DOI 10.1519/JSC.000000000001670.
- McBurnie AJ, Allen KP, Garry M, Martin M, Thomas D, Jones PA, Comfort P, McMahon JJ. 2019. The benefits and limitations of predicting one repetition maximum using the load-velocity relationship. *Strength & Conditioning Journal* 41(6):1.
- Ormsbee MJ, Carzoli JP, Klemp A, Allman BR, Zourdos MC, Kim J-S, Panton LB. 2019. Efficacy of the repetitions in reserve-based rating of perceived exertion for the bench press in

experienced and novice benchers. *Journal of Strength and Conditioning Research* **33**:337–345 DOI 10.1519/JSC.000000000001901.

- Pallarés JG, Sánchez-Medina L, Pérez CE, De La Cruz-Sánchez E, Mora-Rodriguez R. 2014. Imposing a pause between the eccentric and concentric phases increases the reliability of isoinertial strength assessments. *Journal of Sports Sciences* 32:1165–1175 DOI 10.1080/02640414.2014.889844.
- Pérez-Castilla A, Comfort P, McMahon J, Pestaña-Melero F, García-Ramos A. 2018. Comparison of the force-, velocity- and power-time curves between the concentric-only and eccentric-concentric bench press exercises. Epub ahead of print 17 January 2018. *Journal of Strength and Conditioning Research*. DOI 10.1519/JSC.00000000002448.
- Pestaña-Melero F, Haff GG, Rojas FJ, Pérez-Castilla A, García-Ramos A. 2018. Reliability of the load-velocity relationship obtained through linear and polynomial regression models to predict the one-repetition maximum load. *Journal of Applied Biomechanics* 34:184–190 DOI 10.1123/jab.2017-0266.
- Ruf L, Chery C, Taylor K-L. 2018. Validity and reliability of the load-velocity relationship to predict the one-repetition maximum in deadlift. *Journal of Strength and Conditioning Research* 32:681–689 DOI 10.1519/JSC.00000000002369.
- Sánchez-Medina L, González-Badillo JJ, Pérez CE, Pallarés JG. 2014. Velocity- and power-load relationships of the bench pull vs. bench press exercises. *International Journal of Sports Medicine* 35:209–216 DOI 10.1055/s-0033-1351252.