



OPEN ACCESS

# Mapping the complexities of Relative Energy Deficiency in Sport (REDs): development of a physiological model by a subgroup of the International Olympic Committee (IOC) Consensus on REDs

Louise M Burke ,<sup>1</sup> Kathryn E Ackerman ,<sup>2,3,4</sup> Ida A Heikura ,<sup>5,6</sup> Anthony C Hackney ,<sup>7</sup> Trent Stellingwerf

<sup>1</sup>Mary MacKillop Institute for Health Research, Australian Catholic University, Melbourne, Victoria, Australia

<sup>2</sup>Wu Tsai Female Athlete Program, Division of Sports Medicine, Boston Children's Hospital, Boston, Massachusetts, USA

<sup>3</sup>Neuroendocrine Unit, Massachusetts General Hospital, Boston, Massachusetts, USA

<sup>4</sup>Harvard Medical School, Boston, Massachusetts, USA

<sup>5</sup>Canadian Sport Institute Pacific, Victoria, British Columbia, Canada

<sup>6</sup>Exercise Science, Physical & Health Education, University of Victoria, Victoria, British Columbia, Canada

<sup>7</sup>Department of Exercise & Sport Science, University of North Carolina, Chapel Hill, North Carolina, USA

## Correspondence to

Dr Louise M Burke, Australian Catholic University, Melbourne, Victoria, Australia; [louise.burke@acu.edu.au](mailto:louise.burke@acu.edu.au)

Accepted 20 August 2023

## ABSTRACT

The 2023 International Olympic Committee (IOC) consensus statement on Relative Energy Deficiency in Sport (REDs) notes that exposure to low energy availability (LEA) exists on a continuum between *adaptable* and *problematic* LEA, with a range of potential effects on both health and performance. However, there is variability in the outcomes of LEA exposure between and among individuals as well as the specific manifestations of REDs. We outline a framework for a 'systems biology' examination of the effect of LEA on individual body systems, with the eventual goal of creating an integrated map of body system interactions. We provide a template that systematically identifies characteristics of LEA exposure (eg, magnitude, duration, origin) and a variety of moderating factors (eg, medical history, diet and training characteristics) that could exacerbate or attenuate the type and severity of impairments to health and performance faced by an individual athlete. The REDs Physiological Model may assist the diagnosis of underlying causes of problems associated with LEA, with a personalised and nuanced treatment plan promoting compliance and treatment efficacy. It could also be used in the strategic prevention of REDs by drawing attention to scenarios of LEA in which impairments of health and performance are most likely, based on knowledge of the characteristics of the LEA exposure or moderating factors that may increase the risk of harmful outcomes. We challenge researchers and practitioners to create a unifying and dynamic physiological model for each body system that can be continuously updated and mapped as knowledge is gained.

## INTRODUCTION

As far back as historical records can be found, humans have been driven to create maps of their understanding of the world around them. The value of maps, then and now, is to provide a visual representation of the nature, magnitude and inter-relationships of the features in our environment. Maps show us the landscape and help us to make better informed plans. As new explorations and technologies emerge, expert cartographers update maps with greater detail and precision compared with earlier, cruder representations. We are

accustomed to the continual evolution of maps. We also accept the coexistence of different visual representations of the same territory, with varying complexity and features according to the end-user's needs.

Such is the 'cartographic' evolution of Relative Energy Deficiency in Sport (REDs).<sup>1</sup> The International Olympic Committee (IOC) first introduced the REDs concept in 2014<sup>2</sup> (updating it in 2018)<sup>3</sup> to describe the range of negative health and performance sequelae of Low Energy Availability (LEA; see Definitions box) in male and female athletes. REDs expanded the Female Athlete Triad model,<sup>4</sup> which focused on the inter-relationship of LEA, menstrual disturbances and compromised bone health in female athletes; this was itself an update on a previous model<sup>5</sup> and was more recently accompanied by the recognition of the proposed Male Athlete Triad.<sup>6,7</sup> New insights, including contributions from the ~180 new REDs-related papers published since 2018, broadened our understanding of the consequences of LEA and informed the 2023 IOC Consensus Statement on REDs,<sup>1</sup> including its updated REDs Health and Performance Conceptual Models (figure 1) and improved, validated Clinical Assessment Tool (REDs CAT2).<sup>8</sup> The 2023 Consensus Statement and its associated papers<sup>8–13</sup> unveil various changes, address some of the challenges of the underpinning science<sup>14</sup> and discuss future research goals. The REDs Conceptual Models for Health and Performance<sup>1</sup> now clarify that LEA is the exposure and REDs is the eventual outcome (see figure 1 and Definitions box). They contain more precise language and recognise more body systems susceptible to LEA-induced impairments.<sup>1</sup> However, it is important to acknowledge a growing body of scientific<sup>15,16</sup> and practical observations of non-uniform outcomes of LEA exposure, noting variations in both the magnitude and duration of LEA between and among individuals, as well as the specific manifestations of REDs. The REDs Conceptual Models now depict the understanding that some scenarios of LEA can be tolerated with modest and/or reversible perturbations to various biological systems or outcomes; we consider this 'adaptable' LEA.<sup>1</sup> Importantly, scientific/clinical attention is drawn to more severe LEA exposure, termed 'problematic LEA'<sup>1</sup> (see Definitions), which

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Br J Sports Med: first published as 10.1136/bjsports-2023-107335 on 26 September 2023. Downloaded from <http://bjsm.bmj.com/> on April 16, 2025 at Australian Catholic University.



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Burke LM, Ackerman KE, Heikura IA, et al. *Br J Sports Med* 2024;**57**:1098–1110.

### WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ The 2023 International Olympic Committee (IOC) Consensus Statement on Relative Energy Deficiency in Sport (REDs) notes that REDs is a syndrome of impaired physiological and/or psychological functioning experienced by female and male athletes that is caused by exposure to problematic (prolonged and/or severe) low energy availability (LEA).
- ⇒ Some scenarios of LEA can be tolerated with modest and/or reversible perturbations to various biological systems or outcomes (*adaptable* LEA). However, scientific/clinical attention is drawn to more severe LEA exposure (*problematic* LEA), which leads to an increase in the range and magnitude of REDs health and performance concerns.
- ⇒ The REDs Conceptual Models provide a simple representation that LEA exposure exists on a continuum between adaptable and problematic, with a range of potential outcomes in both health and performance metrics. The models demonstrate that increased exposure to LEA increases the risk of developing a variety of health and performance impairments that form the REDs syndrome.

### WHAT THIS STUDY ADDS

- ⇒ The proposed physiological model of REDs provides a framework to expand our knowledge, now and into the future, of the effects of LEA on various body systems in the individual athlete and their specific moderating risk factors, resulting in various health and performance impairments.
- ⇒ The REDs Physiological Model takes a 'systems biology' approach to understanding the specific effects of the characteristics of LEA exposure and moderating factors pertinent to the individual athlete, their environment and their lifestyle on REDs outcomes. It notes that energy reallocation during energy scarcity may affect body systems differently, with systems preferred, essential, reducible or expendable in varying ways between and within athletes.
- ⇒ The REDs Physiological Model could be informed by insights gained from adjacent disciplines, which involve the effect of energy scarcity on human metabolism and health; these include evolutionary biology and the treatment of obesity.

### HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- ⇒ The REDs Physiological Model includes information about moderating factors and characteristics of LEA exposure that might assist in the diagnosis of underlying causes of these problems and help to create a targeted treatment plan. While correcting underlying LEA is the cornerstone of REDs therapy, a more personalised and nuanced approach can assist compliance and enhance treatment efficacy.
- ⇒ The REDs Physiological Model could also be used in the strategic prevention of REDs by drawing attention to scenarios of LEA in which impairments of health and performance are most likely, based on knowledge of the characteristics of the LEA exposure or moderating factors that may exacerbate the risk of harmful outcomes. This may help athletes make informed risk versus reward decisions about deliberate scenarios of LEA (eg, increased training load or body composition manipulation) as well as draw attention to scenarios that might inadvertently lead to problems.


leads to an increase in the range and magnitude of REDs health and performance concerns.

Although recognition of the concept of the continuum between adaptable versus problematic LEA adds some nuance to the REDs Conceptual Models and acknowledges the discordance of real-world outcomes versus laboratory manipulations of LEA exposure, it is unable to fully explain the true complexities and various moderators of REDs manifestations. Therefore, while we appreciate that these models provide valuable visualisations of current knowledge and key messages, we recommend that separate templates be constructed to better explore the multiple layers and interactions between and within LEA and various body systems. A more sophisticated exploration of each spoke within the REDs Conceptual Models ([figure 1](#)) could serve as an ever-evolving dashboard of our increasing knowledge of the complexity of the causes and outputs of LEA exposure in athletes. We suggest that such expanded diagrams be termed the REDs Physiological Model (see Definitions). Such models can inform, and be informed by, the insights of researchers studying a given body system as well as others working in fields involving metabolic adaptation.

### METHODS

This paper emanated from discussions within a subgroup of the IOC working group on REDs to assist the future expansion of scientific knowledge and practice regarding the different outcomes of LEA exposure between and within individual athletes. Our author group consisted of experts with significant clinical and research experience pertaining to LEA—sports physiologists, sports dietitians and a sports endocrinologist. We further consulted with individual members of the full IOC working group on REDs as well as experts in LEA from other disciplines. With multidisciplinary expert consensus, we developed a systematic process for examining LEA exposure's effects on different body systems. This approach included characterising the influence of moderating factors of LEA's effects on health and performance. The aims of an expanded and more nuanced model of LEA exposure were identified, with athlete health and well-being preservation at its core. This model further addresses the concern around scenarios of adaptable LEA to focus attention on exposure to problematic LEA. LEA literature was examined to identify characteristics of LEA exposure, in addition to factors specific to the individual athlete (eg, environment and behaviours), which appear to influence the type and severity of health and performance outcomes. We subsequently produced a standardised template to expand our understanding of the context and complexity of the outcomes of LEA exposure on athletes, including the potential for positive, benign and detrimental effects. The proposed model was tested on a body system (haematological system) that has been less explored in the current LEA and REDs literature, involving a wider group of experts from that field to refine the concept.

Herein we present a potential framework to systematically examine the effects of LEA exposure on individual body systems. With additional research, the eventual goal is to develop a 3-D map that can demonstrate the unique features of each athlete, real-life scenarios of LEA and the interactions within and between body systems; [figure 2](#) demonstrates a template for a 'systems biology' approach to that end. Ideally, the development and production of each individual template will highlight knowledge gaps and inspire further research questions to enhance their accuracy. Our intention is to challenge researchers and practitioners to assist in creating a unifying and dynamic physiological



**Figure 1** The REDs 2023 conceptual models for health (A) and performance (B). LEA, low energy availability; REDs, relative energy deficiency in sport.

model for each body system or REDs outcome that can be continuously updated and mapped as knowledge is gained.

### Equity, diversity and inclusion statement

All contributors to this paper have considerable experience in the research or clinical management of REDs, with skills sets including sports medicine and endocrinology, exercise physiology, nutrition and coaching. The authors include three women and two men, from four different countries.

### Recognition of differing responses to LEA exposure: the reproductive system example

The evolution of insights regarding REDs is illustrated by the study of the reproductive system response to LEA. The initial Female Athlete Triad model and REDs Conceptual Models presented simple causal relationships between LEA and functional impairments of individual body systems. Evidence for these relationships was provided by investigations of biomarker perturbations following short-term (~5 to 7 day), strictly-controlled EA manipulations in sedentary females,<sup>17–20</sup> paired with cross-sectional studies of athletes in which biomarkers of LEA were found in association with impairments to these body systems.<sup>21–23</sup> In the case of the reproductive system, concentrations and pulsatility of luteinising hormone (LH) following LEA exposure in females were robustly demonstrated in short-term investigations in laboratory<sup>18 24</sup> and field settings.<sup>25 26</sup> Meanwhile, differences in LH pulsatility were demonstrated between EA replete/eumenorrheic and LEA/amenorrheic cohorts,<sup>27</sup> with an increased prevalence of functional hypothalamic amenorrhea (FHA) and other menstrual disturbances (eg, oligomenorrhea, luteal phase defects, anovulation) in athletes with LEA that was either directly estimated or determined via measurement of LEA surrogates (eg, decreased resting metabolic rate).<sup>28 29</sup>

The results of a laboratory study<sup>24</sup> involving a stepwise approach to EA reduction supported the development of the narrative that LEA could be tolerated until it reached a specific, universal LEA threshold (eg, 30 kcal/kg FFM/day); below which,

LEA would lead to REDs outcomes, such as menstrual dysfunction. However, other data failed to support the experience of a uniform LEA threshold.<sup>30</sup> For example, older premenopausal females ( $28.7 \pm 2.3$  years, 14–18 years postmenarche) failed to show the alterations in LH pulsatility observed in younger premenopausal females ( $20.5 \pm 0.9$  years, 5–8 y post-menarche) in response to 5 days of a laboratory-controlled reduction in EA (45 vs 10 kcal/kg FFM/day).<sup>15</sup> Such a finding is consistent with the Life History theory that available energy is likely allocated to the various body systems with differing priorities at different phases of life.<sup>31 32</sup> However, even within a narrow age group of females (18–24 years), a longer term intervention with graded reductions in energy balance/availability over three menstrual cycles found a relationship between energy restriction and the prevalence of menstrual disturbances but failed to establish a firm LEA threshold<sup>33</sup> or a relationship with the severity of menstrual disturbances.<sup>34</sup> Cross-sectional studies have also reported inconsistent associations between EA and the range of common menstrual disturbances despite a relationship with the most severe outcomes (ie, amenorrhea).<sup>35</sup> Finally, we now know that LEA inhibits the reproductive system of males, as well. Preliminary evidence from studies of males suggests that they seem less susceptible to signs and symptoms of LEA and/or can tolerate a greater restriction of EA before such negative sequelae are observed.<sup>16</sup> Nonetheless, reductions in testosterone, libido and morning erections have been reported in settings of LEA.<sup>36 37</sup>

Of course, even within the narrow scope of LEA and reproductive outcomes in the REDs model, there have been methodological challenges. These include failure to accurately measure menstrual function,<sup>38</sup> inadequate consideration of the differential diagnosis of menstrual disturbances<sup>39</sup> and usage of surrogate markers of LEA.<sup>21 22</sup> There are also limitations and errors associated with direct assessment of EA in free-living populations.<sup>40 41</sup> Yet, it is clear that the outcomes of LEA exposure differ among athletes according to characteristics of the EA reduction as well as other modifying factors.

Definitions from 2023 IOC consensus statement update on REDs<sup>1</sup>**Energy availability**

Energy availability is the dietary energy left over and available for optimal function of body systems after accounting for the energy expended from exercise. Energy availability is expressed as kcal/kg FFM/day and is defined in the scientific literature in the form of a mathematical formula:

$EA \text{ (Energy Availability)} = [EI \text{ (Energy Intake (kcal))} - EEE \text{ (Exercise Energy Expenditure (kcal))}] / FFM \text{ (Fat-Free Mass (kg))} / \text{day}$

**Low Energy Availability**

LEA is any mismatch between dietary energy intake (EI) and energy expended in exercise that leaves the body's total energy needs unmet, i.e., there is inadequate energy to support the functions required by the body to maintain optimal health and performance. LEA occurs as a continuum between scenarios in which effects are benign (adaptable LEA) and others in which there are substantial and potentially long-term impairments of health and performance (problematic LEA).

**Adaptable LEA**

Adaptable LEA is exposure to a reduction in energy availability that is associated with benign effects, including mild and quickly reversible changes in biomarkers of various body systems that signal an adaptive partitioning of energy and the plasticity of human physiology. In some cases, the scenario that underpins the reduction in energy availability (eg, monitored and mindful manipulation of body composition or scheduled period of intensified training or competition) might be associated with acute health or performance benefits (eg, increased relative  $VO_{2max}$ ). Adaptable LEA is typically a short-term experience with minimal (or no) impact on long-term health, well-being or performance. Moderating factors may also alter the expression of outcomes.

**Problematic LEA**

Problematic LEA is exposure to LEA that is associated with greater and potentially persistent disruption of various body systems, often presenting with signs and/or symptoms, and represents a maladaptive response. The characteristics of problematic LEA exposure (eg, duration, magnitude, frequency) may vary according to the body system and the individual. They may be further affected by interaction with moderating factors that can amplify the disruption to health, well-being, and performance.

**REDs Health Conceptual Model and REDs Performance Conceptual Model**

Separate models (see figure 1) which illustrate the relationship between LEA and REDs for a variety of audiences including the athlete's health and performance team and entourage (eg, coaches, parents). The REDs Conceptual Models provide a simple representation that LEA exposure exists on a continuum between adaptable and problematic, with a range of potential outcomes on both health and performance metrics. The models demonstrate that increased exposure to LEA increases the risk of developing a variety of health and performance impairments that form the REDs syndrome.

**REDs Physiological Model**

Evolving model which explores the complexity of LEA exposure on each of the body systems included in the RED Health Conceptual Model. A template (figure 2) could be used to develop a physiological model for each body system, which

## Definitions from 2023 IOC consensus statement update on REDs 1 Continued

notes the effects of characteristics of the LEA exposure as well as a range of moderating factors on the manifestation health and performance impairments. Ideally, the REDs Physiological Model will develop over time to form a 3-D representation of the REDs syndrome in which the interaction and cross-talk between body systems is recognised.


**The physiological model overview: commencing with individual body systems**

Following consultation and feedback from many REDs experts, we have developed a template to guide the standardised development of a physiological model for each body system (figure 2). This template encourages a systematic approach to understanding how LEA may affect each body system, while incorporating other moderating factors that influence the manifestation of system dysfunction. The goal of this activity is to distinguish between scenarios in which LEA exposure may present as an 'adaptable' form and those that are 'problematic' (see Definitions). Indeed, there is little need to focus on brief and benign exposures to LEA. Rather, it is important to prevent and/or treat REDs by addressing circumstances with likely *problematic* LEA exposures, particularly in the presence of negative moderating factors. Furthermore, this activity will help to illustrate that energy reallocation during energy scarcity may affect body systems differently, with systems being considered preferred, essential, reducible or expendable in varying ways between and within athletes.

The development and application of the physiological model templates could provide a major step in fostering strategies to address individual athletes, scenarios or behaviours that represent a potential for damaging or helpful outcomes. There is existing evidence that specific factors are associated with an altered risk of a body system disturbance within distinct cohorts exposed to LEA. For example, in a group of adolescent runners, a prior stress fracture was a common factor for developing another.<sup>42</sup> However, sex-specific risk factors included later menarche, lower body mass index (BMI) and prior involvement in gymnastics for girls; meanwhile boys with a previous history of a multidirectional movement sport (basketball) had a reduced fracture risk.<sup>42</sup> Similarly, we have already noted that a greater gynaecological age may decrease susceptibility to LH pulsatility perturbations when exposed to LEA.<sup>15</sup> The goal of mapping sophisticated physiological models is to extend and standardise the collection of such information to encourage a targeted approach to LEA treatment and prevention.

Our suggested protocol for undertaking a compartmentalised exploration of each body system is not the end goal; we appreciate the considerable overlap and interaction between many body systems, particularly with the endocrine system. Rather, it provides a dynamic, systematic template that supports the construction of a more intricate model by identifying the commonalities and differences between the effects of various characteristics of LEA, as well as intrinsic and extrinsic moderating factors, on body system outcomes. As the map of each system evolves with new insights, it may prompt discussion of other systems. Importantly, cross-talk, hierarchies and interactions among systems will emerge to link the physiological models into a strong network with the individual athlete at its centre.

Continued



**Table 1** Characteristics of low energy availability (LEA) exposure that are important in promoting energy partitioning or metabolic adaptation in athletes, differentiating adaptable and problematic LEA exposures

| Characteristic            | Comment                                                                                                                                                                                                                                                                                                                                                                    |
|---------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Magnitude/severity        | The severity of acute EA reduction (usually considered over 1 day) from an apparent optimal/normal EA of 40–45 kcal/kg FFM/day could be measured, potentially in zones or ranges of interest (eg, ~30 kcal, ~20 kcal, ~10 kcal)                                                                                                                                            |
| Duration                  | The duration of a consistent exposure to LEA in terms of days, weeks, months. Ranges of interest might include <1 week, 1–2 weeks, 2–4 weeks, weeks-months, months-years.                                                                                                                                                                                                  |
| Frequency                 | The frequency of exposure to LEA could be measured with potential zones of interest including frequent intermittent (eg, 5:2 day intermittent fasting, alternate day intensive competition); infrequent intermittent (eg, 1 week within 1–2 months); longer periodic (eg, 1 month within 6 months); or long-term (eg, 3 month competition phase within a 1 year calendar). |
| Origin                    | Whether the EA reduction is achieved primarily through a restriction in energy intake, an increase in exercise energy expenditure, or a ratio of both scenarios.                                                                                                                                                                                                           |
| Within day energy deficit | Analysis of the spread/timing of energy intake over the day in relation to exercise energy expenditure (eg, large time periods of energy deficit with 'backloading' or 'catch-up' energy intake (eg, exercise occurs early in the day, but majority of caloric intake is in the evening)).                                                                                 |
| Accumulated LEA dose      | A metric that combines duration, severity and frequency of LEA exposure may allow comparisons between different real-life scenarios: LEA dose = (magnitude × length of exposure/frequency)                                                                                                                                                                                 |

Figure 2 displays a proposed protocol for constructing a 'systems biology' approach to each body system or 'spoke' identified on the REDs Health Conceptual Model. To commence the physiological model for each REDs body system, we propose identifying the range of specific health and performance impairments that might occur from LEA exposure, along with details of the criterion tests and metrics that best assess the presence of such disturbances (figure 2, step 1). The full complexity of the model may also be achieved by separately noting the broad differential diagnoses that could be responsible for such impairments (not yet included in the Figure).

### Constructing the model: key characteristics for consideration with LEA exposure


Once the functional impairments of each body system are established, the focus should turn to the identification of key features of LEA exposure that may contribute to these. This step notes that energy mismatch is a primary and shared component of LEA, but each athlete's experience may be different (step 2, figure 2). Table 1 provides various LEA characteristics, along with patterns of such characteristics, which have been tested in the lab or observed in the field, within limitations.<sup>40 41</sup>

### LEA severity

A series of elegant studies conducted by Professor Anne Loucks and colleagues exposed sedentary females to stepwise reductions in EA (40–45, 25–30, ~20 and ~10 kcal/kg Lean Body Mass (LBM)/day) over ~5 days; biomarkers of reproductive health,<sup>24</sup> thyroid function<sup>17</sup> and bone metabolism<sup>20</sup> were measured across the different investigations. Figure 3, which summarises the relative changes in these biomarkers with LEA severity, normalised to their maximal responses, clearly demonstrates that hormonal patterns of specific body systems respond differently to the severity of LEA, at least after a short ~5-day exposure. Unfortunately, most LEA intervention studies have implemented a single high versus low LEA protocol, resulting in possibly dichotomous rather than graded changes.

One of the few longer term observations of graded levels of LEA severity implemented different energy deficits via changes in diet, exercise or both over three menstrual cycles in previously sedentary females.<sup>34</sup> The conditions (approximate energy balance and energy deficits of ~8%, 22% and 42% of daily energy expenditure) were subsequently converted to EA estimations (kcal/FFM/day), allowing comparisons of groups designated by the authors as 'low' EA (23–34 kcal/kg FFM/day), 'moderate' EA

(35–41 kcal/kg FFM/day) and 'high' EA (41–50 kcal/kg FFM/day).<sup>33</sup> Another study involved trained male endurance athletes who undertook a stepwise series of 14-day exposures equivalent to 25%, 50% and 75% reduction in their baseline estimated EA, with stages equivalent to overall EA of ~22, 17 and 9 kcal/kg FFM.<sup>43</sup> Most other controlled field-based prospective investigations of EA exposure in athletes or active people have involved comparison of a single, severe<sup>16 44–49</sup> (eg, 10–15 kcal/kg FFM/day) or moderate<sup>50–53</sup> (20–25 kcal/kg FFM/day) restriction of EA compared with 'adequate'/'high' EA or energy balance (40–50 kcal/kg FFM/day), with interventions lasting from 3 to 9 days. Other prospective observational studies<sup>54–56</sup> have identified self-imposed differences in EA in athletes over periods of 5 days to 12 weeks, enabling differences in health or performance outcomes between LEA exposures to be assessed. In short, although there is a continuum in the perturbations of biomarkers of different body systems to varying increases in LEA severity, the responses are not uniform across all cohorts or individual subjects.



**Figure 3** Dose–response effects of energy availability on luteinising hormone (LH) pulse frequency, plasma glucose (Glucose), markers of bone protein synthesis (PICP—procollagen I carboxyterminal propeptide) and mineralisation (OC—osteocalcin), and anabolic hormones that regulate bone formation (insulin, leptin, triiodothyronine (T3), and insulin-like growth factor-1 (IGF-1)). Created from data collected in sedentary women<sup>20 24</sup> and redrawn from Loucks.<sup>109</sup>

## Investigation of LEA duration

In real-life scenarios, the *duration* of athletes' energy mismatches may be related to the physiological and psychological sustainability of maintaining energy restriction; the sport's competition calendar, which includes periods that BMI/body fat manipulations may impact performance<sup>57–59</sup> and/or extreme in-event or training blocks with excessive energy expenditures.<sup>60–63</sup> Studies demonstrate that significant perturbations to body systems can occur in as little as 3–4 days of moderate to severe LEA exposures.<sup>17–20</sup> However, systematic time-course studies of longer duration are required to isolate the sequence of the translation of perturbations in biomarkers to functional outcomes of growing severity across the range of body systems. Such investigations are currently unavailable.

## Severity and duration

The interaction between the magnitude and duration of EA exposure should be explored. Changes in LH pulsatility and concentrations observed after three menstrual cycles of moderate EA reduction (~30 kcal/kg FFM/day)<sup>64</sup> were found to be similar to changes seen with 4–5 days of severe EA exposure (10 kcal/kg FFM/day).<sup>18 24</sup> Further investigation might warrant the development of a metric that can integrate the interaction of these variables such as an accumulated LEA dose [eg, magnitude×length of exposure (duration×frequency)].

## Patterns of accumulation

The pattern of accumulation of EA exposure might also be considered, as real-life scenarios demonstrate a variety of strategies of purposeful (or sometimes non-purposeful) energy restriction and/or increased exercise loads to periodise their body composition over a sporting season or career.<sup>57–59</sup> In addition, extreme exercise energy expenditure (EEE) in competitions or blocks of intensive training, which might pose another risk of LEA despite proactive nutrition support, can be implemented differently even in the same sport (eg, continuous lengthy exposure<sup>61</sup> or intermittent/alternate day racing<sup>62</sup> in road cycling or ultraendurance events ranging from hours to weeks).<sup>60 63</sup> This is likely to have implications for the frequency and characteristics of LEA exposure.

The discrepancies between the typical investigation of LEA and its real-life implementation should be considered. To date, studies of deliberate LEA exposure, both laboratory-controlled and field-based, have employed protocols of daily repetition of an identical energy restriction (and diet characteristics) and/or exercise load.<sup>15–20 24 47–49 51–53 65</sup> Although this strategy provides a practical way to ensure that certain EA conditions are achieved, they do not mimic real-life scenarios in which athletes alter their daily training and dietary practices according to the goals of their micro-, meso- and macrocycles of periodised training.<sup>66 67</sup> We are aware of only one series of studies with a laboratory-controlled EA intervention, implemented in a more ecologically valid protocol in which highly trained athletes continued their periodised training programme (varying the mode, frequency, intensity and duration of daily workouts) while continually adjusting their energy intake (EI) to achieve the daily EA target.<sup>44–46</sup>

## EI, EEE or both

Within the studies already noted, exposure to LEA has been usually achieved by a reduction in EI, an increase in EEE or a combination of both. Some investigations have included systematic comparisons of the effects of changing one or other of these components to achieve LEA.<sup>16 18 47</sup> Indeed, such a comparison

initiated the birth of the concept of LEA by distinguishing an energy mismatch, rather than low body fatness or high endurance training volumes, as the cause of the Female Athlete Triad symptomatology.<sup>68</sup> However, even though these studies have shown that LEA per se creates perturbations to a range of body systems, closer inspection reveals some subtle differences in outcomes according to relative changes in EI or EEE. Indeed, Loucks *et al* reported that LEA achieved via substantial increase in EEE was associated with less disturbance to LH pulsatility than similar LEA achieved through dietary restriction, with the difference attributed to a greater net reduction in glucose availability via the latter strategy.<sup>18</sup> In addition to direct effects on metabolic/hormonal perturbations, the relative contribution of changes in food intake and exercise patterns to LEA exposure may indirectly alter the manifestation of REDs health and performance outcomes due to their association with moderating factors. As noted in [table 2](#) (moderators), characteristics of some forms of exercise may protect specific body systems from the effects of LEA exposure by diverting energy to repair and adaptation, even though this may decrease the energy available to other systems.

## Within-day variability

Although EA is typically calculated or averaged across a 24-hour period, there has been distinct interest in within-day energy deficits that might occur at times during the day when there is separation between the periods of exercise and occasions of food intake (eg, exercise is undertaken earlier in the day while substantial eating occasions occur in the evening—'back-loading'). A separate metric, *within day energy balance*, has been developed to audit differences between estimates of EI and EEE in hourly increments, with focus on single or consecutive periods of either positive or negative energy balance, particularly by substantial margins that exceed the estimated energy and blood glucose buffer provided by liver glycogen stores (set at 300–400 kcal for females and males, respectively).<sup>37 69–71</sup> Observational studies applying such an assessment in athletes have reported that a greater duration of substantial within-day energy deficits is associated with perturbations in thyroid hormone,<sup>70</sup> resting metabolic rate,<sup>37 71</sup> oestrogen,<sup>71</sup> cortisol<sup>37 71</sup> and testosterone:cortisol ratio.<sup>37</sup> However, this concept has been less investigated than other aspects of LEA, with the few studies relying on self-reported data rather than controlled interventions.<sup>37 69–71</sup>

With these issues in mind, we propose that the REDs Physiological Model develops standardised terminology and metrics to identify characteristics of LEA exposure that may have different effects on perturbations of body systems ([figure 2](#)). Based on the available literature, the map of each body system can be constructed to note these characteristics' effects and identify the scenarios of greatest potential risk. In addition to contributing to greater insight for current practice, this information will highlight gaps in knowledge that might be systematically investigated in the future research.

## Constructing the model: moderating factors that may affect outcomes of LEA exposure

Individual responsiveness in the manifestation of REDs health/performance issues has been a consistent finding in cross-sectional studies of LEA prevalence, with differences in the number and severity of outcomes between and within cohorts.<sup>21 22</sup> Furthermore, prospective investigations have identified differences within<sup>34 48</sup> or between cohorts<sup>15</sup> with respect to immediate physiological/hormonal perturbations or functional outcomes arising from controlled LEA exposure. Therefore, it is logical

**Table 2** Examples of moderating factors that may increase or decrease the likelihood that low energy availability (LEA) is associated with a functional impairment of health or performance involving personal characteristics and medical history

| Categories                          | Potential moderating factors                  | Examples of moderation to (LEA risk), biomarker perturbations or expression of problematic LEA outcomes on (systems) implicated in REDs Health or Performance Conceptual Models                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
|-------------------------------------|-----------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Personal                            | Sex                                           | Males appear to better tolerate EA reduction: males respond to brief exposure to severe LEA with greater individual responses of bone turnover markers (bone health), <sup>48</sup> and less perturbation to T3 (energy metabolism), <sup>16</sup> testosterone <sup>16</sup> (reproductive system) and IGF-1 <sup>16</sup> (growth and development) concentrations. Females are at greater risk of developing stress fractures (bone health).                                                                                                                                                                                                                                                                   |
|                                     | Age/gynaecological age                        | Female: Perturbations of LH pulsatility with brief, severe LEA exposure are seen in females of young (<8y post-menarche) but not mature (>14y gynaecological age) <sup>15</sup> (reproductive system). Menopause is associated with a substantial decrease in bone BMD <sup>110 111</sup> (reproductive system). Female/male: Old(er) age is associated with anabolic resistance: that is, lower rates of muscle protein synthesis and muscle gain in response to resistance exercise and protein intake <sup>112</sup> (skeletal muscle function).                                                                                                                                                              |
|                                     | Genetics/epigenetics                          | Genetic factors that are unrelated to LEA may cause exacerbate several health outcomes identified as possible REDs outcomes, such as FHA <sup>113</sup> (reproductive system), low BMD <sup>114 115</sup> (bone health) and hyperlipidemia <sup>116</sup> (glucose and lipid metabolism).                                                                                                                                                                                                                                                                                                                                                                                                                        |
|                                     | Anatomical/ biomechanical features            | Poor running biomechanics and certain anatomical features may increase the risk of stress fractures <sup>117</sup> (bone health).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| Medical history                     | Co-existing medical disorders                 | Celiac disease can be associated with low BMD <sup>118</sup> (bone health) and iron deficiency <sup>119</sup> (haematological status). Testicular disease can cause male hypogonadism (reproductive system). <sup>120</sup> Traumatic brain injury is associated with hypogonadism in males and females <sup>121</sup> (reproductive system). Some medical conditions found in para-athletes may complicate the assessment of, or require different normative values of, RMR (energy metabolism) or BMD (bone health). <sup>122</sup> The autoimmune condition Hashimoto's Disease is typically associated with hypothyroidism and its various symptoms <sup>123</sup> (energy metabolism/regulation).           |
|                                     | Medication use                                | Non-steroidal anti-inflammatory drugs can cause gastrointestinal bleeding, contributing to iron loss and poor iron status <sup>124</sup> (haematological status). Glucocorticoid steroid use is associated with reductions in BMD and increased risk of fractures <sup>125</sup> (bone health).                                                                                                                                                                                                                                                                                                                                                                                                                  |
|                                     | Past medical history                          | Previous history of bone stress fractures is a risk factor for future stress fractures <sup>126 127</sup> (bone health). Past history of anorexia nervosa is a risk factor for low BMD and stress fractures <sup>127–129</sup> (bone health).                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
|                                     | Menstrual disturbances/low oestrogen (female) | Low oestrogen concentrations exacerbate the effect of LEA on bone loss via enhanced bone breakdown and suppressed bone formation <sup>80</sup> (bone health). Removal of monthly bleeding in amenorrheic females reduces the risk of iron deficiency <sup>77</sup> (haematological status).                                                                                                                                                                                                                                                                                                                                                                                                                      |
|                                     | PCOS/high androgen (female)                   | Female athletes with FHA and clinical/biochemical signs of hyperandrogenism had lower stress fracture risk than those with FHA alone <sup>130</sup> (bone health) (athlete availability).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Training characteristics            | Low impact exercise                           | Athletes undertaking sports involving low impact, non-weight bearing exercise have lower BMD than their counterparts in higher impact sports <sup>131 132</sup> and may even risk losing bone mass annually in the absence of impact loading stimuli <sup>133</sup> (bone health).                                                                                                                                                                                                                                                                                                                                                                                                                               |
|                                     | High impact exercise                          | Repetitive bone strain from dynamic, high impact, weight-bearing modes of exercise increases BMD and bone quality in athletes <sup>131 134</sup> (bone health). Repetitive footstrike (eg, running on hard surfaces) is associated with haemolysis and increased iron losses <sup>135 136</sup> (haematological status).                                                                                                                                                                                                                                                                                                                                                                                         |
|                                     | Training errors                               | Increases in bone strain beyond the adaptive capacity of the bone will independently increase the risk of bone stress injuries; factors include training volume/pattern, type of sport, and inappropriate footwear and training surfaces <sup>42 137–139</sup> (bone health).                                                                                                                                                                                                                                                                                                                                                                                                                                    |
|                                     | Resistance training                           | Resistance exercise (and protein intake) may rescue the decrease in resting muscle protein synthesis seen with LEA <sup>140</sup> and protect against the loss in lean body mass with an energy deficit <sup>141</sup> (skeletal muscle function).                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| Dietary/nutritional characteristics | Energy intake                                 | LEA underpinned primarily by restricted/reduced energy intake increases the magnitude of changes to LH pulsatility compared with an increased energy expenditure <sup>18</sup> (reproductive system).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
|                                     | Carbohydrate (CHO) availability               | Acute exposure to low CHO diet increases marker of exercise-associated bone breakdown and reduces marker of bone formation, <sup>142</sup> with these effects being greater than seen with LEA alone <sup>45</sup> (bone health). Acute exposure to low CHO diet increases the hepcidin response to exercise with likely reduction in iron absorption and recycling, with this effect being greater than associated with LEA alone <sup>46</sup> (haematological status). Acute carbohydrate restriction reduces T3 independently of LEA <sup>143</sup> (energy metabolism). Acute carbohydrate restriction reduces testosterone independently of LEA with exercise training <sup>144</sup> (energy metabolism). |
|                                     | Protein intake                                | Higher protein intake is required to optimise muscle protein synthesis in response to resistance exercise when exposed to LEA; inadequate protein intake likely results in reduced muscle protein gains <sup>140</sup> (training response).                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|                                     | Vitamin D status                              | Vitamin D deficiency/insufficiency may increase the risk of bone stress fractures <sup>145–147</sup> while a higher vitamin D intake/status may be associated with higher BMD and reduced risk of stress fractures <sup>148 149</sup> (bone health) (athlete availability).                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|                                     | Bioavailable iron intake                      | Inadequate intake of bioavailable iron is a risk factor for lower iron status/iron deficiency in athletes <sup>150 151</sup> (haematological status).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
|                                     | Calcium intake                                | Inadequate calcium intake may increase the risk of bone stress fractures, <sup>152</sup> while higher intake of dairy foods and calcium is associated with higher BMD and lower risk of stress fractures <sup>148</sup> (bone health). Consuming calcium prior to non-weight bearing exercise may reduce the PTH-associated increase in bone turnover during and after exercise <sup>153 154</sup> (bone health).                                                                                                                                                                                                                                                                                                |
|                                     | Energy density                                | Diets with low energy density are associated with increased satiety <sup>155</sup> and may reduce energy intake (LEA severity).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
|                                     | Intake of caffeine and other stimulants       | High intake of caffeine and its spread over the day may contribute to sleep disturbances in athletes <sup>156</sup> (sleep quality).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |

Continued

Table 2 Continued

| Categories                                                                                                                                                                                                                                                                                 | Potential moderating factors   | Examples of moderation to (LEA risk), biomarker perturbations or expression of problematic LEA outcomes on (systems) implicated in REDs Health or Performance Conceptual Models                                                                                                                                                                                                                                                                                                                                                                 |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Other                                                                                                                                                                                                                                                                                      | Psychological/lifestyle stress | Psychological stress may lead to FHA independent of physiological stress <sup>157 158</sup> (reproductive function).                                                                                                                                                                                                                                                                                                                                                                                                                            |
|                                                                                                                                                                                                                                                                                            | Environmental stress           | Training/competing at moderate altitude may increase resting metabolic rate <sup>159</sup> exacerbating risk of LEA unless adjustments are made to energy intake [LEA severity] and potentially reducing physiological adaptation <sup>160</sup> [training response] Training/competing at moderate altitude increases iron requirements exacerbating risks to iron status <sup>161</sup> [hematological status], with poor iron status reducing physiological adaptation to the training/altitude stimulus <sup>162</sup> [training response]. |
| BMD, bone mineral density; FHA, functional hypothalamic amenorrhea; IGF-1, insulin-like growth factor 1; LH, luteinizing hormone; PCOS, polycystic ovary syndrome; PTH, parathyroid hormone; REDs, Relative Energy Deficiency in Sport; RMR, resting metabolic rate; T3, triiodothyronine. |                                |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |

that an individual possesses intrinsic and experiences extrinsic factors that may moderate the manifestation of outcomes to LEA exposure on various body systems (ie, *moderating factors*). The mechanisms may aggravate the risk or effects (or interactive effects) of LEA on the body system, exacerbating or attenuating the perturbations or functional endpoints. Examples of such moderating factors and their interaction with various systems are summarised in [table 2](#), noting their potential effects on LEA and/or body system outcomes.

To further develop the REDs Physiological Model, we propose that factors that may moderate the risk of LEA per se or amplify/reduce its impact on each of the body systems or performance outcomes, be identified in a systematic way ([figure 2](#): step 3). The figure displays how moderating factors could be integrated into the physiological model for each body system, linking factors of specific interest to each identified health or performance outcome (step 4) in the REDs Conceptual Models and noting the direction of the effect. Again, a standardised approach to such activity would contribute insights into the prevention, diagnosis and treatment of REDs and promote a systematic programme of targeted research.

### Using the physiological model to predict LEA risks or explain outcomes

As the map of the physiological model for each body system becomes more sophisticated, it could be used to assess the contribution of LEA to the presentation of body system impairments, as well as develop hypotheses regarding the potential for such impairments before they occur. In the case of the athlete who has incurred health or performance decrements, the physiological model includes information about moderating factors and characteristics of LEA exposure that might assist in the diagnosis of underlying causes of these problems. Furthermore, these details could assist in clinical treatment by reducing/removing negative factors and introducing changes that are beneficial as part of a wholistic, multidisciplinary treatment plan. While correcting underlying LEA is the cornerstone of REDs therapy, a more personalised and nuanced approach can assist compliance and enhance treatment efficacy.

The physiological model could also be used in the strategic prevention of REDs. Disordered eating/eating disorders represent pathological conditions that frequently underpin LEA and REDs, with primary, secondary and tertiary prevention (treatment) requiring significant commitment from all within the sporting environment (eg, athlete, coach, performance and medical staff, parents, etc).<sup>72</sup> However, as previously discussed, some scenarios of LEA are integral to the athlete experience (particularly at the elite level) and may sometimes be associated with benefits to performance.<sup>59 73 74</sup> Furthermore, EA mismatches can occur inadvertently due to a variety of factors (eg, food insecurity, poor

food availability in the local environment, inadequate nutrition knowledge and practical skills, suppressed appetite and excessive exercise expenditures).<sup>73</sup> Although each of these scenarios can pose health and performance risks, understanding how each might cause a signature type of LEA of greater or lesser concern might prioritise efforts to educate athletes or organise activities that address the most pressing issues and underlying factors.<sup>12</sup>

### The future: integrating the model to recognise interactions between body systems

Interorgan cross-talk and interactions among various body systems, including in response to exercise, are both a historically recognised phenomenon as well as an area of evolving science.<sup>75 76</sup> Although the REDs Conceptual Models ([figure 1](#)) are drawn in two-dimensional simplicity, the reality of the overlay and interaction among body systems is recognised in research and practice. After complexity is better depicted via the theoretical mapping of the physiological models for each body system, digital online data visualisation skills might be employed to create presentation styles that better integrate the separate models. Features have already been built into this proposal for the REDs Physiological Model to aid this step. These include the interrogation of each body system using a standardised list of *moderating factors* that may assist in identifying the similarities and overlaps of some responses. Furthermore, the inclusion of some features of body systems into the list of *potential moderating factors* for another body system also accentuates the interaction. For example, the list of suggested moderating factors acknowledges that the cessation of menstrual periods (reproductive system) reduces iron losses<sup>77</sup> and is, therefore, a moderating (attenuating) factor for low iron status (haematological system) in females. Similarly, reproductive dysfunction and low levels of sex hormones (reproductive system) are associated with low BMD or bone stress injuries in both men<sup>78 79</sup> and women<sup>80–84</sup> (bone health) and represent a moderating (amplifying) factor to the effects of LEA. Opportunities to establish the hierarchy of energy allocation to different body systems and how this can differ between and within individuals in different scenarios may also be explored. This process of integrating the separate physiological models will challenge both the currently published science as well as provide visualisation opportunities; it will benefit from interdisciplinary input, including expertise and insights from adjacent fields pertaining to energy constraint and energy allocation during scenarios of energy scarcity.

### Learning from other perspectives of energy scarcity

Interest in the health and performance effects of LEA is a relatively new theme within the field of sports science/sports medicine. Although impairments of reproductive and/or bone status

were identified as far back as the 1980s in both male<sup>85–88</sup> and female athletes,<sup>89–90</sup> and the ‘Female Athlete Triad’ (Triad) was coined in 1993<sup>91</sup> and officially described in 1997,<sup>5</sup> the specific involvement of LEA was not recognised until the Triad update in 2007.<sup>4</sup> Since then, the evolution of Triad and REDs models has occurred largely via insights gained within sports science/sports medicine research and practice. Only recently have other perspectives on responses to energy mismatches been considered within the sports domain, with forays into evolutionary biology<sup>92–94</sup> enabled by recognition that “nothing in biology makes sense except in the light of evolution”.<sup>95</sup> Additionally, anthropologists have developed the reverse interest in athletes as a modern cohort with unique profiles of energy expenditure.<sup>60–96–97</sup>

Life History Theory proposes that humans have finite energy resources for which various biological processes related to growth, health, activity and reproduction compete with different outcomes over their life span.<sup>31–32–92</sup> During periods of energy scarcity, which occur for numerous reasons over our individual and collective history, energy insufficiency is managed by allocating it to tissues and functions in a hierarchical manner, with priority afforded to those that offer the greatest immediate survival value. Such dynamic energy trade-offs are context specific, varying according to factors such as sex, life stage, environment, severity/duration of energy scarcity and other stressors; our capacity to dynamically redistribute energy resources derives from phenotypic plasticity shaped by human history.<sup>31–32–92</sup> Aspects of life history theory explain many of the features of the updated REDs Conceptual Models: energy partitioning can occur at different levels and timescales across and within athletes; it can range from transient and reversible, representing a normal and necessary, adaptive response through to more severe impairments to body systems that nevertheless represent survival priorities.<sup>92</sup> These perspectives reinforce the caveats identified in the updated REDs Conceptual Models,<sup>1</sup> emphasising that LEA and the response to it are not uniform and always negative,<sup>92–94</sup> and confirming that the focus should be on *problematic* LEA exposure. Observations on athletes via the life history lens have included commentary on priorities of energy allocation during energetic stress, noting considerable reliance on the context of both the energy restriction and the individual athlete/environment,<sup>92–94</sup> but also the apparent importance of preserving physical capacity.<sup>98</sup>

Athletes have allowed a further examination of the constrained energy model of Pontzer and colleagues.<sup>99–100</sup> The constrained energy model proposes that humans share a set of evolved mechanisms to maintain total energy expenditure (TEE) within a narrow range, and to dynamically reduce behavioural and non-behavioural (mainly basal metabolic rate; BMR) components of energy expenditure to compensate for increases in physical activity.<sup>99–100</sup> While compensation may include adjustment to other activities in the day, reduction in whole body BMR via decreased metabolic expenditure on various body systems may occur to limit TEE.<sup>99</sup> This hypothesis opposes the additive energy model in which deliberate high levels of physical activity in athletes, hunter gatherers, labourers and other populations are simply added to the other components of their energy expenditure. The constrained energy model is intriguing and explains REDs as a necessary reduction in metabolic rate in athletes with high EEE. However, it has been noted that its support comes from cross-sectional observational data and statistical models that compare extremely varied populations.<sup>101</sup> It is likely that the level of Constraint vs Addition with respect to physical activity varies across populations, according to factors such as

individual characteristics, the level of energy expenditure and the presence of energy balance vs energy deficit.<sup>96–101</sup> Indeed, insights from endurance athletes have contributed to a model that predicts that at low activity levels, exercise is added to total energy expenditure, with moderate to high EEE leading to some partial compensation to achieve energy savings, including behavioural changes to reduce non-exercise activity thermogenesis (NEAT).<sup>94</sup> Meanwhile, very high activity levels that increase TEE beyond alimentary limits to cause LEA will accrue the greatest energy compensations and energy reallocation among biological processes.<sup>94</sup>

The study of obesity also provides insight into REDs-related issues; weight loss is often accompanied by metabolic adaptation or adaptive thermogenesis, as displayed by a reduction in TEE and resting metabolic rate that exceeds that predicted from reduced BM/FFM and decreases in thermic effect of food intake and BM-associated EEE.<sup>102–104</sup> Because this reduction in TEE may contribute to failure to maintain weight loss in the long-term,<sup>102–104</sup> there is interest in understanding and combatting adaptive thermogenesis and its components.<sup>105–106</sup> In the obesity literature, there is also evidence of phenotypic and genetic differences in the presence and components of adaptive thermogenesis.<sup>105–106</sup> Ultimately, a better understanding of REDs will be gained from interdisciplinary expertise and insights from outside the sports medicine/sports science environment. Of course, we recognise the complex differences between obese populations, current and past hunter gatherer populations and athletes, but also some overlaps. Indeed, understanding differences as well as similarities in responses to energy mismatches may help to refine the physiological model of LEA in athletes.

## CONCLUSION

Our understanding and presentation of REDs have evolved from the concept first introduced in 2014,<sup>2</sup> with the REDs Conceptual Models for Health and Performance from the 2023 REDs Consensus Statement<sup>1</sup> showcasing a more nuanced and rigorous depiction of the science and practical experiences of the syndrome. Nevertheless, continued development of the knowledge and messaging around REDs to support prevention, diagnosis<sup>107</sup> and treatment strategies<sup>12</sup> that accommodate the range of real-life outcomes seen in athletes is needed. Recognising that REDs entails the accumulation of health and performance outcomes stemming from problematic exposure to LEA, the field requires greater insights into the continuum between *adaptable* LEA vs *problematic* LEA exposure. We have proposed the development of a more complex physiological model approach for each body system implicated in REDs, underpinned by a ‘systems biology mindset’. We encourage a sequence of activities to develop individual maps for each body system, which could then be integrated to acknowledge interrelationships and cross-talk among organs/systems. This approach will enable a more nuanced assessment of the individual athlete—considering their specific combination of LEA exposure and secondary moderators to determine if the outcome is likely to be positive, neutral or negative. In short, the evolution of the scientific and practical REDs concept requires an interdisciplinary field of study that integrates complex biological interactions and uses a holistic approach towards functional outcomes.<sup>108</sup> We hope that the development and continued evolution of these individual body system, physiologically-based models will enhance the understanding and exploration of the complex science underpinning REDs; that they assist clinicians in providing better education, prevention and treatment programmes for REDs; and most

importantly, that they aid athletes in improving how they manage their own training and nutrition.

**Twitter** Louise M Burke @LouiseMBurke, Kathryn E Ackerman @DrKateAckerman, Ida A Heikura @IdaHeikura, Anthony C Hackney @AC\_Hackney and Trent Stellingwerff @TStellingwerff

**Acknowledgements** The authors would like to thank the International Olympic Committee for their prioritisation and support of athlete health, Professor Margo Mountjoy for leading the Working Group for the 2023 consensus statement on Relative Energy Deficiency in Sport (REDs), and all members of the Group for their insights and discussions. We acknowledge all who have contributed to our understanding of Low Energy Availability and invite new contributions to the suggested model discussed in this paper. We note interactions with Drs Alannah McKay, Peter Peeling, Naama Constantini and Claire Badenhorst to test-drive a REDs Physiological Model within the theme of the hematological system, and earlier discussions with Professors Mountjoy and Jorunn Sundgot-Borgen in devising the model goals. In particular, the pioneering work of Professor Anne Loucks is recognised. We thank Grace Saville for assistance in the preparation of the manuscript.

**Contributors** LMB was responsible for leading and coordinating this manuscript. All authors were involved in the drafting, revising, and approval of the final manuscript prior to submission.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent for publication** Not applicable.

**Ethics approval** Not applicable.

**Provenance and peer review** Not commissioned; externally peer-reviewed.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iDs

Louise M Burke <http://orcid.org/0000-0001-8866-5637>  
Kathryn E Ackerman <http://orcid.org/0000-0003-2626-7785>  
Ida A Heikura <http://orcid.org/0000-0002-1088-428X>  
Anthony C Hackney <http://orcid.org/0000-0002-6607-1472>  
Trent Stellingwerff <http://orcid.org/0000-0002-4704-8250>

#### REFERENCES

- Mountjoy ML, Ackerman KE, Bailey DM, et al. The 2023 International Olympic Committee's (IOC) consensus statement on relative energy deficiency in sports (Reds). *BJSM* 2023.
- Mountjoy M, Sundgot-Borgen J, Burke L, et al. The IOC consensus statement: beyond the female athlete Triad—relative energy deficiency in sport (RED-S). *Br J Sports Med* 2014;48:491–7.
- Mountjoy M, Sundgot-Borgen JK, Burke LM, et al. IOC consensus statement on relative energy deficiency in sport (RED-S): 2018 update. *Br J Sports Med* 2018;52:687–97.
- Nattiv A, Loucks AB, Manore MM, et al. American college of sports medicine position stand. The female athlete Triad. *Med Sci Sports Exerc* 2007;39:1867–82.
- Otis CL, Drinkwater B, Johnson M, et al. American college of sports medicine position stand. The female athlete Triad. *Med Sci Sports Exerc* 1997;29:i–ix.
- Nattiv A, De Souza MJ, Koltun KJ, et al. The male athlete Triad-A consensus statement from the female and male athlete Triad coalition part 1: definition and scientific basis. *Clin J Sport Med* 2021;31:345–53.
- Fredericson M, Kussman A, Misra M, et al. The male athlete Triad-A consensus statement from the female and male athlete Triad coalition part II: diagnosis, treatment, and return-to-play. *Clin J Sport Med* 2021;31:349–66.
- Stellingwerff T, Mountjoy ML, McClusky WTP, et al. A review of the scientific rationale, development, and validation of the IOC relative energy deficiency in sport clinical assessment tool - version 2 (IOC Reds Cat2): by a sub-group of the IOC consensus on. *REDs British Journal of Sports Medicine* 2023.
- Mountjoy M et al. Avoiding the 'Reds card': an introductory editorial to the International Olympic Committee's (IOC) consensus statement on relative energy deficiency in sports (Reds). *Br J Sports Med* 2023.
- Mathisen Tfer et al. Body composition practices as antecedents of Reds: minimizing risks and optimizing benefits of assessment - a narrative review by a sub-group of the IOC consensus on relative energy deficiency in sport (Reds). *Br J Sports Med* 2023.
- Hackney AC, Melin AK, Ackerman KE, et al. REDS alert: male athletes be wary and scientists take action! *Br J Sports Med* 2023.
- Torstveit MK, Ackerman KE, Constantini N, et al. Primary, secondary, and tertiary prevention of Reds - a narrative review by a sub-group of the IOC consensus on relative energy deficiency in sport (Reds). *Br J Sports Med* 2023.
- Holtzman B, Rogers M, Heikura IA, et al. Methods to study Reds - a narrative review by a sub-group of the IOC consensus on relative energy deficiency in sport (Reds). *Br J Sports Med* 2023.
- De Souza MJ, Strock NCA, Ricker EA, et al. The path towards progress: A critical review to advance the science of the female and male athlete Triad and relative energy deficiency in sport. *Sports Med* 2022;52:13–23.
- Loucks AB. The response of luteinizing hormone Pulsatility to 5 days of low energy availability disappears by 14 years of gynecological age. *J Clin Endocrinol Metab* 2006;91:3158–64.
- Koehler K, Hoerner NR, Gibbs JC, et al. Low energy availability in exercising men is associated with reduced Leptin and insulin but not with changes in other metabolic hormones. *J Sports Sci* 2016;34:1921–9.
- Loucks AB, Heath EM. Induction of low-T3 syndrome in exercising women occurs at a threshold of energy availability. *Am J Physiol* 1994;266(3 Pt 2):R817–23.
- Loucks AB, Verdun M, Heath EM. Low energy availability, not stress of exercise, alters LH Pulsatility in exercising women. *J Appl Physiol (1985)* 1998;84:37–46.
- Hilton LK, Loucks AB. Low energy availability, not exercise stress, suppresses the diurnal rhythm of Leptin in healthy young women. *Am J Physiol Endocrinol Metab* 2000;278:E43–9.
- Ilhe R, Loucks AB. Dose-response relationships between energy availability and bone turnover in young exercising women. *J Bone Miner Res* 2004;19:1231–40.
- Logue D, Madigan SM, Delahunt E, et al. Low energy availability in athletes: A review of prevalence, dietary patterns, physiological health, and sports performance. *Sports Med* 2018;48:73–96.
- Logue DM, Madigan SM, Melin A, et al. Low energy availability in athletes 2020: an updated narrative review of prevalence, risk, within-day energy balance, knowledge, and impact on sports performance. *Nutrients* 2020;12:835.
- Stellingwerff T, Heikura IA, Meeusen R, et al. Overtraining syndrome (OTS) and relative energy deficiency in sport (RED-S): shared pathways, symptoms and complexities. *Sports Med* 2021;51:2251–80.
- Loucks AB, Thuma JR. Luteinizing hormone Pulsatility is disrupted at a threshold of energy availability in regularly Menstruating women. *J Clin Endocrinol Metab* 2003;88:297–311.
- Williams NI, Young JC, McArthur JW, et al. Strenuous exercise with caloric restriction: effect on luteinizing hormone secretion. *Med Sci Sports Exerc* 1995;27:1390–8.
- Ruffing KM, Koltun KJ, De Souza MJ, et al. Moderate weight loss is associated with reductions in LH pulse frequency and increases in 24-hour Cortisol with no change in perceived stress in young Ovulatory women. *Physiology & Behavior* 2022;254:113885.
- Baer JT. Endocrine parameters in Amenorrheic and Eumenorrheic adolescent female runners. *Int J Sports Med* 1993;14:191–5.
- Gibbs JC, Williams NI, Scheid JL, et al. The Association of a high drive for thinness with energy deficiency and severe Menstrual disturbances: confirmation in a large population of exercising women. *Int J Sport Nutr Exerc Metab* 2011;21:280–90.
- Gibbs JC, Williams NI, Mallinson RJ, et al. Effect of high dietary restraint on energy availability and Menstrual status. *Med Sci Sports Exerc* 2013;45:1790–7.
- De Souza MJ, Koltun KJ, Strock NC, et al. Rethinking the concept of an energy availability threshold and its role in the female athlete Triad. *Current Opinion in Physiology* 2019;10:35–42.
- Gadgil M, Bossert WH. Life historical consequences of natural selection. *The American Naturalist* 1970;104:1–24.
- Stearns SC. *The evolution of life histories Oxford*. New York: Oxford University Press, 1992.
- Lieberman JL, De Souza MJ, Wagstaff DA, et al. Menstrual disruption with exercise is not linked to an energy availability threshold. *Med Sci Sports Exerc* 2018;50:551–61.
- Williams NI, Leidy HJ, Hill BR, et al. Magnitude of daily energy deficit predicts frequency but not severity of Menstrual disturbances associated with exercise and caloric restriction. *Am J Physiol Endocrinol Metab* 2015;308:E29–39.
- Reed JL, De Souza MJ, Mallinson RJ, et al. Energy availability discriminates clinical Menstrual status in exercising women. *J Int Soc Sports Nutr* 2015;12:11.
- Lundy B, Torstveit MK, Stenqvist TB, et al. Screening for low energy availability in male athletes: attempted validation of LEAM-Q. *Nutrients* 2022;14:1873.
- Torstveit MK, Fahrenholtz I, Stenqvist TB, et al. Within-day energy deficiency and metabolic perturbation in male endurance athletes. *Int J Sport Nutr Exerc Metab* 2018;28:419–27.
- Elliott-Sale KJ, Minahan CL, de Jonge XAKJ, et al. Methodological considerations for studies in sport and exercise science with women as participants: A working guide for standards of practice for research on women. *Sports Med* 2021;51:843–61.
- Loucks AB. No evidence of absence of an energy availability threshold for Menstrual disturbances. *Med Sci Sports Exerc* 2019;51:1790.
- Areta JL, Taylor HL, Koehler K. Low energy availability: history, definition and evidence of its endocrine, metabolic and physiological effects in prospective studies in females and males. *Eur J Appl Physiol* 2021;121:1–21.

- 41 Burke LM, Lundy B, Fahrenholtz IL, *et al.* Pitfalls of conducting and interpreting estimates of energy availability in free-living athletes. *Int J Sport Nutr Exerc Metab* 2018;28:350–63.
- 42 Tenforde AS, Fredericson M, Sayres LC, *et al.* Identifying sex-specific risk factors for low bone mineral density in adolescent runners. *Am J Sports Med* 2015;43:1494–504.
- 43 Jurov I, Keay N, Rauter S. Reducing energy availability in male endurance athletes: a randomized trial with a three-step energy reduction. *J Int Soc Sports Nutr* 2022;19:179–95.
- 44 Burke LM, Whitfield J, Ross MLR, *et al.* Short severe energy restriction with refueling reduces body mass without altering training-associated performance improvement. *Med Sci Sports Exerc* 2023;55:1487–98.
- 45 Fensham NC, Heikura IA, McKay AKA, *et al.* Short-term carbohydrate restriction impairs bone formation at rest and during prolonged exercise to a greater degree than low energy availability. *J Bone Miner Res* 2022;37:1915–25.
- 46 McKay AKA, Peeling P, Pyne DB, *et al.* Six days of low carbohydrate, not energy availability, alters the iron and immune response to exercise in elite athletes. *Med Sci Sports Exerc* 2022;54:377–87.
- 47 Martin D, Papageorgiou M, Colgan H, *et al.* The effects of short-term low energy availability, achieved through diet or exercise, on cognitive function in oral contraceptive users and Eumenorrheic women. *Appl Physiol Nutr Metab* 2021;46:781–9.
- 48 Papageorgiou M, Elliott-Sale KJ, Parsons A, *et al.* Effects of reduced energy availability on bone metabolism in women and men. *Bone* 2017;105:191–9.
- 49 Papageorgiou M, Martin D, Colgan H, *et al.* Bone metabolic responses to low energy availability achieved by diet or exercise in active Eumenorrheic women. *Bone* 2018;114:181–8.
- 50 Jurov I, Keay N, Spudić D, *et al.* Inducing low energy availability in trained endurance male athletes results in poorer explosive power. *Eur J Appl Physiol* 2022;122:503–13.
- 51 Kojima C, Ishibashi A, Ebi K, *et al.* Exogenous glucose oxidation during endurance exercise under low energy availability. *PLoS One* 2022;17:e0276002.
- 52 Kojima C, Ishibashi A, Tanabe Y, *et al.* Muscle Glycogen content during endurance training under low energy availability. *Med Sci Sports Exerc* 2020;52:187–95.
- 53 Ishibashi A, Kojima C, Tanabe Y, *et al.* Effect of low energy availability during three consecutive days of endurance training on iron metabolism in male long distance runners. *Physiol Rep* 2020;8:e14494.
- 54 Woods AL, Garvican-Lewis LA, Lundy B, *et al.* New approaches to determine fatigue in elite athletes during intensified training: resting metabolic rate and pacing profile. *PLoS One* 2017;12:e0173807.
- 55 Kettunen O, Ihalainen JK, Ohtonen O, *et al.* Energy availability during training camp is associated with signs of Overreaching and changes in performance in young female cross-country skiers. *Bioméd Hum Kinet* 2021;13:246–54.
- 56 VanHeest JL, Rodgers CD, Mahoney CE, *et al.* Ovarian suppression impairs sport performance in Junior elite female swimmers. *Med Sci Sports Exerc* 2014;46:156–66.
- 57 Heydenreich J, Kayser B, Schutz Y, *et al.* Total energy expenditure, energy intake, and body composition in endurance athletes across the training season: A systematic review. *Sports Med - Open* 2017;3.
- 58 Burke LM, Slater GJ, Matthews JJ, *et al.* ACSM expert consensus statement on weight loss in weight-category sports. *Curr Sports Med Rep* 2021;20:199–217.
- 59 Stellingwerff T. Case study: body composition Periodization in an Olympic-level female middle-distance runner over a 9-year career. *Int J Sport Nutr Exerc Metab* 2018;28:428–33.
- 60 Longman DP, Dolan E, Wells JCK, *et al.* Patterns of energy allocation during energetic scarcity: evolutionary insights from ultra-endurance events. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* 2023;281:111422.
- 61 Lombardi G, Lanteri P, Graziani R, *et al.* Bone and energy metabolism parameters in professional cyclists during the Giro D'Italia 3-weeks stage race. *PLoS ONE* 2012;7:e42077.
- 62 Heikura IA, Quod M, Strobel N, *et al.* n.d. Alternate-day low energy availability during spring classics in professional cyclists. *Int J Sports Physiol Perform*;14:1233–43.
- 63 Nikolaidis PT, Veniamakis E, Rosemann T, *et al.* Nutrition in ultra-endurance: state of the art. *Nutrients* 2018;10:1995.
- 64 Koltun KJ, De Souza MJ, Scheid JL, *et al.* Energy availability is associated with luteinizing hormone pulse frequency and induction of Luteal phase defects. *The Journal of Clinical Endocrinology & Metabolism* 2020;105:185–93.
- 65 Loucks AB, Heath EM, Law T. Dietary restriction reduces luteinizing hormone (LH) pulse frequency during waking hours and increases LH pulse amplitude during sleep in young Menstruating women. *J Clin Endocrinol Metab* 1994;78:910–5.
- 66 Mujika I, Halson S, Burke LM, *et al.* An integrated, Multifactorial approach to Periodization for optimal performance in individual and team sports. *Int J Sports Physiol Perform* 2018;13:538–61.
- 67 Stellingwerff T, Maughan RJ, Burke LM. Nutrition for power sports: middle-distance running, track Cycling, rowing, canoeing/Kayaking, and swimming. *Journal of Sports Sciences* 2011;29:579–89.
- 68 Loucks AB. Energy availability, not body fatness, regulates reproductive function in women. *Exerc Sport Sci Rev* 2003;31:144–8.
- 69 Deutz RC, Benardot D, Martin DE, *et al.* Relationship between energy deficits and body composition in elite female gymnasts and runners. *Medicine & Science in Sports & Exercise* 2000;32:659–68.
- 70 Lundstrom EA, De Souza MJ, Canil HN, *et al.* Sex differences and indications of metabolic compensation in within-day energy balance in elite division 1 swimmers. *Appl Physiol Nutr Metab* 2023;48:74–87.
- 71 Fahrenholtz IL, Sjödin A, Benardot D, *et al.* Within-day energy deficiency and reproductive function in female endurance athletes. *Scand J Med Sci Sports* 2018;28:1139–46. 10.1111/sms.13030 Available: <http://doi.wiley.com/10.1111/sms.13030> Available: 2018.28.issue-3
- 72 Wells KR, Jeacocke NA, Appaneal R, *et al.* The Australian Institute of sport (AIS) and national eating disorders collaboration (NEDC) position statement on disordered eating in high performance sport [British journal of sports medicine 2020:bjsports-2019-1]. *Br J Sports Med* 2020;54:1247–58.
- 73 Burke L, Fahrenholtz I, Garthe I, *et al.* Low energy availability: challenges and approaches to measurement and treatment. In: Burke L, Deakin V, Minehan M, eds. *Clinical Sports Nutrition*. 6th ed. Sydney: McGraw-Hill 2021:110–43.
- 74 Melin AK, Areta JL, Heikura IA, *et al.* Direct and indirect impact of low energy availability on sports performance. *Scandinavian Med Sci Sports* 2023.
- 75 Sabaratnam R, Wojtaszewski JFP, Højlund K. Factors mediating Exercise-Induced organ Crosstalk. *Acta Physiol (Oxf)* 2022;234:e13766.
- 76 Gonzalez-Gil AM, Elizondo-Montemayor L. The role of exercise in the interplay between Myokines, Hepatokines, Osteokines, Adipokines, and modulation of inflammation for energy substrate redistribution and fat mass loss: A review. *Nutrients* 2020;12:1899.
- 77 Petkus DL, Murray-Kolb LE, Scott SP, *et al.* Iron status at opposite ends of the Menstrual function spectrum. *J Trace Elem Med Biol* 2019;51:169–75.
- 78 Dolan E, McGoldrick A, Davenport C, *et al.* An altered hormonal profile and elevated rate of bone loss are associated with low bone mass in professional horse-racing jockeys. *J Bone Miner Metab* 2012;30:534–42.
- 79 Heikura IA, Uusitalo AL, Stellingwerff T, *et al.* Low energy availability is difficult to assess but outcomes have large impact on bone injury rates in elite distance athletes. *Int J Sport Nutr Exerc Metab* 2018;28:403–11.
- 80 De Souza MJ, West SL, Jamal SA, *et al.* The presence of both an energy deficiency and estrogen deficiency exacerbate alterations of bone metabolism in exercising women. *Bone* 2008;43:140–8.
- 81 Gibbs JC, Nattiv A, Barrack MT, *et al.* Low bone density risk is higher in exercising women with multiple Triad risk factors. *Med Sci Sports Exerc* 2014;46:167–76.
- 82 Mitchell DM, Tuck P, Ackerman KE, *et al.* Altered trabecular bone morphology in adolescent and young adult athletes with Menstrual dysfunction. *Bone* 2015;81:24–30.
- 83 Barrack MT, Rauh MJ, Nichols JF. Prevalence of and traits associated with low BMD among female adolescent runners. *Med Sci Sports Exerc* 2008;40:2015–21.
- 84 Ackerman KE, Cano Sokoloff N, DE Nardo Maffazioli G, *et al.* Fractures in relation to Menstrual status and bone parameters in young athletes. *Med Sci Sports Exerc* 2015;47:1577–86.
- 85 De Souza MJ, Arce JC, Pescatello LS, *et al.* Gonadal hormones and semen quality in male runners. A volume threshold effect of endurance training. *Int J Sports Med* 1994;15:383–91.
- 86 Hetland ML, Haarbo J, Christiansen C. Low bone mass and high bone turnover in male long distance runners. *J Clin Endocrinol Metab* 1993;77:770–5.
- 87 Hackney AC, Sinning WE, Bruot BC. Reproductive hormonal profiles of endurance-trained and untrained males. *Med Sci Sports Exerc* 1988;20:60–5.
- 88 Hackney AC, Dolny DG, Ness RJ. Comparison of resting reproductive hormonal profiles in select athletic groups. *Biol Sport* 1988;4:200–8.
- 89 Dale E, Gerlach DH, Wilhite AL. Menstrual dysfunction in distance runners. *Obstet Gynecol* 1979;54:47–53.
- 90 Hetland ML, Haarbo J, Christiansen C, *et al.* Running induces Menstrual disturbances but bone mass is unaffected, except in Amenorrheic women. *Am J Med* 1993;95:53–60.
- 91 Yeager KK, Agostini R, Nattiv A, *et al.* The female athlete Triad: disordered eating, Amenorrhea, osteoporosis. *Med Sci Sports Exerc* 1993;25:775–7.
- 92 Shirley MK, Longman DP, Elliott-Sale KJ, *et al.* A life history perspective on athletes with low energy availability. *Sports Med* 2022;52:1233–34.
- 93 Oliveira-Junior G, Pinto RS, Shirley MK, *et al.* The Skeletal muscle response to energy deficiency: A life history perspective. *Adaptive Human Behavior and Physiology* 2022;8:114–29.
- 94 Dolan E, Koehler K, Areta J, *et al.* Energy constraint and compensation: insights from endurance athletes. *COMP Biochem Physiol A Mol Integr Physiol* 2023;285:111500.
- 95 Dobzhansky T. Nothing in biology makes sense except in the light of evolution. *The American Biology Teacher* 1973;35:125–9.
- 96 Areta JL. Physical performance during energy deficiency in humans: an evolutionary perspective. *Comp Biochem Physiol A Mol Integr Physiol* 2023;284:111473.
- 97 Pontzer H, Yamada Y, Sagayama H, *et al.* Daily energy expenditure through the human life course. *Science* 2021;373:808–12.

- 98 Thurber C, Dugas LR, Ocobock C, *et al.* Extreme events reveal an Alimentary limit on sustained maximal human energy expenditure. *Sci Adv* 2019;5:eaaw0341.
- 99 Pontzer H, Durazo-Arvizu R, Dugas LR, *et al.* Constrained total energy expenditure and metabolic adaptation to physical activity in adult humans. *Curr Biol* 2016;26:410–7.
- 100 Pontzer H. Constrained total energy expenditure and the evolutionary biology of energy balance. *Exerc Sport Sci Rev* 2015;43:110–6.
- 101 Gonzalez JT, Batterham AM, Atkinson G, *et al.* Perspective: is the response of human energy expenditure to increased physical activity additive or constrained? *Adv Nutr* 2023;14:406–19.
- 102 Fothergill E, Guo J, Howard L, *et al.* "Persistent metabolic adaptation 6 years after "the biggest loser" competition". *Obesity (Silver Spring)* 2016;24:1612–9.
- 103 Hall KD. "Energy compensation and metabolic adaptation: "the biggest loser" study reinterpreted". *Obesity* 2022;30:11–3.
- 104 Johannsen DL, Knuth ND, Huizenga R, *et al.* Metabolic slowing with massive weight loss despite preservation of fat-free mass. *J Clin Endocrinol Metab* 2012;97:2489–96.
- 105 Aronne LJ, Hall KD, M. Jakicic J, *et al.* Describing the weight-reduced state: physiology, behavior, and interventions. *Obesity* 2021;29:59–24. 10.1002/oby.23086 Available: <https://onlinelibrary.wiley.com/doi/10.1002/oby.23086>
- 106 Ravussin E, Smith SR, Ferrante AW Jr. Physiology of energy expenditure in the weight-reduced state. *Obesity* 2021;29:531–8. 10.1002/oby.23095 Available: <https://onlinelibrary.wiley.com/doi/10.1002/oby.23095>
- 107 Stellingwerff T, Mountjoy ML, McClusky WTP, *et al.* Review of the scientific rationale, development and validation of the International Olympic committee relative energy deficiency in sport clinical assessment tool: V.2 (IOC Reds Cat2)—By a subgroup of the IOC consensus on Reds. *Br J Sports Med* 2023.
- 108 Nielsen J. Systems biology of metabolism. *Annu Rev Biochem* 2017;86:245–75.
- 109 Loucks AB. Commentary 2. female athlete triad and energy availability. In: Burke L, Deakin V, eds. *Clinical Sports Nutrition*. 5th ed. Sydney: McGraw Hill 2015:140–45.
- 110 Sipilä S, Törmäkangas T, Sillanpää E, *et al.* Muscle and bone mass in middle-aged women: role of menopausal status and physical activity. *J Cachexia Sarcopenia Muscle* 2020;11:698–709.
- 111 Hadjidakis D, Kokkinakis E, Sfakianakis M, *et al.* The type and time of Menopause as decisive factors for bone mass changes. *Eur J Clin Invest* 1999;29:877–85.
- 112 Phillips BE, Williams JP, Greenhaff PL, *et al.* Physiological adaptations to resistance exercise as a function of age. *JCI Insight* 2017;2:e95581.
- 113 Fontana L, Garzia E, Marfia G, *et al.* Epigenetics of functional hypothalamic Amenorrhea. *Front Endocrinol (Lausanne)* 2022;13:953431.
- 114 Curtis EM, Fuggle NR, Cooper C, *et al.* Epigenetic regulation of bone mass. *Best Pract Res Clin Endocrinol Metab* 2022;36:101612.
- 115 Ralston SH, Uitterlinden AG. Genetics of osteoporosis. *Endocr Rev* 2010;31:629–62.
- 116 Bilen O, Pokharel Y, Ballantyne CM. Genetic testing in hyperlipidemia. *Cardiol Clin* 2015;33:267–75.
- 117 Korpelainen R, Orava S, Karpakka J, *et al.* Risk factors for recurrent stress fractures in athletes. *Am J Sports Med* 2001;29:304–10.
- 118 Lungaro L, Manza F, Costanzini A, *et al.* Osteoporosis and celiac disease: updates and hidden pitfalls. *Nutrients* 2023;15:1089.
- 119 Talarico V, Giancotti L, Mazza GA, *et al.* Iron deficiency anemia in celiac disease. *Nutrients* 2021;13:1695.
- 120 Basaria S. Male Hypogonadism. *Lancet* 2014;383:1250–63.
- 121 Rothman MS, Arciniegas DB, Filley CM, *et al.* The Neuroendocrine effects of traumatic brain injury. *J Neuropsychiatry Clin Neurosci* 2007;19:363–72.
- 122 Jonvik KL, Vardardottir B, Broad E. How do we assess energy availability and RED-S risk factors in para athletes *Nutrients* 2022;14:1068.
- 123 Klubo-Gwiedzinska J, Wartofsky L. Hashimoto thyroiditis: an evidence-based guide to etiology, diagnosis and treatment. *Pol Arch Intern Med* 2022;132:16222.
- 124 Tai FWD, McAlindon ME. Non-Steroidal anti-inflammatory drugs and the gastrointestinal tract. *Clin Med (Lond)* 2021;21:131–4.
- 125 Van Staa TP, Leufkens HG, Abenham L, *et al.* Use of oral corticosteroids and risk of fractures. *J Bone Miner Res* 2000;15:993–1000.
- 126 Kelsey JL, Bachrach LK, Procter-Gray E, *et al.* Risk factors for stress fracture among young female cross-country runners. *Med Sci Sports Exerc* 2007;39:1457–63.
- 127 Tenforde AS, Sayres LC, McCurdy ML, *et al.* Identifying sex-specific risk factors for stress fractures in adolescent runners. *Med Sci Sports Exerc* 2013;45:1843–51.
- 128 Herzog W, Minne H, Deter C, *et al.* Outcome of bone mineral density in anorexia Nervosa patients 11.7 years after first admission. *J Bone Miner Res* 1993;8:597–605.
- 129 Hartman D, Crisp A, Rooney B, *et al.* Bone density of women who have recovered from anorexia Nervosa. *Int J Eat Disord* 2000;28:107–12.
- 130 Javed A, Kashyap R, Lteif AN. Hyperandrogenism in female athletes with functional hypothalamic Amenorrhea: a distinct phenotype. *Int J Womens Health* 2015;7:103–11.
- 131 Bellver M, Del Rio L, Jovell E, *et al.* Bone mineral density and bone mineral content among female elite athletes. *Bone* 2019;127:393–400.
- 132 Scofield KL, Hecht S. Bone health in endurance athletes: runners, cyclists, and swimmers. *Curr Sports Med Rep* 2012;11:328–34.
- 133 Sherk VD, Barry DW, Villalon KL, *et al.* Bone loss over 1 year of training and competition in female cyclists. *Clin J Sport Med* 2014;24:331–6.
- 134 Stojanović E, Radovanović D, Dalbo VJ, *et al.* Basketball players possess a higher bone mineral density than matched non-athletes, swimming, soccer, and volleyball athletes: a systematic review and meta-analysis. *Arch Osteoporos* 2020;15:123.
- 135 Miller BJ, Pate RR, Burgess W. Foot impact force and Intravascular hemolysis during distance running. *Int J Sports Med* 1988;9:56–60.
- 136 Telford RD, Sly GJ, Hahn AG, *et al.* Footstrike is the major cause of hemolysis during running. *Journal of Applied Physiology* 2003;94:38–42.
- 137 Warden SJ, Edwards WB, Willy RW. Preventing bone stress injuries in runners with optimal workload. *Curr Osteoporos Rep* 2021;19:298–307.
- 138 Tenforde AS, DeLuca S, Wu AC, *et al.* Prevalence and factors associated with bone stress injury in middle school runners. *PM R* 2022;14:1056–67.
- 139 Matheson GO, Clement DB, McKenzie DC, *et al.* Stress fractures in athletes. A study of 320 cases. *Am J Sports Med* 1987;15:46–58.
- 140 Areta JL, Burke LM, Camera DM, *et al.* Reduced resting Skeletal muscle protein synthesis is rescued by resistance exercise and protein ingestion following short-term energy deficit. *Am J Physiol Endocrinol Metab* 2014;306:E989–97.
- 141 Murphy C, Koehler K. Energy deficiency impairs resistance training gains in lean mass but not strength: A meta-analysis and meta-regression. *Scand J Med Sci Sports* 2022;32:125–37. 10.1111/sms.14075 Available: <https://onlinelibrary.wiley.com/doi/10.1111/sms.14075>
- 142 Heikura IA, Burke LM, Hawley JA, *et al.* A short-term Ketogenic diet impairs markers of bone health in response to exercise. *Front Endocrinol (Lausanne)* 2019;10:880.
- 143 Spaulding SW, Chopra IJ, Sherwin RS, *et al.* Effect of caloric restriction and dietary composition of serum T3 and reverse T3 in man. *J Clin Endocrinol Metab* 1976;42:197–200.
- 144 Lane AR, Duke JW, Hackney AC. Influence of dietary carbohydrate intake on the free testosterone: Cortisol ratio responses to short-term intensive exercise training. *Eur J Appl Physiol* 2010;108:1125–31.
- 145 Shimasaki Y, Nagao M, Miyamori T, *et al.* Evaluating the risk of a fifth Metatarsal stress fracture by measuring the serum 25-Hydroxyvitamin D levels. *Foot Ankle Int* 2016;37:307–11.
- 146 Millward D, Root AD, Dubois J, *et al.* Association of serum vitamin D levels and stress fractures in collegiate athletes. *Orthopaedic Journal of Sports Medicine* 2020;8:232596712096696.
- 147 Ruohola J-P, Laaksi I, Ylikomi T, *et al.* Association between serum 25(OH)D concentrations and bone stress fractures in Finnish young men. *J Bone Miner Res* 2006;21:1483–8.
- 148 Nieves JW, Melsop K, Curtis M, *et al.* Nutritional factors that influence change in bone density and stress fracture risk among young female cross-country runners. *PM R* 2010;2:740–50.
- 149 Burgi AA, Gorham ED, Garland CF, *et al.* High serum 25-Hydroxyvitamin D is associated with a low incidence of stress fractures. *J Bone Miner Res* 2011;26:2371–7.
- 150 Snyder AC, Dvorak LL, Roepke JB. Influence of dietary iron source on measures of iron status among female runners. *Medicine & Science in Sports & Exercise* 1989;21:7–10.
- 151 Gibson-Smith E, Storey R, Ranchordas M. Dietary intake, body composition and iron status in experienced and elite climbers. *Front Nutr* 2020;7:122.
- 152 Barrack MT, Fredericson M, Tenforde AS, *et al.* Evidence of a cumulative effect for risk factors predicting low bone mass among male adolescent athletes. *Br J Sports Med* 2017;51:200–5.
- 153 Haakonssen EC, Ross ML, Knight EJ, *et al.* The effects of a calcium-rich pre-exercise meal on biomarkers of calcium homeostasis in competitive female cyclists: a randomised crossover trial. *PLoS One* 2015;10:e0123302.
- 154 Lundy B, McKay AKA, Fensham NC, *et al.* The impact of acute calcium intake on bone turnover markers during a training day in elite male rowers. *Med Sci Sports Exerc* 2023;55:55–65.
- 155 Reed JL, Bowell JL, Hill BR, *et al.* n.d. Exercising women with Menstrual disturbances consume low energy dense Foods and Beverages.
- 156 Gardiner C, Weakley J, Burke LM, *et al.* The effect of caffeine on subsequent sleep: A systematic review and meta-analysis. *Sleep Med Rev* 2023;69:101764.
- 157 Berga SL, Marcus MD, Loucks TL, *et al.* Recovery of ovarian activity in women with functional hypothalamic Amenorrhea who were treated with cognitive behavior therapy. *Fertil Steril* 2003;80:976–81.
- 158 Fioroni L, Fava M, Genazzani AD, *et al.* Life events impact in patients with secondary Amenorrhea. *J Psychosom Res* 1994;38:617–22.
- 159 Woods AL, Sharma AP, Garvican-Lewis LA, *et al.* Four weeks of classical altitude training increases resting metabolic rate in highly trained middle-distance runners. *Int J Sport Nutr Exerc Metab* 2017;27:83–90.
- 160 McLean BD, Buttifant D, Gore CJ, *et al.* Year-to-year variability in Haemoglobin mass response to two altitude training camps. *Br J Sports Med* 2013;47 Suppl 1(Suppl 1):i51–8.
- 161 Govus AD, Garvican-Lewis LA, Abbiss CR, *et al.* Pre-altitude serum Ferritin levels and daily oral iron supplement dose mediate iron parameter and hemoglobin mass responses to altitude exposure. *PLoS One* 2015;10:e0135120.
- 162 Okazaki K, Stray-Gundersen J, Chapman RF, *et al.* Iron insufficiency diminishes the erythropoietic response to moderate altitude exposure. *J Appl Physiol (1985)* 2019;127:1569–78.