

Research Bank

Journal article

Distinct profiles of social skill in adults with autism spectrum disorder and schizophrenia

Morrison, Kerriane E., Pinkham, Amy E., Penn, David L., Kelsven, Skylar, Ludwig, Kelsea and Sasson, Noah J.

This is the peer reviewed version of the following article:

Morrison, K. E., Pinkham, A. E., Penn, D. L., Kelsven, S., Ludwig, K. and Sasson, N. J. (2017). Distinct profiles of social skill in adults with autism spectrum disorder and schizophrenia. *Autism Research*, 10(5), pp. 878-887, which has been published in final form at <https://doi.org/10.1002/aur.1734>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions. This article may not be enhanced, enriched or otherwise transformed into a derivative work, without express permission from Wiley or by statutory rights under applicable legislation. Copyright notices must not be removed, obscured or modified. The article must be linked to Wiley's version of record on Wiley Online Library and any embedding, framing or otherwise making available the article or pages thereof by third parties from platforms, services and websites other than Wiley Online Library must be prohibited.



Published in final edited form as:

Autism Res. 2017 May ; 10(5): 878–887. doi:10.1002/aur.1734.

Distinct Profiles of Social Skill in Adults with Autism Spectrum Disorder and Schizophrenia

Kerriane E. Morrison, B.S., Amy E. Pinkham, Ph.D., David L. Penn, Ph.D., Skylar Kelsven, B.A., Kelsey Ludwig, B.S., and Noah J. Sasson, Ph.D

School of Behavioral and Brain Sciences, The University of Texas at Dallas, Richardson, TX 75080 (K.E.M., A.E.P., S.K., N.J.S); Department of Psychiatry, University of Texas Southwestern Medical School, Dallas, TX 75390 (A.E.P); Department of Psychology, University of North Carolina, Chapel Hill, NC 27599 (D.L.P, K.L); Department of Psychology, Australian Catholic University, Melbourne, Victoria, Australia (D.L.P)

Scientific Abstract

Overlapping social impairments in Autism Spectrum Disorder (ASD) and Schizophrenia (SCZ) contributed to decades of diagnostic confusion that continues to this day in some clinical settings. The current study provides the first direct and detailed comparison of social behavior in the two disorders by identifying profiles of social skill in adults with ASD ($n=54$), SCZ ($n=54$), and typically-developing (TD) controls ($n=56$) during a real-world social interaction. ASD and SCZ groups exhibited poorer social skill, both overall and on most discrete abilities, relative to the TD group. Direct comparison of ASD to SCZ revealed distinct behavioral profiles, with ASD uniquely characterized by fewer interactive behaviors, and SCZ characterized by greater impaired gaze and flat/inappropriate affective responses. Additionally, IQ was associated with both overall social skill and many discrete social skills in SCZ, but was largely unrelated to social skill in ASD. These results indicate that overlapping social deficits in ASD and SCZ are comprised of both shared and distinct social skill impairments. The largest distinctions — reduced social reciprocity but better expressivity in ASD relative to SCZ, and a greater role of IQ in social skill for SCZ than ASD — highlight disorder-specific features that can improve etiological understanding, diagnostic differentiation, and treatment strategies.

Lay Abstract

Adults with Autism Spectrum Disorder (ASD) and those with Schizophrenia (SCZ) share impairments in social interaction that has led to diagnostic confusion in some settings. In the current study, we provide a direct comparison of social behavior in these two groups to determine the ways in which social skill in ASD and SCZ are similar and different. Compared to typically-developing individuals, ASD and SCZ adults exhibited poorer social skill, both overall and on most discrete social abilities. However, when compared to each other, distinctions between ASD and SCZ emerged. ASD was uniquely characterized by fewer interactive behaviors, whereas SCZ

Corresponding author: Noah Sasson, Ph.D. Phone: (972) 883-2541, fax: (972) 883-2491, nsasson@utdallas.edu.

Address for Correspondence and Reprints: Noah Sasson, Ph.D., School of Behavioral and Brain Sciences, The University of Texas at Dallas, 800 W. Campbell Rd., Richardson, TX 75080

Conflict of Interest Disclosure: None reported.

was characterized by greater impaired gaze and flat or inappropriate expressiveness. Additionally, IQ was associated with both overall social skill and many discrete social skills in SCZ, but was largely unrelated to social skill in ASD. These distinctions highlight features specific to each condition that can improve diagnosis and treatment of both ASD and SCZ.

Keywords

Social Skills; Social Interaction; Social Cognition; Social Behavior

Introduction

Adults with Autism Spectrum Disorder (ASD) and Schizophrenia (SCZ) are both characterized by social dysfunction, a broad term for maladaptive behaviors that hinder optimal functioning in social contexts [Bellack et al., 1990; Klin et al., 1992]. Social impairments in ASD and SCZ are associated with occupational challenges [Marwaha & Johnson, 2004; Taylor et al., 2015], interpersonal difficulties [Horan et al., 2006; Howlin et al., 2004] (e.g., small social networks, low rates of friendship), and a reduced quality of life [Barneveld et al., 2014; Eack & Newhill, 2007]. These outcomes are present in adults with “high-functioning” ASD, who despite being intellectually-capable, still struggle with the social demands of adult responsibilities [Seltzer et al., 2004]. Similarly, social dysfunction in individuals with SCZ persists even when pharmacological treatment improves symptom presentation in other domains [Pinkham, et al., 2012]. For both groups, psychosocial treatments have yielded some gains in social functioning [Bishop-Fitzpatrick et al., 2014; Wykes et al., 2008], but the extent of improvement and generalization to real-world outcomes are often limited.

Similarities in the social features of ASD and SCZ contributed to decades of diagnostic confusion [Sasson et al., 2011] that continue to this day in some clinical settings [Mandell et al., 2012]. ASD and SCZ do share some behavioral, genetic, and neurobiological characteristics [American Psychiatric Association, 2013; Crespi et al., 2010; Pinkham et al., 2008; Rapoport et al., 2009; Sullivan et al., 2012], yet they have distinct developmental trajectories [Sasson et al., 2011], prominent non-shared features (e.g., positive symptoms in SCZ and restricted interests in ASD) [Leifker et al., 2009; Nadig et al., 2010], and respond differently to psychosocial treatment [Turner-Brown et al., 2008]. These distinctions suggest that the nature and mechanisms of social impairment in the two disorders may differ, but their identification has been hindered by a lack of direct comparisons of ASD and SCZ using systematically-matched designs. Direct comparisons can reveal disorder-specific aspects of social impairment missed when using the traditional approach of comparing each clinical group to typically-developing (TD) controls [Sasson et al., 2011].

Indeed, previous direct comparisons between ASD and SCZ suggest that the two disorders share general impairments in social cognition and emotional understanding (Couture et al., 2010; Craig et al., 2004; Konstantareas & Hewitt, 2001), yet the processes underlying these impairments may differ [Sasson et al., 2011; Pinkham et al., 2012]. For instance, individuals with ASD demonstrate greater impairments in orienting to relevant social and emotional

information [Sasson et al., 2007; 2015] whereas individuals with SCZ may exhibit a greater tendency to “jump to conclusions” in social situations [Freeman 2007]. Some evidence also suggests that social impairments in the two disorders may manifest in divergent ways, with ASD characterized more by hyposociality and under-mentalizing, but SCZ—at least in its paranoid variant—characterized more by hypersociality and over-mentalizing [Crespi et al 2010; Crespi & Badcock, 2008]. Further, although IQ is positively associated with social outcomes for both groups (Green et al, 2004; Eaves & Ho, 2008; Howlin et al., 2004), general intellectual ability may be more central in SCZ than ASD for both social cognitive skill [Sasson et al., 2015] and social functioning [Green et al., 2004; Saulnier & Klin, 2007]. A large portion of the variance in functional outcome is attributable to general intelligence in SCZ [Green et al., 2004] whereas social disability and poor functional outcomes persist in ASD even in individuals with average to above-average intellectual ability [Klin, 2000; Nicpon et al., 2010; Saulnier & Klin, 2007]. Ultimately, refined understanding of these and other distinctions in social cognition and behavior in the two disorders can be used to improve diagnostic practices, inform etiological understanding, and guide targeted treatment approaches.

A primary component of social behavior is social skill, defined as the set of behaviors employed to achieve social goals in different contexts and environments [Mueser & Bellack, 1998]. Social skill is impaired in ASD and SCZ relative to TD controls, both globally [Constantino et al., 2000; Pinkham & Penn, 2006] and in more discrete social behaviors such as pragmatic language usage and nonverbal skills [Bishop, 1998; Patterson et al., 2001]. However, studies of social skill in ASD and SCZ have to this point been conducted independently, and large differences in sample characteristics—mostly children in ASD but adults in SCZ— and methodological distinctions between studies prevent the ability to identify patterns of social skill convergence and divergence in ASD and SCZ.

The current study compares social skills between ASD and SCZ, both relative to TD adults and to each other, using systematic direct observation of social behaviors during a real-world social situation common to adult daily functioning. We hypothesized that relative to TD controls, the two clinical groups would exhibit reduced social skill, both overall and across most discrete social behaviors [Mueser et al., 1990; Nadig et al., 2010; Paul et al., 2009; Verhoeven et al., 2013]. However, based upon social aspects emphasized in prior research and clinical reports, we predicted several differences between the two clinical groups: whereas the SCZ group would be characterized by inappropriate affective behaviors and more impaired paralinguistic behaviors (i.e., how words are spoken), including verbal clarity and fluency [Bellack et al., 1990; Verhoeven et al., 2013], the ASD group was expected to demonstrate less reciprocity [Paul et al., 2009], and more repetitive verbal content and movement [Bodfish et al., 2000]. Further, given prior research indicating a greater role of general cognition to social impairment in SCZ relative to ASD [Sasson et al., 2015], intellectual ability was predicted to correlate with social skills in SCZ but not ASD. Finally, discriminant function analysis explored the constellation of social skills that best distinguished the groups. Collectively, these analyses sought to identify distinctions in social behavior in ASD and SCZ that can improve clinical differentiation and inform treatment.

Method

Participants

Participants ($N=164$) were drawn from a multisite study assessing social cognition and functioning in ASD and SCZ. The three groups (ASD $n=54$, SCZ $n=54$, TD controls $n=56$) were matched on gender and comparable on age, ethnicity, years of education completed, and IQ as estimated by the verbal section of the WRAT-3 [Wilkinson, 1993] (see Table 1), a common measure of premorbid IQ in schizophrenia. All participants were between 18 and 65, had an estimated IQ over 70, and had no substance abuse or dependence in the previous six months. Using G Power, we found that with a value $=.80$ and $\alpha=.05$ sample sizes of 31 participants per group would be sufficient to detect medium sized effects (e.g. $F\text{-squared}=.15$). Thus, we recruited over 50 individuals per group to enhance our ability to detect smaller effects.

Adults with ASD were recruited from the UT-Dallas Autism Research Collaborative, a registry of local adults with a confirmed diagnosis of ASD via the Autism Diagnostic Observation Schedule (ADOS) [Lord et al., 2000]. Participants with SCZ were recruited from Metrocare Services, a nonprofit mental health services provider in Dallas, TX and from the Outreach and Support Intervention Services (OASIS) program affiliated with the University of North Carolina at Chapel Hill (UNC-CH). All SCZ diagnoses were confirmed with the Structured Clinical Interview for DSM-IV. The SCZ group exhibited more positive and negative symptoms than the ASD group on the Positive and Negative Syndrome Scale (PANSS; Kay et al., 1987), and had higher rates of antipsychotic medication usage (see Table 1). TD adults reporting no developmental disabilities or mental illnesses were recruited from the local community in Dallas, TX, and Chapel Hill, NC. All participants completed the study at university facilities between October 2012 and October 2015, provided written informed consent, and received compensation. The Institutional Review Boards of UT-Dallas and UNC-CH approved this protocol.

Procedure

Participants completed scene 1 of the Social Skills Performance Assessment (SSPA) [Patterson et al., 2001], a three-minute role-play in which individuals initiate and maintain a conversation within the context of meeting a new neighbor, played by the experimenter. Role-plays were videotaped, and social skills were coded using scoring criteria from the Conversation Probe (CP) [Pinkham & Penn, 2006], a related social skills role-playing paradigm used previously with adults with SCZ and with individuals exhibiting the Broad Autism Phenotype [Sasson et al., 2013] that, like the SSPA, also assesses social behavior upon meeting somebody for the first time. Although the CP has not previously been applied to the SSPA, it was used here because it is largely redundant with the SSPA in the behaviors it captures and their coding criteria, but is more comprehensive in that it assesses additional discrete social behaviors (e.g., separate codes for flat and appropriate affect) and allows for greater sensitivity (i.e., a nine-point scale vs. the five-point scale of the SSPA).

Coded behaviors include: *clarity* (easily understood speech); *fluency* (ease of speech production); *meshing* (ease of conversational turn-taking); *gaze* (frequency and duration of

appropriate eye contact); *flat affect* (nonverbal emotional expressivity); *appropriate affect* (expression of suitable emotion); *involvement* (interest and investment in the conversation); *asks questions* (number of questions asked by the participant); *verbal content* (topics discussed are appropriate to the situation); *social anxiety* (behavioral indicators of anxiousness); and *overall social skill* (a global measure of the participant's ability to interact and communicate effectively). Because the SSPA was initially developed for capturing social skills relevant to SCZ, three additional behaviors known to affect social interaction quality in ASD were also measured: *repetitive verbal content* (repeating phrases or topics), *repetitive movement* (presence of motor or sensory stereotypies), and *verbosity* (total amount of speech). Repetitive behaviors, though often conceptualized as non-social behaviors, are related to social deficits in ASD [Lam et al., 2008] and impact social interaction quality [Nadig et al., 2010; Paul et al., 2009], and verbosity (e.g., the tendency to monologue) is common in cognitively-able adults with ASD [Laugeson & Ellingsen, 2014].

Two coders were trained to reliability at the beginning of the study, and reliability was also assessed at the study mid-point and study end (ICCs > .66 for all behaviors; see Table 3). Coders were blind to group membership and rated the 164 videos independently. The average of both coders' ratings constituted the final scores.

Statistical Analysis

Groups were compared on social skills using MANOVA with follow-up two-tailed univariate tests corrected for multiple comparisons ($\alpha_{pc} = 0.0036$) using SPSS version 23 (IBM). Significant univariate results were followed up with post-hoc Tukey tests. Data used in the MANOVA did not violate assumptions of normality, and variances did not differ between compared groups.

Discriminant function analysis (DFA) was also used to determine the constellation of social skills that together best characterize group membership. DFA is a multivariate technique that predicts group membership based on an optimally weighted linear combination of variables. When predicting three groups, DFA plots the groups along two orthogonal functions (i.e., factors or components) according to the linear combination of behaviors. The distance between the multivariate group means on each function describes how the groups are separated; therefore functions describe how one group is uniquely characterized relative to the other groups. The contribution of each social skill to group discrimination is determined by examining the magnitude and direction of the standardized coefficients of the linear combination of social skills separating the groups. To focus on profiles of discrete behaviors, the global behaviors (overall social skill and social anxiety) were excluded from this analysis. Because tolerance values for the remaining social skills indicated that the DFA assumption of non-colinearity was violated [Tabachnick & Fidell, 2007], colinearity was reduced by condensing the number of discrete behaviors with a factor analysis. Four factors accounting for 62% of the variance (see Table 2) emerged that align with prior factor analyses of social skill [Mueser & Bellack, 1998]: paralinguistic (clarity, fluency, and meshing), interactive (asks questions and involvement), nonverbal (gaze, appropriate affect, and flat affect), and verbal content behaviors. These four factors and the three ASD-related behaviors were included in the DFA analysis.

We also examined the bivariate correlations of IQ with social skills for each group. Further, as symptom severity differed between the clinical groups, we examined the correlations between positive, negative, and general symptoms and overall social skill. Only negative symptom severity was significantly correlated with overall social skill in the SCZ group ($r = -.56, p < .001$, all other correlations below $-.27$). Because the clinical groups also differed in antipsychotic medication usage, we used a MANCOVA with negative symptoms and medication usage as a covariate to explore whether any reported group effects would remain when controlling for these differences between the groups.

Results

Group Differences in Social Skills

Table 3 displays group means for all social behaviors. MANOVA results indicated group differences across all social skills, $\lambda = .488, F(28, 296) = 4.56, p < .001$. The groups significantly differed on overall social skill, verbal content, fluency, gaze, involvement, asks questions, appropriate affect, social anxiety, and repetitive movement (p 's $< .003$; univariate tests displayed in Table 3). Groups did not significantly differ on flat affect, meshing, clarity, repetitive verbal content, and verbosity (p 's $> .009$). Post-hoc tests demonstrated that the ASD and SCZ groups each scored lower than the TD group on overall social skill, verbal content, fluency, gaze, asks questions, appropriate affect, and social anxiety (p 's $< .003$), and the ASD group, but not the SCZ group, scored lower than the TD group on involvement (ASD $p < .001$, SCZ $p = .072$) and higher on repetitive movement (ASD $p < .001$, SCZ $p = .728$).

When the clinical groups were directly compared to each other, the ASD group demonstrated significantly more repetitive movement ($p < .001$) and asked significantly fewer questions ($p = .021$) than the SCZ group. The ASD group was also rated as more impaired, scoring lower than the SCZ group at the mean level across all social skills except clarity and flat affect, where SCZ showed greater impairment than ASD, and verbosity, where the groups appeared comparable (comparisons and Cohen's d values are displayed in Table 3). Other moderate effect size differences favoring the SCZ group over the ASD group (Cohen's d s between $.31$ and $.48$; see Table 3) emerged for verbal content, involvement, repetitive verbal content, and overall social skill, but these did not reach statistical significance.

Discriminant Function Analysis

The DFA resulted in one function separating the ASD group from the SCZ and TD groups that accounted for 84% of the variance, $\lambda(14) = .632, p < .001$, canonical $R^2 = .560$, 95% confidence interval (CI) [0.42, 0.64]. A second function was also significant, separating the SCZ group from the ASD and TD groups, $\lambda(6) = .920, p = .041$, canonical $R^2 = .283$, 95% CI [0.13, 0.37]. Figure 1 displays the discriminant function plot. The pattern of the standardized coefficients (see Table 4) suggests that the ASD group is best differentiated from the SCZ and TD groups by a profile of more repetitive movement, less appropriate verbal content, and fewer interactive behaviors (i.e., involvement and asking questions), whereas the SCZ group is best separated from the ASD and TD groups by a profile of less appropriate nonverbal behaviors (e.g., appropriate gaze and affect), less repetitive moment, and more interactive behaviors.

Correlation of IQ with Social Skills

As shown in Table 5, IQ was positively associated with verbal content, clarity, fluency, meshing, involvement, social anxiety, and overall social skill for the SCZ group, and involvement, asks questions, appropriate affect, verbosity, and overall social skill in the TD group, but only verbosity in the ASD group. To support that IQ significantly correlated across social skills in SCZ but not ASD, we used the randomization test procedure [Sherman & Funder, 2009] in the R programming environment to determine whether the number of correlations observed for each group was significant compared to the number of correlations that would be observed by chance alone. The pattern of the correlations between IQ and social skills was significant for the SCZ ($p < .01$) and TD groups ($p = .02$) but not the ASD group ($p = .40$).

Post Hoc Examination of Negative Symptoms and Medication Usage

Finally, we repeated the MANOVA with negative symptoms and antipsychotic medication usage as a covariate, and results changed only minimally. The overall MANCOVA was still significant, $\lambda = .705$, $F(14, 91) = 2.732$, $p = .002$, with the ASD group continuing to score significantly lower than the SCZ group on repetitive movement, but interactive behaviors no longer survived correction ($p = .02$). However, now the ASD group also demonstrated worse overall social skill ($p < .001$) than the SCZ group, and also approached the corrected alpha on demonstrating poorer involvement ($p = .004$) and appropriate affect ($p = .004$).

Discussion

This study compared social skill profiles of adults with ASD, SCZ, and TD controls. Consistent with prior work [Patterson et al., 2001; Verhoeven et al., 2013], the ASD and SCZ groups demonstrated poorer social skill than TD controls during a real-world social interaction, both overall and on most specific behaviors, with the ASD but not the SCZ group less socially involved and demonstrating more repetitive movement. Although these results provide further confirmation that ASD and SCZ share broad social impairments relative to TD controls, several notable differences between the clinical groups emerged when they were directly compared.

First, the ASD group asked significantly fewer questions of their conversation partner than did the SCZ group. This deficit in reciprocity is consistent with prior studies of ASD [Paul et al., 2009; Verhoeven et al., 2013], and is particularly salient given that participants were specifically tasked with “getting to know” a new neighbor. Further, given that the three groups did not differ in the amount of speech (i.e., verbosity) used during the task, the low rate of questions asked by the ASD group cannot be explained by lower rates of talking generally. Instead, it suggests that the function of the interaction differed for those with ASD, with a greater focus on providing information and discussing personal interests rather than attempting to learn more about their conversational partner. In contrast, the higher rate of questions by the SCZ group suggests that their social impairments are less related to deficits in social reciprocity. Additionally, when controlling for the higher negative symptoms and greater medication use of the SCZ group in post-hoc analyses, not only did patterns largely remain similar, but overall social skill also emerged as significantly worse in

ASD compared to SCZ. Collectively, the size and consistent direction of effects favoring the SCZ group over the ASD group highlights the significant social disability experienced by adults with ASD, and underscores the tremendous need for increasing research and services directed towards adults within a population that historically has focused on children.

DFA analyses revealed that ASD was uniquely characterized by a profile of reduced interactive behaviors and increased repetitive movement and inappropriate conversational content, a combination that corresponds with the ASD phenotype [American Psychiatric Association, 2013]. Collectively, this profile is consistent with prior research indicating that individuals with ASD demonstrate reduced social motivation [Chevallier et al., 2012], a greater tendency to monologue [Laugeson & Ellingson, 2014], and a penchant for directing conversation towards the self and personal interests [Nadig et al., 2010]. In contrast, the SCZ group was uniquely characterized by poor nonverbal behavior (i.e., presence of poorly modulated eye gaze and affective responses) along with relatively higher levels of interactive behaviors. Taken together, these findings hint at a discrepancy in the two groups between social motivation and social expressivity, with the SCZ group demonstrating an interest in getting to know their social partner but reduced expressive competency while doing so, whereas the ASD group exhibited the reverse pattern. Given the history of diagnostic overlap and social functioning similarities in ASD and SCZ [Sasson et al., 2011], particularly when symptoms seemingly overlap (e.g., negative symptoms), these discrepant social skill patterns may offer clinical value for diagnostic differentiation. The predictive accuracy of DFA can be evaluated statistically [Tabachnick & Fidell, 2007], suggesting that DFA offers promise as a tool for diagnostically classifying individuals based on their social skill profiles.

Our results also have implications for treatment. Developing targeted social skills interventions for ASD and SCZ may prove more beneficial than employing a “one size fits all” approach or administering treatments developed for one disorder on the other [Turner-Brown et al., 2008]. Rather, results here suggest that social skill programs should differ between the two disorders by emphasizing reciprocity in ASD, but affective responses and neurocognitive remediation in SCZ. IQ was significantly related to overall social skill and many discrete social skills in SCZ, but largely unrelated to social skill in ASD. This finding is consistent with studies suggesting that general cognitive ability accounts for a large portion of the variance in social and functional outcomes in SCZ [Green et al., 2004], whereas social deficits in ASD persist even in the intellectually-gifted [Nicpon et al., 2010]. Future work is encouraged to explore whether general cognitive remediation may produce downstream effects for improving social skills in SCZ more so than ASD and examine comprehensively how general cognitive ability relates to broader social disability in each disorder.

Our results should be interpreted in the context of several limitations. First, the SSPA task is a structured role-play that mimics a real-world scenario but may not fully capture naturally occurring social behavior. The SSPA requires the participant to take on a specific role (e.g., a neighbor) and engage in a social interaction with an explicit goal in mind (e.g., introducing themselves). These task demands may not translate to real world situations, especially for the clinical groups, who may be less inclined to initiate or maintain a social interaction with

an unfamiliar social partner. The task demands used here assume that social interaction would occur, which may not be the case. It is unclear whether results would differ if the nature of the social task were changed. Second, although the SSPA and CP code for a wide range of social skills, these codes are not exhaustive and may have missed other relevant social skills for the group comparison. Third, although the sample used here is larger and better matched than previous direct comparisons of ASD and SCZ, it was not possible to match the groups on medication use and the presence of positive, negative, and general symptoms. However, performance was generally better in SCZ than in ASD despite their greater medication use and symptoms, and this pattern only strengthened when negative symptoms were co-varied in analyses. Thus, the lower social skill exhibited by the ASD group despite lower levels of medication and symptoms suggest that their performance may be a conservative estimate of their social skill impairment. Finally, although the current findings suggest both similarities and discrepancies in social behavior between ASD and SCZ, they cannot address what cognitive, neural, or developmental mechanisms may be driving these patterns. ASD and SCZ do share many behavioral, genetic, and neurobiological features [Crespi et al., 2010; Pinkham et al., 2008; Rapoport et al., 2009; Sasson et al., 2011; Sullivan et al., 2012], and continued direct comparisons are needed to further refine our understanding of shared and divergent aspects of pathophysiology in the two disorders.

Notwithstanding these limitations, the current study highlights distinct social skill impairments in SCZ and ASD. Using a large, well-matched sample of ASD, SCZ, and TD adults on a task of real-world social behavior, we found that overall social skill impairments in ASD and SCZ are subserved by unique constellations of social skills. Whereas the ASD group demonstrated the largest social skill deficits and a distinct deficit in social reciprocity, the SCZ group was differentiated by impaired affective and nonverbal behaviors. These differences, along with the differential relationship of IQ and social skills for each group, have important implications for treating social reciprocity in ASD and social expressivity in SCZ. Future work is encouraged to examine how these patterns of social skills relate to real-world outcomes, as well as explore the cognitive and neurobiological mechanisms underlying social skill deficits in ASD and SCZ.

Acknowledgments

This research was funded by grants from the National Institute of Mental Health at National Institutes of Health (R01 MH093432, PIs Harvey, Penn, Pinkham; R15 MH1015945, PI Sasson). We thank all the individuals who participated in this study and Isis Nelson-Graham (The University of Texas at Dallas) for her assistance with data collection and management.

References

- American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders DSM-5. Washington DC: American Psychiatric Association; 2013.
- Barneveld PS, Swaab H, Fagel S, van Engeland H, de Sonneville LM. Quality of life: a case-controlled long-term follow-up study, comparing young high-functioning adults with autism spectrum disorders with adults with other psychiatric disorders diagnosed in childhood. *Comprehensive Psychiatry*. 2014; 55:302–310. [PubMed: 24290884]
- Bellack AS, Morrison RL, Wixted JT, Mueser KT. An analysis of social competence in schizophrenia. *British Journal of Psychiatry*. 1990; 156:809–818. [PubMed: 2207511]

- Bishop-Fitzpatrick, L., Minshew, NJ., Eack, SM. A systematic review of psychosocial interventions for adults with autism spectrum disorders. In: Volkmar, FR.Reichow, B., McPartland, JC., editors. *Adolescents and Adults with Autism Spectrum Disorders*. New York: Springer; 2014. p. 315-327.
- Bishop DV. Development of the Children's Communication Checklist (CCC): a method for assessing qualitative aspects of communicative impairment in children. *Journal of Child Psychology and Psychiatry*. 1998; 39:879–891. [PubMed: 9758196]
- Bodfish JW, Symons FJ, Parker DE, Lewis MH. Varieties of repetitive behavior in autism: comparisons to mental retardation. *Journal of Autism and Developmental Disorders*. 2000; 30:237–243. [PubMed: 11055459]
- Chevallier C, Kohls G, Troiani V, Brodtkin ES, Schultz RT. The social motivation theory of autism. *Trends in Cognitive Sciences*. 2012; 16:231–239. [PubMed: 22425667]
- Constantino JN, Przybeck T, Friesen D, Todd RD. Reciprocal social behavior in children with and without pervasive developmental disorders. *Journal of Developmental and Behavioral Pediatrics*. 2000; 21:2–11. [PubMed: 10706343]
- Couture SM, Penn DL, Losh M, Adolphs R, Hurley R, Piven J. Comparison of social cognitive functioning in schizophrenia and high functioning autism: more convergence than divergence. *Psychological Medicine*. 2010; 40(4):569–579. [PubMed: 19671209]
- Craig JS, Hatton C, Craig FB, Bental RP. Persecutory beliefs, attributions and theory of mind: comparison of patients with paranoid delusions, Asperger's syndrome and healthy controls. *Schizophrenia Research*. 2004; 69(1):29–33. [PubMed: 15145468]
- Crespi B, Badcock C. Psychosis and autism as diametrical disorders of the social brain. *Behavioral Brain Sciences*. 2008; 31(3):241–261. [PubMed: 18578904]
- Crespi B, Stead P, Elliot M. Comparative genomics of autism and schizophrenia. *Proceedings of the National Academy of Sciences of the United States of America*. 2010; 107:1736–1741. [PubMed: 19955444]
- Eack SM, Newhill CE. Psychiatric symptoms and quality of life in schizophrenia: a meta-analysis. *Schizophrenia Bulletin*. 2007; 33:1225–1237. [PubMed: 17204532]
- Eaves LC, Ho HH. Young adult outcome of autism spectrum disorders. *Journal of Autism and Developmental Disorders*. 2008; 38(4):739–747. [PubMed: 17764027]
- Freeman D. Suspicious minds: the psychology of persecutory delusions. *Clinical Psychology Review*. 2007; 27(4):425–457. [PubMed: 17258852]
- Green MF, Kern RS, Heaton RK. Longitudinal studies of cognition and functional outcome in schizophrenia: implications for MATRICS. *Schizophrenia Research*. 2004; 72:41–51. [PubMed: 15531406]
- Horan WP, Subotnik KL, Snyder KS, Nuechterlein KH. Do recent-onset schizophrenia patients experience a “social network crisis”? *Psychiatry*. 2006; 69:115–129. [PubMed: 16822191]
- Howlin P, Goode S, Hutton J, Rutter M. Adult outcome for children with autism. *Journal of Child Psychology and Psychiatry*. 2004; 45:212–229. [PubMed: 14982237]
- Kay SR, Flszbein A, Opfer LA. The Positive and Negative Syndrome Scale (PANSS) for schizophrenia. *Schizophrenia Bulletin*. 1987; 13:261–276. [PubMed: 3616518]
- Klin A. Attributing social meaning to ambiguous visual stimuli in higher- functioning autism and Asperger syndrome: the social attribution task. *Journal of Child Psychology and Psychiatry*. 2000; 41(7):831–846. [PubMed: 11079426]
- Klin A, Volkmar FR, Sparrow SS. Autistic social dysfunction: some limitations of the theory of mind hypothesis. *Journal of Child Psychology and Psychiatry*. 1992; 33:861–876. [PubMed: 1378847]
- Konstantareas MM, Hewitt T. Autistic disorder and schizophrenia: Diagnostic overlaps. *Journal of Autism and Developmental Disorders*. 2001; 31(1):19–28. [PubMed: 11439750]
- Lam KS, Bodfish JW, Piven J. Evidence for three subtypes of repetitive behavior in autism that differ in familiarity and association with other symptoms. *Journal of Child Psychology and Psychiatry*. 2008; 49:1193–1200. [PubMed: 19017031]
- Laugeson, EA., Ellingsen, R. Social skills training for adolescents and adults with autism spectrum disorder. In: Volkmar, FR.Reichow, B., McPartland, JC., editors. *Adolescents and Adults with Autism Spectrum Disorders*. New York: Springer; 2014. p. 61-85.

- Leifker FR, Bowie CR, Harvey PD. Determinants of everyday outcomes in schizophrenia: the influences of cognitive impairment, functional capacity, and symptoms. *Schizophrenia Research*. 2009; 115:82–87. [PubMed: 19775869]
- Lord C, Risi S, Lambrecht L, Cook EH Jr, Leventhal BL, DiLavore PC, et al. The Autism Diagnostic Observation Schedule—Generic: a standard measure of social and communication deficits associated with the spectrum of autism. *Journal of Autism and Developmental Disorders*. 2000; 30:205–223. [PubMed: 11055457]
- Mandell DS, Lawer LJ, Branch K, Brodtkin ES, Healey K, Witalec R, et al. Prevalence and correlates of autism in a state psychiatric hospital. *Autism*. 2012; 16:557–567. [PubMed: 21846667]
- Marwaha S, Johnson S. Schizophrenia and employment. *Social Psychiatry Psychiatric Epidemiology*. 2004; 39:337–349. [PubMed: 15133589]
- Mueser, KT., Bellack, AS. Social skills and social functioning. In: Mueser, KT., Tarrrier, N., editors. *Handbook of Social Functioning in Schizophrenia*. Needham Heights: Allyn & Bacon; 1998. p. 79-96.
- Mueser KT, Bellack AS, Morrison RL, Wixted JT. Social competence in schizophrenia: premorbid adjustment, social skill, and domains of functioning. *Journal of Psychiatric Research*. 1990; 24:51–63.
- Nadig A, Lee I, Singh L, Bosshart K, Ozonoff S. How does the topic of conversation affect verbal exchange and eye gaze? A comparison between typical development and high-functioning autism. *Neuropsychologia*. 2010; 48:2730–2739. [PubMed: 20493890]
- Nicpon MF, Doobay AF, Assouline SG. Parent, teacher, and self perceptions of psychosocial functioning in intellectually gifted children and adolescents with autism spectrum disorder. *Journal of Autism and Developmental Disorders*. 2010; 40:1028–1038. [PubMed: 20143145]
- Patterson TL, Moscona S, McKibbin CL, Davidson K, Jeste DV. Social skills performance assessment among older patients with schizophrenia. *Schizophrenia Research*. 2001; 48:351–360. [PubMed: 11295387]
- Paul R, Orlovski SM, Marcinko HC, Volkmar F. Conversational behaviors in youth with high-functioning ASD and Asperger syndrome. *Journal of Autism and Developmental Disorders*. 2009; 39:115–125. [PubMed: 18607708]
- Pinkham AE, Penn DL. Neurocognitive and social cognitive predictors of interpersonal skill in schizophrenia. *Psychiatry Research*. 2006; 143:167–178. [PubMed: 16859754]
- Pinkham AE, Hopfinger JB, Pelphey KA, Piven J, Penn DL. Neural bases for impaired social cognition in schizophrenia and autism spectrum disorders. *Schizophrenia Research*. 2008; 99:164–175. [PubMed: 18053686]
- Pinkham, AE., Mueser, KT., Penn, DL., Glyn, SM., McGurk, SR., Addington, J. Social and functional impairments. In: Lieberman, JA. Stroup, TS., Perkins, DO., editors. *Essentials of Schizophrenia*. Arlington: American Psychiatric Publishing Inc; 2012. p. 93-130.
- Pinkham AE, Sasson NJ, Beaton D, Abdi H, Kohler CG, Penn DL. Qualitatively distinct factors contribute to elevated rates of paranoia in autism and schizophrenia. *Journal of Abnormal Psychology*. 2012; 121(3):767–777. [PubMed: 22686868]
- Rapoport J, Chavez A, Greenstein D, Addington A, Gogtay N. Autism spectrum disorders and childhood-onset schizophrenia: clinical and biological contributions to a relation revisited. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2009; 48:10–18. [PubMed: 19218893]
- Sasson NJ, Nowlin RB, Pinkham AE. Social cognition, social skill, and the broad autism phenotype. *Autism*. 2013; 17:655–667. [PubMed: 22987889]
- Sasson NJ, Pinkham AE, Carpenter KL, Belger A. The benefit of directly comparing autism and schizophrenia for revealing mechanisms of social cognitive impairment. *Journal of Neurodevelopmental Disorders*. 2011; 3:87–100. [PubMed: 21484194]
- Sasson NJ, Pinkham AE, Weittenhiller LP, Faso DJ, Simpson C. Context effects on facial affect recognition in schizophrenia and autism: behavioral and eye-tracking evidence. *Schizophrenia Bulletin*. 2015; 42:675–683. [PubMed: 26645375]

- Sasson N, Tsuchiya N, Hurley R, Couture SM, Penn DL, Adolphs R, et al. Orienting to social stimuli differentiates social cognitive impairment in autism and schizophrenia. *Neuropsychologia*. 2007; 45(11):2580–2588. [PubMed: 17459428]
- Saulnier CA, Klin A. Brief report: Social and communication abilities and disabilities in higher functioning individuals with autism and Asperger syndrome. *Journal of Autism and Developmental Disorders*. 2007; 37(4):788–793.
- Seltzer MM, Shattuck P, Abbeduto L, Greenberg JS. Trajectory of development in adolescents and adults with autism. *Mental Retardation and Developmental Disabilities Research Reviews*. 2004; 10:234–247. [PubMed: 15666341]
- Sherman RA, Funder DC. Evaluating correlations in studies of personality and behavior: beyond the number of significant findings to be expected by chance. *Journal of Research in Personality*. 2009; 43:1053–1063.
- Sullivan PF, Magnusson C, Reichenberg A, Boman M, Dalman C, Davidson M, et al. Family history of schizophrenia and bipolar disorder as risk factors for autism. *Archives of General Psychiatry*. 2012; 69:1099–1103. [PubMed: 22752149]
- Tabachnick, BG., Fidell, LS. Discriminant analysis. In: Tabachnick, BG., Fidell, LS., editors. *Using Multivariate Statistics*, 5e. Needham Heights: Allyn & Bacon; 2007. p. 375-436.
- Taylor JL, Henninger NA, Mailick MR. Longitudinal patterns of employment and postsecondary education for adults with autism and average-range IQ. *Autism*. 2015; 19:785–793. [PubMed: 26019306]
- Turner-Brown LM, Perry TD, Dichter GS, Bodfish JW, Penn DL. Brief report: feasibility of social cognition and interaction training for adults with high functioning autism. *Journal of Autism and Developmental Disorders*. 2008; 38:1777–1784. [PubMed: 18246419]
- Verhoeven EW, Smeekens I, Didden R. Brief report: suitability of the Social Skills Performance Assessment (SSPA) for the assessment of social skills in adults with autism spectrum disorders. *Journal of Autism and Developmental Disorders*. 2013; 43:2990–2996. [PubMed: 23624879]
- Wilkinson, G. *WRAT-3 Wide Range Achievement Test Administration Manual*, 3e. Wilmington: Wide Range Inc; 1993.
- Wykes T, Steel C, Everitt B, Tarrier N. Cognitive behavior therapy for schizophrenia: effect sizes, clinical models, and methodological rigor. *Schizophrenia Bulletin*. 2008; 34:523–537. [PubMed: 17962231]




Figure 1.

Table 1

Sample Characteristics

	ASD (n=54)		SCZ (n=54)		TD (n=56) ^a		χ^2	p
	n(%)	n(%)	n(%)	n(%)	n(%)	n(%)		
Gender							--	--
Male	47(87%)		47(87%)		49(88%)			
Ethnicity							10.11	.120
Caucasian	48(89%)		43(80%)		51(91%)			
African American	1(2%)		6(11%)		2(4%)			
Asian	5(9%)		3(5%)		3(5%)			
Other	--		2(4%)		--			
Medication							49.31	<.001
Atypical Only	11(20%)		41(76%)		--			
Typical Only	1(2%)		3(6%)		--			
Both	--		4(7%)		--			
	<i>Mean(SD)</i>	<i>Mean(SD)</i>	<i>Mean(SD)</i>	<i>Mean(SD)</i>	<i>Mean(SD)</i>	<i>F</i>	<i>p</i>	
Age	25.67(7.17)	28.67(10.11)	26.89(9.23)	1.53	.221			
Education (years)	13.89*1.78)	13.46(1.99)	13.80(1.68)	0.83	.439			
WRAT IQ	106.02(12.83)	103.32(10.98)	105.34(11.10)	0.79	.457			
PANSS								
Positive Total	8.94(3.16)	15.59(4.99)	--	68.45	<.001			
Negative Total	11.07(4.35)	14.50(4.60)	--	15.82	<.001			
General Total	24.02(5.46)	33.00(6.99)	--	55.33	<.001			

^aTD controls did not complete PANSS and were not taking any psychotropic medications.

Abbreviations: *SD*= Standard deviation; WRAT = Wide Range Achievement Test; PANSS = Positive and Negative Syndrome Scale.

Table 2

Factor Analysis Pattern Matrix for Verbal Content, Nonverbal, Paralinguistic, and Interactive Measures of Social Behavior^a

	Factor 1	Factor 2	Factor 3	Factor 4
Social Behavior	Paralinguistic	Nonverbal	Interactive	Verbal Content
Meshing	.86	.12	-.12	-.08
Fluency	.68	-.12	.05	.13
Clarity	.45	.07	.08	.10
Flat Affect	.06	.89	-.08	-.18
Appropriate Affect	-.11	.67	-.05	.33
Gaze	.06	.47	.08	.17
Asks Questions	-.05	-.10	.96	-.02
Involvement	.16	.35	.49	-.03
Verbal Content	.07	-.06	-.03	.90

^a*N* = 164. Principal Axis Factoring with Promax rotation.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Reliability Coefficients, and Univariate Tests for Social Behaviors, and Cohen's *d* for Group Comparisons

Table 3

Social Behavior	ICC (n=30)		TD (n=56)		SCZ (n=54)		ASD (n=54)		<i>F</i> (2, 161)	<i>p</i>	TD v. ASD		TD v. ASD v. SCZ	
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>			<i>d</i>	<i>d</i>	<i>d</i>	<i>d</i>
Verbal Content	.87	7.01	0.74	6.28	1.37	6.04	1.18	11.27*	<.001	0.9 [†]	1.31 [†]	-0.32		
Clarity	.84	6.24	1.12	5.54	1.15	5.86	1.27	4.92	.008	0.63	0.34	0.29		
Fluency	.84	6.18	1.16	5.27	1.14	4.97	1.49	13.55*	<.001	0.78 [†]	1.04 [†]	-0.26		
Meshing	.92	5.70	1.37	5.29	1.39	5.23	1.56	1.72	.183	0.30	0.34	-0.04		
Gaze	.95	7.22	1.32	6.03	1.59	5.94	1.83	11.17*	<.001	0.90 [†]	0.97 [†]	-0.07		
Involvement	.93	6.64	1.17	6.07	1.35	5.58	1.51	8.53*	<.001	0.49	0.91 [†]	-0.42		
Asks Questions	.99	5.61	2.94	4.29	2.77	2.94	1.97	14.50*	<.001	0.45 [†]	0.91 [†]	-0.46 [†]		
Appropriate Affect	.66	7.09	0.55	6.78	0.66	6.65	0.83	6.04*	<.003	0.56 [†]	0.80 [†]	-0.24		
Flat Affect	.88	6.70	0.77	6.16	1.05	6.34	0.95	4.84	.009	0.70	0.47	0.23		
Social Anxiety	.74	6.24	1.30	5.38	1.18	5.07	1.43	11.86*	<.001	0.66 [†]	0.90 [†]	-0.24		
Repetitive Verbal Content	.80	7.21	0.74	6.88	1.03	6.65	0.98	5.22	.006	0.45	0.76	-0.31		
Repetitive Movement	.84	7.68	1.01	7.51	1.16	6.69	1.33	11.26*	<.001	0.17	0.98 [†]	-0.81 [†]		
Verbosity	.96	5.30	1.6	4.52	1.69	4.63	1.42	4.02	.020	0.49	0.42	0.07		
Overall Social Skill	.95	6.70	1.07	5.65	1.58	5.14	1.59	17.01*	<.001	0.98 [†]	1.46 [†]	-0.48		

Note Cohen's *d* calculated with the standard deviation of the TD group used as the standardizer. Repetitive Verbal Content and Repetitive Movement have been reversed coded for consistency.

ICC = Intra-class correlation coefficient; *d* = Cohen's *d*; *M* = Mean; *SD* = Standard deviation.

* Significant at Bonferroni corrected alpha, *p* < 0.0035.

[†] Tukey post-hoc test significant at *p* < .05.

Table 4

Standardized Coefficients for Social Behaviors for Discriminant Functions 1 and 2. Coefficients greater than .40 load onto the function.

	Discriminant Function	
	1	2
Interactive	.75	-.48
Repetitive Movement	.64	-.46
Verbal Content	.54	.14
Nonverbal	-.09	.71
Paralinguistic	-.19	.30
Verbosity	<.01	.35
Repetitive Verbal Content	.03	-.11

Note: Repetitive Verbal Content and Repetitive Movement have been reversed coded for consistency.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 5

Bivariate Correlations of Social Behaviors with IQ for Each Group

Social Behavior	ASD (n = 54)	SCZ (n = 54)	TD (n = 56)
	<i>r</i>	<i>r</i>	<i>r</i>
Verbal Content	.02	.33 ^a	.02
Clarity	.25	.39 ^b	.25
Fluency	-.13	.35 ^a	.04
Meshing	.03	.29 ^a	.04
Gaze	.02	.24	.17
Involvement	.26	.33 ^a	.32 ^a
Asks Questions	.18	.17	.47 ^b
Appropriate Affect	.16	.15	.29 ^a
Flat Affect	.22	.12	.13
Social Anxiety	.01	.31 ^a	.20
Repetitive Verbal Content	.15	.12	.06
Repetitive Movement	-.12	-.25	.06
Verbosity	.37 ^b	.25	.35 ^b
Overall Social Skill	.23	.42 ^b	.27 ^a

^a*p* < .05.^b*p* < .01.

Note: Repetitive Verbal Content and Repetitive Movement have been reversed coded for consistency.