

Research Bank

Journal article

Repeated high-intensity running and sprinting in elite women's soccer competition

Gabbett, Tim J., Wiig, Håvard and Spencer, Matt

Accepted author manuscript version reprinted, by permission, from *International Journal of Sports Physiology and Performance*, 2013, 8 (2): 130-138, <https://doi.org/10.1123/ijssp.8.2.130>. © 2013 Human Kinetics, Inc.

1 Running title: Repeated high-intensity exercise in soccer

2
3 **Repeated High-Intensity Running and Sprinting in Elite Women's Soccer Competition**

4
5
6 Tim J. Gabbett

7 *School of Exercise Science, Australian Catholic University, Brisbane, Australia*
8 *School of Human Movement Studies, The University of Queensland, Brisbane, Australia*

9
10
11 Håvard Wiig

12 *Norwegian Research Centre for Training and Performance, Norwegian School of Sports*
13 *Sciences, Oslo, Norway*

14
15
16 Matt Spencer

17 *Norwegian Research Centre for Training and Performance, Norwegian School of Sports*
18 *Sciences, Oslo, Norway*

19
20
21 Address correspondence to:

22 Dr. Tim J. Gabbett

23 School of Exercise Science,
24 Australian Catholic University,

25 Brisbane, AUSTRALIA 4014

26 Email: tim_gabbett@yahoo.com.au

27
28
29 *Submission Type:* Original Investigation

30 *Abstract Word Count:* 250 Words

31
32
33 *Text-Only Word Count:* 4014 Words

34
35 *Number of Figures and Tables:* 4 Figures, 2 Tables

36

37 **Abstract**

38 **Background:** To our knowledge, no study has investigated the concurrent repeated, high-
39 intensity (RHIA) and repeated-sprint activity (RSA) of intermittent team sport competition.

40

41 **Purpose:** In this study, we report on the RSA of elite Women's football competition. In
42 addition, we describe the nature of RHIA (e.g. striding and sprinting activities) that involve a
43 high energy cost and are associated with short duration (i.e. ≤ 20 seconds) recovery periods.

44

45 **Methods:** Thirteen elite women soccer players underwent video-based time-motion analysis
46 on 34 occasions during national and international standard matches. RSA and RHIA were
47 defined as successive (i.e. 2) sprints, or striding and sprinting efforts that occurred with ≤ 20
48 seconds between efforts.

49

50 **Results:** The number of RSA and RHIA bouts performed was similar between the first and
51 second half of matches. Sprinting and striding/sprinting durations tended to remain relatively
52 stable irrespective of the number of efforts in a RSA or RHIA bout, or the period of play.
53 However, recovery duration between efforts increased in the second half, and when a greater
54 number of efforts were performed per bout.

55

56 **Conclusion:** These findings suggest that first to second half reductions in RHIA and RSA do
57 not occur in elite Women's soccer competition. However, players increase the amount of
58 low-intensity recovery undertaken between RHIA and RSA efforts, most likely in an attempt
59 to maintain RHIA and RSA performance. These findings emphasize the importance of
60 repeated-sprint and effort ability to elite Women's soccer, and highlight the importance of
61 training this quality to prevent reductions in performance during competitive match-play.

62

63 **Keywords:** time-motion analysis, high-speed running, repeated-sprint ability, training, team
64 sports

65

66

67 **Introduction**

68 Repeated-sprint and prolonged high-intensity running ability are widely accepted as critical
69 components of high-intensity intermittent team sports (e.g. soccer). In soccer competition it
70 has been shown that periods of fatigue follow the most intense bouts of high-intensity
71 running.¹ Reports on the relative importance of high-intensity running to elite male and
72 female soccer performance are equivocal, with some,^{2,3} but not all^{4,5} studies demonstrating
73 that elite players perform more high-intensity running than sub-elite competitors, and when
74 the demands of competition are greatest. The number and intensity of repeated-sprint bouts
75 has also been reported to be a significant discriminator of elite and sub-elite female soccer
76 competition.⁶ In a study of female soccer players, Gabbett and Mulvey⁶ demonstrated that
77 players performed on average, more repeated-sprint bouts in international matches (4.8 bouts
78 per match) than national (1.0 bout per match) and domestic (1.4 bouts per match) matches.
79 Importantly, the average number of sprints was greater, and recovery between sprints shorter
80 in international competition. Collectively, these results suggest that repeated-sprint activity
81 and the amount of high-intensity running performed may differ according to the competitive
82 standard.

83
84 Despite the importance of high-intensity running^{1,2} and repeated-sprint ability⁶⁻⁸ to
85 competitive success in high-intensity intermittent team sports, studies investigating the nature
86 of repeated high-intensity activity in these sports are limited.⁹⁻¹² Spencer et al.⁷ subjectively
87 assessed the sprint and repeated-sprint demands of elite field hockey using video time-motion
88 analysis. The authors reported that repeated-sprint bouts (defined as a minimum of 3 sprints,
89 with ≤ 20 seconds recovery between sprints) occurred on 17 occasions throughout the match.
90 The average number of sprints within the repeated-sprint bouts was 4 (± 1), with a mean
91 recovery time of 14.9 seconds between sprint efforts. It has been suggested that while
92 repeated-sprints occur infrequently, the ability or inability to perform these activities may
93 prove critical to the outcome of the match,⁷ although to date, limited evidence exists to
94 support this claim.⁸

95
96 Although repeated-sprint activity provides an indication of the most demanding passages
97 experienced during match-play, operational definitions of repeated-sprint ability (i.e. 3 or
98 more sprints with ≤ 20 seconds recovery between sprints) may only provide a limited picture
99 of the physically demanding running activities that occur in high-intensity, intermittent team
100 sports. For example, repeated-sprint bouts may include a range of sprint effort frequencies,
101 with relatively long or short recovery between efforts; reporting of the average demands
102 alone may underestimate the most extreme demands that might be expected during
103 competition. In addition, limiting repeated-sprint bouts to 3 or more sprints, effectively
104 eliminates successive, short-recovery sprints that may also be physically demanding but fail
105 to meet the traditional repeated-sprint criteria.⁷ Equally, high-speed running (i.e. striding)
106 efforts, which may also be separated by short recovery periods, are likely to make a
107 substantial contribution to the energetic cost of competition, despite failing to qualify as
108 repeated-sprint activity. An understanding of these repeated, high-intensity activity bouts
109 would provide strength and conditioning coaches with evidence to inform conditioning
110 programs in order to adequately prepare players for the repeated high-intensity (both striding
111 and sprinting) demands of competition.

112
113 To our knowledge, no study has investigated the concurrent repeated, high-intensity activity
114 (i.e. striding and sprinting) and repeated-sprint demands of intermittent team sport
115 competition. In this study, we report on the repeated-sprint demands of elite Women's soccer
116 competition, with special reference to the number of sprints within a bout, and the typical

117 sprint and recovery durations performed within these high-intensity activities. In addition,
118 we describe the nature of repeated, high-intensity activities (e.g. striding and sprinting) that
119 involve a high energy cost and are associated with short duration (i.e. ≤ 20 seconds) recovery
120 periods, in order to document repeated high and very-high intensity exercise bouts in this
121 sport.

122

123 **Methods**

124 *Subjects*

125 Thirteen elite women soccer players (mean \pm SD age, 21 ± 2 years) participated in this study.
126 Athletes were scholarship holders with the Queensland Academy of Sport women's soccer
127 program and/or members of the Australian Matildas women's soccer team. All participants
128 received a clear explanation of the study, including the risks and benefits of participation, and
129 written consent was obtained. The institutional review board for human investigation
130 approved all experimental procedures.

131

132 *Time-Motion Analysis*

133 Video-based time-motion analysis was performed during 10 national and 5 international
134 standard competitive matches. Players were filmed and analyzed on 34 occasions (19
135 national and 15 international player appearances) over the course of the 15 matches. The
136 mean \pm SD number of national and international matches analyzed per player was 2.7 ± 1.1
137 and 2.1 ± 1.5 , respectively. All matches were 90 minutes in duration, and played on a full-
138 sized pitch (100 m length x 65 m width). Only outfield players (i.e., strikers, mid-fielders,
139 and defenders) were filmed. No substitutes were filmed or analyzed in this study. All
140 comparisons between the first and second half were made on the same player. Video
141 recordings were made using digital video cameras (Sony, DCR-TRV 950E). Cameras were
142 positioned in the stadium, on the half-way line, approximately 30 m above the field of play to
143 cover the entire playing arena. The zoom function of the video camera was utilized during
144 recording so that the image of the player and a 10 m radius of her surrounds was maintained.
145 Up to four cameras were used in any given match, and players had no knowledge of who was
146 being filmed for each match.

147

148 The video recordings were analyzed by one experienced observer by logging frequency and
149 duration of the activities performed using purpose built software (Gab-Trakka, Brisbane,
150 Australia). A description of all match-play activities is provided in Table 1. The initiation
151 and completion of each individual activity were recorded, and the duration of each activity
152 was calculated. The time spent standing, walking, and jogging was considered to be low-
153 intensity exercise, with striding and sprinting regarded as high-intensity exercise. The typical
154 error of measurement (% coefficient of variation, CV) for test-retest reliability for the
155 activities of standing, walking, jogging, striding, and sprinting were 0.6%, 0.3%, 2.4%, 4.6%,
156 and 3.5%, respectively.

157

158

Insert Table 1 About Here

159

160 *Repeated-Sprint and Repeated High-Intensity Activity*

161 Repeated-sprint activity (RSA) was defined in two ways. First, RSA was defined as a
162 minimum of three sprints, with ≤ 20 seconds between sprints.⁷ The mean, maximal, and
163 minimal duration of sprints, number of sprint repetitions, and recovery duration were also
164 recorded. To account for physically demanding repeated sprints that did not meet the
165 traditional repeated-sprint definition, we also recorded successive (i.e. 2) sprints that occurred
166 with ≤ 20 seconds between sprints. We also adapted this definition to record the frequency of

167 repeated high-intensity activity (i.e. a combination of striding and/or sprinting; RHIA) bouts.
168 RHIA was defined as a minimum of two consecutive sprints or striding efforts, with ≤ 20
169 seconds between efforts.

170 171 *Statistical Analysis*

172 Differences in RSA and RHIA (striding and/or sprinting) demands between the first and
173 second half of matches were analyzed using a practical approach based on the real world
174 relevance of the results.¹³ Differences in the RSA and RHIA demands between the first and
175 second half of matches, and national and international competition were analyzed using
176 Cohen's effect size (ES) statistic and 90% confidence limits (CL). Effect sizes of <0.2 , 0.2 -
177 0.6 , 0.6 - 1.2 , 1.2 - 2.0 , and >2.0 were considered trivial, small, moderate, large, and very large,
178 respectively.¹³ Given our small sample size and that a wide between-match variability has
179 been shown for the amount of high-speed running and sprinting performed in soccer,¹⁴ a
180 moderate effect size was chosen as our lower limit for observed differences. All data are
181 reported as mean \pm SD.

182 183 **Results**

184 The number of repeated-sprint and striding actions per bout when analyzed as sets of 2, 3, 4,
185 5, or 6 efforts are shown in Table 2. In general, there was wide variability in the number of
186 RSA bouts performed among the different players, with 5.1 ± 5.1 (range = 0-23) RSA bouts
187 performed per player per match. Sprint bouts consisting of successive sprints (i.e. 2 sprint
188 efforts) were the most common type of RSA bout. Some players ($N = 6$, 17.7%) performed
189 no repeated-sprint bouts, and others performed up to 23 repeated-sprint bouts in a match.
190 The greatest number of sprints in any single bout was 7. The average sprint duration
191 throughout the various RSA bouts was 2.17 ± 0.13 s, with sprint duration remaining
192 relatively stable across short (i.e. 2 sprints; 2.27 ± 0.56 s) and long (i.e. 6 sprints; 2.08 ± 0.19
193 s) RSA bouts. However, the mean recovery duration progressively increased with a greater
194 number of sprints per bout.

195
196 RHIA bouts that included a combination of striding and/or sprinting were more common than
197 RSA bouts; players performed on average 31.2 ± 18.7 RHIA bouts per game. The most
198 common type of RHIA bout involved 2 efforts. Repeated striding and sprinting activities that
199 involved 6 efforts occurred on 11.1 ± 13.8 occasions per player, per game. The average
200 stride and sprint duration throughout the various RHIA exercise bouts was 2.94 ± 0.05 s, with
201 sprinting and striding durations remaining relatively stable across short (i.e. 2 sprint/stride
202 efforts; 2.98 ± 0.66 s) and long (i.e. 6 sprint/stride efforts; 2.93 ± 0.67 s) RHIA exercise
203 bouts. Consistent with RSA bouts, the mean recovery duration between sprinting and
204 striding efforts progressively increased with a greater number of efforts per bout.

205
206 When analyzed as sets of 6 efforts, the average sprint duration in RSA bouts of international
207 matches was greater ($26.0 \pm 24.2\%$, $ES = 2.48$) than national matches. No other meaningful
208 differences ($ES = 0.03$ - 0.50) were found between national and international matches for
209 average sprint duration in RSA bouts. No meaningful differences ($ES = 0.08$ - 0.29) were
210 detected between national and international matches for the average effort duration of RHIA
211 bouts.

212
213 The recovery duration of sets of 4 RSA bouts was moderately shorter ($22.0 \pm 20.3\%$, $ES =$
214 0.64) in international than national matches. However, only trivial to small differences ($ES =$
215 0.02 - 0.40) were observed between national and international matches for recovery durations
216 between sprinting/striding efforts in RHIA exercise bouts.

217
218
219

Insert Table 2 About Here

220 While the small sample size limited our analysis of positional differences, the number of
221 RSA bouts performed was generally higher in central mid-fielders (7.4 ± 4.5 national, $N = 5$;
222 10.0 ± 11.3 international, $N = 3$) than wide mid-fielders (1.0 ± 1.0 national, $N = 3$; 6
223 international, $N = 1$), strikers (6.0 ± 6.7 national, $N = 5$; 3.6 ± 2.3 international, $N = 5$), and
224 defenders (5.3 ± 4.4 national, $N = 6$; 3.2 ± 2.3 international, $N = 6$). The number of RHIA
225 bouts performed was also generally higher in central mid-fielders (37.8 ± 19.7 national, $N =$
226 5 ; 39.7 ± 35.2 international, $N = 3$) than wide mid-fielders (24.7 ± 9.1 national, $N = 3$; 6
227 international, $N = 1$), strikers (42.2 ± 20.6 national, $N = 5$; 26.8 ± 9.2 international, $N = 5$),
228 and defenders (32.1 ± 21.1 national, $N = 6$; 20.7 ± 13.6 international, $N = 6$).

229

230 Possibly small differences (30%, $CL \pm 31\%$, $ES = 0.29$) were found between the first and
231 second half of matches for the number of RSA (3.0 ± 3.2 vs. 2.1 ± 2.8) bouts performed. In
232 addition, there was a possibly small decrease ($14\% \pm 11\%$, $ES = 0.22$) in the number of
233 RHIA bouts from the first (16.8 ± 11.3) to the second half (14.4 ± 10.0) of matches. In
234 general, there was a greater frequency of RSA and RHIA bouts in the first 15 minutes of the
235 match, with the frequency of RSA and RHIA bouts declining over the course of the match for
236 both national and international players (Figure 1). After the half-time break there was a trend
237 towards an increased number of RSA and RHIA bouts during the 45-60 min period, when
238 compared to the two previous and subsequent periods.

239

Insert Figure 1 About Here

240

241
242 Figure 2 shows the first and second half recovery durations per RSA and RHIA bout, when
243 analyzed as sets of 2, 3, 4, 5 or 6 efforts. RSA bouts were associated with very likely greater
244 recovery durations between efforts for sets of 4 ($22.3 \pm 18.5\%$, $ES = 0.67$) and 5 ($26.2 \pm$
245 5.7% , $ES = 0.93$) sprints, in the second half compared to the first half of matches.
246 Approximately 88-90% of motion involved low-intensity activity. The time spent standing
247 (5.4% vs. 8.5%), walking (36.5% vs. 30.4%), jogging (47.6% vs. 49.8%), and striding (10.4%
248 vs. 11.3%) between RSA bouts changed differentially from the first to second half. Only
249 trivial to small differences ($ES = 0.01-0.33$) were observed between the first and second half
250 of matches for recovery durations between sprinting/striding efforts in RHIA exercise bouts.
251 The percentage of time spent standing, walking, and jogging between RHIA bouts in the first
252 and second half was 7.0% and 7.2% , 39.4% and 44.6% , and 53.6% and 48.2% , respectively.

253

Insert Figure 2 About Here

254

255
256 Figure 3 shows the percentage change in recovery duration between RSA bouts and RHIA
257 bouts, expressed relative to sets of 2 RSA or RHIA exercise bouts. Moderate ($ES = 0.6-1.2$)
258 to large ($ES = 1.2-2.0$) increases in recovery durations were observed for both RSA and
259 RHIA bouts as the number of efforts in the respective bouts increased. In comparison to
260 RHIA bouts, larger recovery durations were observed in RSA bouts when sprint activity
261 increased to 6 efforts per bout ($11.4 \pm 6.8\%$, $ES = 1.2$).

262

Insert Figure 3 About Here

263

264
265 Figure 4 shows the percentage change in recovery duration between RSA bouts and RHIA
266 bouts for national and international matches, expressed relative to sets of 2 RSA or RHIA

267 exercise bouts. Very large differences in recovery durations between national and
268 international matches were observed for RSA bouts involving 3 ($23.5 \pm 3.8\%$, ES = 2.33) and
269 4 ($32.9 \pm 5.8\%$, ES = 3.01) efforts, with national matches associated with greater recovery
270 durations than international matches. Conversely, the recovery duration between RHIA
271 efforts for international matches was greater than national matches for RHIA bouts involving
272 3 ($14.5 \pm 5.0\%$, ES = 2.15), 4 ($11.8 \pm 1.4\%$, ES = 1.21), and 5 ($11.2 \pm 1.7\%$, ES = 0.89)
273 efforts.

274
275 *Insert Figure 4 About Here*
276

277 **Discussion**

278 This study investigated the repeated-sprinting and high-intensity exercise (e.g. striding and
279 sprinting) demands of elite Women's soccer match-play. The number of repeated-sprint and
280 repeated high-intensity exercise (i.e. striding and sprinting) bouts performed were similar
281 between the first and second half of matches. While sprint duration tended to remain
282 relatively stable, irrespective of the number of sprints in a repeated-sprint bout, or the period
283 of play (i.e. first or second half of matches), recovery between sprints increased in the second
284 half of matches, and when a greater number of efforts were performed per bout. These
285 findings suggest that first to second half reductions in repeated high-intensity exercise
286 activity and repeated-sprint activity do not occur in elite Women's soccer competition.
287 However, players increase the amount of low-intensity recovery undertaken between repeated
288 high-intensity exercise and repeated-sprint efforts, most likely in an attempt to maintain
289 repeated high-intensity activity and repeated-sprint performance.

290
291 The results of this study demonstrate large individual variations in the repeated-sprint
292 demands of elite Women's soccer match-play, with some players performing no repeated-
293 sprint bouts, and others performing up to 23 repeated-sprint bouts in a match. While our
294 sample size prevented a detailed analysis of the repeated-sprint and high-intensity activity
295 demands of different playing positions, it is likely that this variability is explained by the
296 different positional demands experienced in soccer.⁵ The number of sprints also varied
297 considerably, with as few as 2 sprints in a bout (53.5%), and as many as 7 sprints (0.6%) in a
298 bout. The player with the single highest number of repeated-sprint bouts in a match
299 performed 23 repeated-sprints bouts, of which 9 bouts involved 4 sprints, 6 bouts involved 5
300 sprints, 4 bouts involved 6 sprints, and 2 bouts involved as many as 7 sprints. These findings
301 highlight the stochastic nature of Women's soccer, while also emphasizing the highly
302 variable nature of repeated-sprint activity in this sport. Clearly, conditioning programs
303 designed to improve repeated-sprint ability should focus on bouts with varying sprint
304 frequencies and recovery durations that mimic the intense intermittent and unplanned nature
305 of sprinting in the sport. Moreover, individualized and position-specific conditioning
306 programs may be necessary for some players in order to adequately prepare them for the
307 large amounts of repeated-sprint and repeated high-intensity activity required in competition.

308
309 We found that the average number of repeated-sprint bouts and mean sprint duration was
310 maintained across the duration of the match, although consistent with previous studies of the
311 high-intensity running demands of elite soccer,^{1,3} the frequency of RSA and RHIA bouts in
312 the first 15 minutes of the match tended to be higher than during other 15 minute periods.
313 The finding of maintained repeated-sprint activity across the first and second half of matches
314 in the present study is consistent with the results from male soccer players; sprinting
315 performance was also maintained across the duration of a match, despite reductions in total
316 distance covered and distances covered in high-speed running.¹⁵ We also found that recovery

317 duration progressively increased with increases in the number of efforts performed per bout.
318 In addition, while mean sprint duration was relatively stable from the first to the second half
319 of the match, recovery durations between sprint efforts were longer in the second half. It has
320 previously been shown that points scored or conceded in elite team sport competition occur in
321 close proximity to a repeated effort bout,⁸ suggesting that the ability (or inability) to perform
322 repeated-sprint activity may prove critical to the outcome of the match. It is unclear if the
323 increase in recovery durations in the second half of matches, and with greater sprint
324 frequencies, observed in the present study represents conscious control by players. However,
325 these results may suggest a pacing strategy employed by players in an attempt to preserve
326 repeated-sprint performance.¹⁶

327
328 A novel aspect of this study was the expansion of previous repeated-sprint definitions to
329 include repeated high-intensity exercise (i.e. striding and sprinting) activities, and repeated,
330 successive (i.e. 2) sprints. Previous studies have defined repeated-sprint activity as 3 or more
331 sprints, with short (≤ 20 s) recovery between sprints.⁷ While this definition has provided a
332 consistent method of assessing repeated-sprint ability, limiting repeated-sprint bouts to 3 or
333 more sprints effectively eliminates successive, short-recovery sprints that may also be
334 physically demanding but fail to meet the traditional (i.e. 3 or more sprints) repeated-sprint
335 criteria. Equally, high-speed running (i.e. striding) efforts, which may also be separated by
336 short recovery periods, are likely to make a substantial contribution to the energetic cost of
337 competition, despite failing to qualify as repeated-sprint activity. Consistent with studies
338 from other high-intensity intermittent team sports (e.g. water polo),¹⁰ the number of repeated
339 high-intensity efforts increased considerably when considering repeated-sprint bouts that only
340 included 2 sprints. Furthermore, the number of repeated-effort bouts performed per player
341 increased from 5.1 to 31.2, when considering repeated-striding and sprinting, rather than
342 repeated-sprinting alone. However, rather than over-emphasizing the repeated-sprint
343 demands of competition, we believe these findings provide a more complete picture of the
344 repeated high-intensity effort activity that is required in elite Women's soccer match-play.
345 These findings provide specific information for the design of conditioning programs to meet
346 the repeated high-intensity exercise (i.e. striding and sprinting) demands of intermittent team
347 sports. Importantly, these findings consider activities that are physically demanding,
348 including those activities that do not involve sprinting.

349
350 In general, few differences were observed between national and international matches for the
351 durations of sprinting efforts (in RSA bouts), sprinting and striding efforts (in RHIA bouts),
352 and recovery between efforts (for both RSA and RHIA bouts). However, the average sprint
353 duration was longer in international matches (2.32 ± 0.15 s vs. 1.83 ± 0.24 s) when RSA
354 bouts consisted of 6 efforts, and the recovery duration between efforts was shorter in
355 international matches (11.97 ± 4.09 s vs. 14.60 ± 4.18 s) when RSA bouts consisted of 4
356 efforts. Of interest was the percentage change in recovery duration between RSA and RHIA
357 bouts for national and international matches when recovery durations were expressed relative
358 to sets of 2 RSA and RHIA exercise bouts. RSA bouts involving 3 and 4 efforts were
359 associated with greater recovery durations for national matches, while the recovery duration
360 between RHIA efforts for international matches was greater than national matches for RHIA
361 bouts involving 3, 4, and 5 efforts. While previous studies have investigated the physical
362 demands of soccer match-play, reports on the relative importance of high-intensity running to
363 soccer performance are equivocal, with some,^{2,3} but not all^{4,5} studies demonstrating that elite
364 players perform more high-intensity running than sub-elite competitors, and when the
365 demands of competition are greatest. Our results are generally consistent with the hypothesis
366 that the RSA demands are similar between national and international competition, although in

367 international matches the absolute sprint durations are slightly longer and absolute recovery
368 durations slightly shorter than national matches. Furthermore, in comparison to national
369 matches, elite female soccer players have higher relative recovery durations between RHIA
370 efforts in international matches, perhaps due to the relatively short recovery durations
371 between RSA efforts.

372
373 In this study, the physical demands of national and international female soccer matches were
374 assessed using video-based time-motion analysis. Due to the time-consuming and labour-
375 intensive nature of this methodology, our sample size is limited. Consequently, while the
376 RSA and RHIA demands tended to be greater in central mid-fielders, our ability to perform a
377 comprehensive analysis and make definitive conclusions on the differences in RSA and
378 RHIA among playing positions is restricted, and could be viewed as a limitation of this study.
379 While the use of global position system (GPS) technology would permit a much larger study
380 to be performed, including a detailed analysis of positional differences, current restrictions
381 applied by the International Federation of Association Football (FIFA) preventing players
382 from wearing microsensor technology in competition, may limit further progress in the field
383 of repeated-sprint and repeated high-intensity exercise activity. In addition, while it has
384 recently been shown that performance, physiological, and perceptual responses may be
385 influenced by the magnitude of directional change performed during repeated-sprint bouts,¹⁷
386 we made no attempt to quantify repeated change of direction activities. Although this
387 decision may have resulted in an underestimation of the repeated sprint and high-intensity
388 exercise demands of Women's soccer match-play, we reconciled this decision with the
389 knowledge that the coding of our other match-play activities was highly reproducible.

390
391 In conclusion, this study investigated the repeated-sprinting and high-intensity exercise (e.g.
392 striding and sprinting) demands of elite Women's soccer match-play. A major new finding
393 of this study was the manner in which repeated-sprint and repeated high-intensity exercise
394 (i.e. striding and sprinting) activity changed across the duration of a competitive match.
395 When comparing the repeated-sprint and repeated high-intensity exercise demands of the first
396 and second half, no differences were found for the number of repeated-sprint or repeated
397 high-intensity exercise bouts performed. Furthermore, the recovery between repeated-sprint
398 efforts increased from the first to the second half, while the recovery between repeated high-
399 intensity exercise efforts remained relatively stable. These findings suggest that reductions in
400 performance manifest in different ways for repeated-sprint and repeated high-intensity
401 exercise ability. It appears that repeated-sprint activity is protected by decreasing the amount
402 of repeated high-intensity exercise performed or by increasing the amount of low-intensity
403 recovery activity undertaken between repeated-sprints.

404 405 **Practical Applications**

406 This study described the unique repeated high-intensity exercise demands of elite Women's
407 soccer match-play. Repeated-sprint bouts involved as few as 2 sprint efforts, and as many as
408 7 sprint efforts. As the number of sprints performed per bout increased, the recovery
409 durations between sprints also increased. Despite the intense nature of the repeated-sprint
410 and repeated high-intensity exercise bouts, players were able to maintain repeated-effort
411 performance over the duration of the match. These findings can be used by coaches to
412 adequately prepare players for the repeated high-intensity (both striding and sprinting)
413 demands of competition. Critically, these results emphasize the importance of repeated-
414 sprint and effort ability to elite Women's soccer, and highlight the importance of training this
415 quality to prevent reductions in performance during competitive match-play.

416

417 **References**

- 418 1. Mohr, M, Krustup, P, Bangsbo, J. Fatigue in soccer: a brief review. *J. Sports. Sci.*
419 2005;23:593-599.
- 420 2. Mohr, M, Krustup, P, Andersson, H, Kirkendal, D, Bangsbo, J. Match activities of elite
421 women soccer players at different performance levels. *J. Strength. Cond. Res.* 2008;22:341-
422 349.
- 423 3. Andersson, HA, Randers, MB, Heiner-Moller, A, Krustup, P, Mohr, M. Elite female
424 soccer players perform more high-intensity running when playing in international games
425 compared with domestic league games. *J. Strength. Cond. Res.* 2010;24:912-919.
- 426 4. Rampinini, E, Impellizzeri, FM, Castagna, C, Coutts, AJ, Wisloff, U. Technical
427 performance during soccer matches of the Italian Serie A league: effect of fatigue and
428 competitive level. *J. Sci. Med. Sport.* 2009;12:227-233.
- 429 5. Di Salvo, V, Gregson, W, Atkinson, G, Tordoff, P, Drust, B. Analysis of high intensity
430 activity in Premier League soccer. *Int. J. Sports. Med.* 2009;30:205-212.
- 431 6. Gabbett, TJ, Mulvey, MJ. Time-motion analysis of small-sided training games and
432 competition in elite women soccer players. *J. Strength. Cond. Res.* 2008;22:543-552.
- 433 7. Spencer, M, Lawrence, S, Rechichi, C, Bishop, D, Goodman, C. Time-motion analysis of
434 elite field hockey, with special reference to repeated-sprint activity. *J. Sports. Sci.*
435 2004;22:843-850.
- 436 8. Austin, DJ, Gabbett, TJ, Jenkins, DG. Repeated high-intensity exercise in a professional
437 rugby league. *J. Strength. Cond. Res.* 2011;25:1898-1904.
- 438 9. Buchheit, M, Mendez-Villanueva, A, Simpson, BM, Pourdon, PC. Repeated-sprint
439 sequences during youth soccer matches. *Int. J. Sports. Med.* 2010;31:709-716.
- 440 10. D'Auria, S, Gabbett, T. A time-motion analysis of international women's water polo
441 match play. *Int. J. Sports. Physiol. Perform.* 2008;3:305-319.
- 442 11. Gabbett, TJ, Jenkins, DG, Abernethy, B. Physical demands of professional rugby league
443 training and competition using microtechnology. *J. Sci. Med. Sport.* 2012;15:80-86.
- 444 12. Tan, F, Polglaze, T, Dawson, B. Activity profiles and physical demands of elite women's
445 water polo match play. *J. Sports. Sci.* 2009;27:1095-1104.
- 446 13. Batterham, AM, Hopkins, WG. Making meaningful inferences about magnitudes. *Int. J.*
447 *Sports. Physiol. Perform.* 2006;1:50-57.
- 448 14. Gregson, W, Drust, B, Atkinson, G, Di Salvo, V. Match-to-match variability of high-
449 speed activities in Premier League soccer. *Int. J. Sports. Med.* 2010;31:237-242.
- 450 15. Weston, M, Drust, B, Gregson, W. Intensities of exercise during match-play in FA
451 Premier League referees and players. *J. Sports. Sci.* 2011;29:527-532.
- 452 16. Duffield, R, Coutts, AJ, Quinn, J. Core temperature responses and match running
453 performance during intermittent-sprint exercise competition in warm conditions. *J. Strength.*
454 *Cond. Res.* 2009;23:1238-1244.
- 455 17. Buchheit, M, Haydar, B, Ahmaidi, S. Repeated sprints with directional changes: do
456 angles matter? *J. Sports. Sci.* 2012; in press.
- 457

458 **FIGURE CAPTIONS**

459

460 **Figure 1.** Number of repeated-sprint and repeated high-intensity activity (i.e. sprinting and/or
461 striding) effort bouts when analyzed as sets of 2, 3, or 4+ efforts in 15 minute periods
462 throughout the game for national and international players. (A) repeated-sprint activity (RSA)
463 for national players; (B) repeated high-intensity activity (i.e. striding and/or sprinting, RHIA)
464 for national players; (C) repeated-sprint activity for international players; (D) repeated high-
465 intensity activity (i.e. striding and/or sprinting) for international players. Data are presented
466 as the mean number of RSA and RHIA bouts performed in each 15 minute period of the
467 match.

468

469 **Figure 2.** Comparison of first and second half recovery durations per bout for (A) repeated-
470 sprint, and (B) repeated high-intensity activity (i.e. striding and/or sprinting), when analyzed
471 as sets of 2, 3, 4, 5 or 6 efforts, with the recovery criteria being ≤ 20 s between efforts. Data
472 are mean \pm SD. *Moderate effect size (ES = 0.6–1.2).

473

474 **Figure 3.** Comparison of mean recovery duration between repeated-sprint (RSA) and
475 repeated high-intensity activity (i.e. striding and/or sprinting; RHIA) bouts when analyzed as
476 sets of 2, 3, 4, 5 or 6 efforts, with the recovery criteria being ≤ 20 s between efforts. Data are
477 presented as percentage change from sets of 2 bouts (mean \pm SD). *Moderate effect size (0.6–
478 1.2) between repeated-sprint and repeated high-intensity activity (i.e. striding and/or
479 sprinting); †Moderate and ‡Large effect size (1.2–2.0) between bout 2 and subsequent bouts
480 for repeated-sprint; °Moderate and §Large effect size between bout 2 and subsequent bouts
481 for repeated high-intensity activity (i.e. striding and/or sprinting).

482

483 **Figure 4.** Comparison of mean recovery duration for national and international players
484 between (A) repeated-sprint (RSA) and (B) repeated high-intensity activity (i.e. striding
485 and/or sprinting; RHIA) bouts when analyzed as sets of 2, 3, 4, 5 or 6 efforts, with the
486 recovery criteria being ≤ 20 s between efforts. Data are presented as percentage change from
487 sets of 2 bouts (mean \pm SD). *Very large effect size (>2.0) between groups; †Large effect size
488 (1.2–2.0) between groups.

489

490 **Table 1.** Match-play activities used during time-motion analysis of women soccer players.

Match-play activity	Definition
Standing	No locomotor activity
Walking	Movement involves at least one foot being in continual contact with the ground
Jogging	Movement involves a flight phase and minimal arm swing
Striding	Movement is similar to jogging but involves a longer stride and more pronounced arm swing
Sprinting	Maximal effort with a greater extension of the lower leg during forward swing and higher heel lift relative to striding


491

492 **Table 2.** Number of repeated-sprint and repeated high-intensity activity (i.e. sprinting and/or
 493 striding) efforts per bout when analyzed as sets of 2, 3, 4, 5 or 6 efforts, with the recovery criteria
 494 being ≤ 20 s between efforts for national and international players.

Actions per bout	2	3	4	5	6
<i>Sprint</i>					
<i>National</i>					
No. per player/per game	5.3 ± 5.0	2.6 ± 2.8	1.2 ± 1.8	0.4 ± 0.8	0.1 ± 0.3
Sprint Duration (s)	2.21 ± 0.57	2.03 ± 0.49	2.30 ± 0.83	1.96 ± 0.52	1.83 ± 0.24 ^b
Recovery Duration (s)	9.73 ± 4.93	13.64 ± 4.29	14.60 ± 4.18 ^a	14.56 ± 2.22	15.92 ± 2.47
<i>International</i>					
No. per player/per game	4.9 ± 5.4	2.5 ± 3.4	1.1 ± 2.3	0.6 ± 1.6	0.4 ± 1.1
Sprint Duration (s)	2.35 ± 0.56	2.32 ± 0.66	2.12 ± 0.48	1.94 ± 0.56	2.32 ± 0.15
Recovery Duration (s)	10.22 ± 4.42	11.93 ± 4.67	11.97 ± 4.09	15.36 ± 5.99	17.22 ± 2.50
<i>All Matches</i>					
No. per player/per game	5.1 ± 5.1	2.5 ± 3.0 [*]	1.1 ± 2.0 [*]	0.5 ± 1.2 ^{†‡}	0.2 ± 0.7 ^{†‡}
Sprint Duration (s)	2.27 ± 0.56	2.16 ± 0.60	2.24 ± 0.77	1.96 ± 0.51	2.08 ± 0.19
Recovery Duration (s)	9.94 ± 4.73	12.95 ± 4.34 [*]	13.28 ± 3.99 [*]	15.11 ± 3.74 [†]	16.57 ± 2.49 ^{†‡§}
<i>Sprint/Stride</i>					
<i>National</i>					
No. per player/per game	34.4 ± 19.2	24.0 ± 16.9	19.5 ± 17.0	14.9 ± 14.1	12.0 ± 13.1
Sprint/Stride Duration (s)	3.05 ± 0.76	3.02 ± 0.84	3.05 ± 0.86	2.86 ± 0.75	2.96 ± 0.77
Recovery Duration (s)	9.55 ± 2.58	11.34 ± 2.99	12.78 ± 3.04	13.96 ± 2.76	14.76 ± 2.49
<i>International</i>					
No. per player/per game	27.3 ± 17.9	20.0 ± 16.1	14.8 ± 15.9	12.1 ± 15.0	9.9 ± 15.1
Sprint/Stride Duration (s)	2.98 ± 0.66	2.94 ± 0.72	2.98 ± 0.76	2.87 ± 0.65	2.93 ± 0.67
Recovery Duration (s)	9.42 ± 2.68	12.55 ± 3.08	13.72 ± 3.38	14.66 ± 3.16	14.70 ± 3.18
<i>All Matches</i>					
No. per player/per game	31.2 ± 18.7	22.2 ± 16.4 [*]	17.4 ± 16.4 [*]	13.7 ± 14.4 [*]	11.1 ± 13.8 ^{†‡}
Sprint/Stride Duration (s)	2.98 ± 0.66	2.94 ± 0.72	2.98 ± 0.76	2.87 ± 0.65	2.93 ± 0.67
Recovery Duration (s)	9.49 ± 2.60	11.86 ± 3.07 [*]	13.21 ± 3.17 [†]	14.23 ± 2.90 ^{†‡}	14.74 ± 2.66 ^{†‡}

495 Data are mean ± SD. ^{*}Moderate effect size (0.6–1.2) between bouts of 2 and subsequent bouts;
 496 [†]Large effect size (1.2–2.0) between bouts of 2 and subsequent bouts; [‡]Moderate effect size (0.6–
 497 1.2) between bouts of 3 and subsequent bouts; [§]Moderate effect size (0.6–1.2) between bouts of 4
 498 and subsequent bouts. ^aModerate effect size (0.6–1.2) between national and international players;
 499 ^bVery large effect size (>2.0) between national and international players.

500
 501



502
503
504
505

Figure 1.

506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535




Figure 2.

536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553





Figure 3.

554



555

556

557 **Figure 4.**