

Design, Synthesis and Biological Activity of 1,2,3-Triazolobenzodiazepine BET Bromodomain Inhibitors

Phillip P. Sharp, Jean-Marc Garnier, Tamas Hatfaludi, Zhen Xu, David Segal, Kate E. Jarman, Helene Jousset Sabroux, Alexandre Garnham, John T. Feutrill, Anthony Cuzzupe, Peter D. Hall, Scott Taylor, Carl Walkley, Dean Tyler, Mark Dawson, Peter E Czabotar, Andrew F. Wilks, Stefan Glaser, David C. S. Huang, and Christopher J. Burns

ACS Med. Chem. Lett., **Just Accepted Manuscript** • DOI: 10.1021/acsmedchemlett.7b00389 • Publication Date (Web): 14 Nov 2017

Downloaded from <http://pubs.acs.org> on November 22, 2017

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Design, Synthesis and Biological Activity of 1,2,3-Triazolobenzodiazepine BET Bromodomain Inhibitors

Phillip P. Sharp,^{a,b,†,‡} Jean-Marc Garnier,^{a,b,‡} Tamas Hatfaludi,^{a,b,‡} Zhen Xu,^{a,b} David Segal,^{a,b} Kate E. Jarman,^{a,b} H el ene Jousset,^{a,b} Alexandra Garnham,^{a,b} John T. Feutrill,^c Anthony Cuzzupe,^c Peter Hall,^{a,b} Scott Taylor,^d Carl R. Walkley,^d Dean Tyler,^e Mark A. Dawson,^{e,f,g} Peter Czabotar,^{a,b} Andrew F. Wilks,^c Stefan Glaser,^{a,b} David C. S. Huang,^{a,b} Christopher J. Burns^{*,a,b,h}

a) Walter and Eliza Hall Institute of Medical Research, 1G Royal Parade, Parkville, VIC 3052, Australia

b) Department of Medical Biology, The University of Melbourne, VIC 3010, Australia

c) Synthesis MedChem, 399 Royal Parade, Parkville, VIC 3052, Australia

d) St. Vincent's Institute of Medical Research and Department of Medicine, St Vincent's Hospital, University of Melbourne, Fitzroy, VIC 3052, Australia

e) Cancer Research Division, Peter MacCallum Cancer Centre, East Melbourne, VIC 3002, Australia

f) Department of Haematology, Peter MacCallum Cancer Centre, East Melbourne, Victoria 3002, Australia

g) Sir Peter MacCallum Department of Oncology, The University of Melbourne, East Melbourne, Victoria 3002, Australia

h) School of Chemistry, The Bio21 Institute, The University of Melbourne, VIC 3010, Australia

KEYWORDS. *bromodomain, JQ1, triazole, leukemia, benzodiazepine*

ABSTRACT: A number of diazepines are known to inhibit bromo- and extra-terminal domain (BET) proteins. Their BET inhibitory activity derives from the fusion of an acetyl-lysine mimetic heterocycle onto the diazepine framework. Herein we describe a straightforward, modular synthesis of novel 1,2,3-triazolobenzodiazepines and show that the 1,2,3-triazole acts as an effective acetyl-lysine mimetic heterocycle. Structure-based optimization of this series of compounds led to the development of potent BET bromodomain inhibitors with excellent activity against AF9-MLL-driven leukemic cells, concomitant with a reduction in *c-MYC* expression. These novel benzodiazepines therefore represent a promising class of therapeutic BET inhibitors.

The bromo- and extra-terminal domain (BET) family of proteins are epigenetic reader proteins involved in transcription regulation and chromatin remodelling.^{1,2} The family consists of BRD2, BRD3 and BRD4, which are ubiquitously expressed, and BRDT, which is expressed only in the testes. Each BET protein contains two bromodomain structural motifs, here designated as D1 and D2, which have been shown to bind acetylated lysines on histones H3 and H4, and existing data indicates that the biology of BET proteins is largely regulated through the first bromodomain.^{3,4} The acetylated histone lysine residues bind into a hydrophobic pocket of the BET proteins making specific hydrogen bonding interactions with a conserved asparagine and tyrosine residue, the latter through a water molecule.⁵ In the last six years a number of research teams have identified high affinity, small molecule ligands for this hydrophobic pocket that block binding to the cognate acetylated histones.⁶⁻¹⁰ Such small molecule BET inhibitors, along with genetic deletion studies, have demonstrated that inhibition of the BET – histone interaction can result in profound disruption of transcriptional programs resulting in anticancer and anti-inflammatory activity, amongst others. For example, treatment of cancer


cells dependent on the oncogene *c-MYC* with BET inhibitors, either in vitro or in vivo, can result in significant anti-proliferative and cytotoxic effects. As a consequence of these and other data, at least ten BET inhibitors have now entered clinical trials for the treatment of a range of hematological cancers (incl. leukemia, lymphoma and myeloma), certain solid tumors (including glioblastoma multiforme and NUT-midline carcinoma) and atherosclerosis.^{2,11}

The first potent BET inhibitors described in the literature were the benzodiazepine IBET-762 (**1**)^{12,13} and the related thienodiazepine (*S*)-JQ-1 (**2**) (Chart 1).¹⁴ Co-crystal structures of these compounds with BRD4(D1) revealed that the 1,2,4-triazole within these inhibitors acts as an acetyl-lysine mimic by interacting with Asn140 and a conserved water molecule in the KAc recognition pocket. Constellation Pharmaceuticals have reported isoxazole azepines (e.g. **3**)¹⁵ and benzotriazololo[4,3-*d*][1,4]diazepine **4**¹⁶ that also inhibit BET bromodomains, while Knapp and coworkers reported that benzotriazepine **5** binds to BRD4.¹⁷ Interestingly, based on structural similarities to IBET-762, a range of benzodiazepine drugs were retrospectively tested for inhibition of BRD4(D1).

Of these, alprazolam (**6**) was found to be a weak inhibitor ($K_d = 2.5 \mu\text{M}$) and X-ray crystallography confirmed the 1,2,4-triazole acts as an acetyl lysine mimetic. Other closely related analogues of alprazolam that were devoid of the 1,2,4-triazole motif were found to be inactive,¹⁷ underscoring the importance of the acetyl lysine mimicking heterocycle for BET activity.


We have been interested in understanding the impact of the KAc mimicking pharmacophore on bromodomain inhibition. Based on previous studies¹⁸ we designed and tested compounds based on a 5*H*-benzo[*f*][1,2,3]triazolo[1,5-*d*][1,4]diazepine nucleus. The 1,2,3-triazole motif was predicted to have appropriate hydrogen-bond accepting vector at N2 to interact with the conserved asparagine [Asn140 in BRD4(D1)] and molecular modeling indicated that a phenyl substituent at N7 of the diazepine ring would project into the WPF shelf, closely mapping over chlorophenyl (C6) of 3*H*-benzo-1,4-diazepines JQ-1 and IBET-762. We found that this is an excellent BET bromodomain-binding framework and, through subsequent structure-based optimization, developed a series of 1,2,3-triazolobenzodiazepines that display potent BET bromodomain inhibition and excellent cellular activity.

Chart 1. Example azepine BET inhibitors



Initially, we prepared 1,2,3-triazole-containing benzodiazepine **7** via the route shown in Scheme 1. Starting from 1,2-diiodobenzene (**8**), Buchwald coupling with aniline afforded diarylaniline **9**. The propyne was introduced at the remaining iodinated position using Sonogashira cross-coupling and the product (**10**) was acetylated with chloroacetyl chloride providing chloroacetamide **11**. Based on a report by Majumdar *et al.* and others,^{19,21} we performed a one-pot S_N2 /Hüigsen 1,3-dipolar cycloaddition reaction cascade to directly prepare the 1,2,3-triazolobenzodiazepine framework by heating **11** to 150°C in the presence of sodium azide. While this strategy provided triazole **12** as expected, the un-optimized yield in this case was low (13%). Nevertheless, with the diazepinone in hand, selective deletion of the carbonyl oxygen with borane provided the desired 1,2,3-triazolobenzodiazepine **7**. Using an optimized version of this general strategy, followed by additional late-stage cross-coupling reactions and other

functional group interconversions, we synthesized a range of analogs shown in Table 1 (see electronic supporting information for full synthetic details).



Scheme 1. Synthesis of 1,2,3-triazolobenzodiazepines: Synthesis of **7**.

Initially we assessed the BET inhibitory activity of **7** using AlphaScreenTM, a well-validated competition binding assay. Gratifyingly, **7** showed good activity in this assay against all bromodomains tested with the strongest activity against BRD4(D2) ($IC_{50} = 22 \text{ nM}$), but significantly weaker activity against BRD2(D1) ($IC_{50} = 264 \text{ nM}$) (Figure 1 and Table 1). Control compound, JQ-1, gave BRD4(D2) $IC_{50} = 12 \text{ nM}$ and BRD2(D1) $IC_{50} = 26 \text{ nM}$ (Figure 1). We also tested the diazepinone (**12**) but this was essentially inactive (IC_{50} greater than $10 \mu\text{M}$, data not shown).




Figure 1. Dose response plots for AlphaScreenTM assays. Results of at least 2 independent experiments reported as mean ± SD.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The good ligand efficiency [0.51 kcal.mol⁻¹ per non-hydrogen atom for BRD4(D2)] of **7** prompted us to expand this series and explore structure-activity relationships with the view to improve the overall BET activity. First we briefly examined modifications to the D-ring. Benzyl substitution at N7 (**13**) led to a reduction in activity by greater than 50-fold against all bromodomains tested, while addition of either *meta* or *para* substituents on the D-ring were tolerated in terms of overall activity with the 4-chlorophenyl-substituted analog **15** provided a modest improvement over the activity of parent compound **7** [about 2-fold improvement against BRD2(D1)], consistent with previous studies on BET inhibitors, and we therefore retained this modification in subsequent compounds. Next we investigated modifications to the A-ring. Crystal structures of both IBET-762 (pdb code 2YEK) or alprazolam (pdb code 3U5J) bound to BET bromodomains indicated that substitution at C9 (C8 in IBET-762 numbering) projected into the ZA channel and we extrapolated that our series of benzodiazepines would also accommodate functionalization at this position. A set of compounds was therefore designed to incorporate a substituent from the C9 vector. From the activity of the set of C9 analogs it was clear that sp³ substitution provided no improvement in activity (compounds **16**, **17** and **27**), whereas heteroaryl substituents significantly improved activity, with pyrazole **18** and aminopyridine **20** giving IC₅₀ values as low as 7 and 6 nM respectively against BRD4(D2). Interestingly, tetrazole **22** showed the greatest activity across each bromodomain with IC₅₀ values ranging from 7 nM against BRD2(D2) to 2 nM against BRD4(D2). At this point we established the binding mode of the aminopyridyl-substituted compound (**20**) in BRD2(D2) by single crystal X-ray analysis (Figure 2). The structure confirmed that the 1,2,3-triazole motif acts as an acetyl-lysine mimetic, accepting a hydrogen bond from both Asn429 and the conserved water molecule coordinated to Tyr386, as expected. Additionally, the chlorophenyl ring lies within the WPF shelf and the aminopyridine ring extends through the ZA channel and towards solvent exposed space. The positioning of the aminopyridine towards solvent indicated that modifications to this position should be well tolerated. Based on these structural insights we reasoned that other sp²-hybridized moieties, such as carbonyls, might also improve potency and allow for greater diversity of functionalization. Initially we compared carboxylic acids, esters and simple amides. In general, these derivatives showed good activity. Of the initial carbonyl series, primary amide **25** was significantly more potent (~3-fold) against all BET bromodomains tested compared with ester **23** and prompted us to explore the amide derivatives further. The benzyl amide series (compounds **29** to **33**) were also exceptionally potent, particularly chiral benzyl amides **29**, **30** and **31**. Further, in the amide series the *S*-configured amides **29** and **30** displayed significant domain selectivity [BRD2(D2) vs BRD2(D1)] relative to the *R*-configured analog **31**. In light of the potent activity of the tetrazole, **22**, we explored bioisosteric acylsulfonamides at the C9 position. These compounds, although somewhat less active than the tetrazole, also displayed low nanomolar activity against BRD4(D2) despite the significantly increased steric

bulk, further attesting to the accommodation of sp²-modifications at C9. Lastly, we investigated substitutions to the benzodiazepine ring. Methyl substitution at C5 (Compounds **36**, **37**) improved activity against BRD4(D1) and the *R*-stereogenic configuration was preferred consistent with similar modification of previously reported benzodiazepine BET inhibitors.¹⁶

From this chemical series we selected compounds **18** and **20** for further study because they displayed potent activity and were predicted to have good cell permeability (for example the tetrazole **22**, while potent, is likely an efflux transporter substrate).²² First, we assessed the bromodomain selectivity of **20** across a panel of 32 recombinant human bromodomains in a bromoMAX™ assay (see electronic supporting information for full data). In these assays, **20** showed excellent selectivity for the BET family bromodomains. We also assessed the binding kinetics of **20** against individual bromodomains using surface plasmon resonance (SPR). The dissociation-rate measured for compound **20** against BRD2 was approximately 3-fold slower for domain 2 than domain 1 ($k_d = 0.039 \text{ s}^{-1}$ vs $k_d = 0.13 \text{ s}^{-1}$, respectively), whereas association rates were similar ($k_a = 2.90 \times 10^5 \text{ M}^{-1}\text{s}^{-1}$ vs $k_a = 2.62 \times 10^5 \text{ M}^{-1}\text{s}^{-1}$, respectively) (see electronic supporting information for full data).




Figure 2. A X-ray crystal structure of **20** (magenta) in complex with BRD2(D2) PDB code = 5U6V overlaid with co-crystal structure of JQ-1 (grey) bound to BRD2(D2) PDB = 3ONI.

BET inhibition has been shown to have a remarkable effect on certain primary cancer cells and cell lines, which is thought to be a consequence of downregulation of oncogenes *c-MYC*, *BCL-2* and others by displacement of BRD4 from hyperacetylated histone tails near their respective promoters and enhancers.^{23,24} To test the anti-leukemic effect of this chemical series we profiled **18** and **20** against a cancer cell panel (Table 2). Compound **20** potently inhibited proliferation of known BRD4-dependant leukemic line MLL-AF9, and showed significant inhibition of leukemic macrophage cell line MV4-11, (EC₅₀ = 140 nM), whereas B cell lymphoma lines DOHH2, SU-DHL-4 and Raji, and myeloma cell line RPMI-8226 were less sensitive (EC₅₀ > 500 nM). Consistent with BET-targeted anti-proliferative effects, BCR-Abl-driven cell line K562 was completely resistant to **20** (>1000 nM).

Table 1. Inhibition of BET bromodomains as determined by AlphaScreen™ assays. Numbers represent mean IC₅₀ in nanomolar (nM) of at least two independent experiments. **A.** D-ring modifications. **B** C9 sp³ functionalized derivatives. **C** C9 sp²-functionalized derivatives. **D** C9 carbonyl derivatives. **E** miscellaneous C9 derivatives. **F** C9 benzamide derivatives. **G** Acylsulphonamides. **H** C5 Alkyl derivatives.


IC ₅₀ (nM)	BRD2			BRD3			BRD4		
	domain 1	domain 2		domain 1	domain 2		domain 1	domain 2	
7	264	19	28	>999	>999		129	21	29
	39		12	>999	>999		44		17
13									
14									
15							129	25	35
							43		32
16							273	52	45
							147		38
17							388	36	79
							40		30
18							35	10	14
							25		7
19							64	14	23
							38		12
20							33	9	13
							21		6
21							53	10	16
							28		7
22							6		3
							7		2
23							116	18	29
							52		18
24							185	16	28
							50		17
25							34		6
							17		2
26							79	14	21
							18		13
27							204		43
							370		23
28							56		11
							24		13
29							95	23	34
							4		17
30							51	12	10
							3		6
31							68	13	14
							20		13
32							136	21	35
							37		15
33							151	39	44
							44		14
34							54	21	21
							25		10
35							44		13
							25		7
36							53	13	15
							34		10
37							38	11	14
							25		7
38							340	44	52
							45		36

Osteosarcoma (OS) cells are also known to be sensitive to BET inhibition, but through a MYC-independent mechanism.²⁵ We were therefore interested to characterize the effect of **18** and **20** on primary mouse OS cells (fibroblastic 494H and osteoblastic 148I). Both compounds **18** and **20** inhibited proliferation of both primary OS cell types, serving to highlight the broad utility of 1,2,3-triazolobenzodiazepines derivatives in cancer studies. The cellular potency profile of **18** and **20** also closely correlated with positive control JQ-1.

Table 2. Treatment of various cancer cell lines with **18**, **20** or JQ-1 for 72 h. Cell viability measure by Cell-TiterGlo™

Cell line	EC ₅₀ (nM)		
	20	18	JQ-1
MLL-AF9	77	111	64
HL-60	152	139	51
MV4-11	140	200	39
DOHH2	561	285	279
RPMI-8226	631	495	258
SU-DHL-4	977	570	660
CCRF-CEM	1365	2221	2037
Raji	560	780	330
HEL92.1.7	400	720	180
K562	>10000	>10000	>10000
494H	166	194	122
148I	1285	973	284

We used RNA-seq to compare the effect of **20** and JQ-1 on global gene expression in THP-1 leukemia cells. These experiments revealed that the effect on gene expression between these two compounds is highly correlated for both up and down regulated genes (Figure 3A). In particular, BRD4-dependent oncogene *c-MYC* demonstrated robust downregulation on treatment with **20** (Figure 3B) whilst *HEXIM1* levels were upregulated.²⁶ Similar effects on *c-MYC* and *BCL-2* downregulation were observed for JQ-1, **18**, **20** and non-diazepine BET inhibitor IBET-151 in MV4-11 cells (data not shown, see electronic supporting information for full data).



benzo[*f*][1,2,3]triazolo[1,5-*d*][1,4]diazepines. Structure-

Figure 3. A. GENAS plot comparing effects of **20** and JQ-1 on THP-1 cells. B. Identity of 10 highest up and down regulated genes by **20** and JQ-1.

activity relationship studies and crystal structure-guided

optimization led to the development of analogs with single-digit nanomolar activity. Compound **20**, showed good BET bromodomain selectivity and potent activity against a number of BRD4-dependant leukaemia cell lines and caused down-regulation of oncogene *MYC*. Taken together these data show that the benzodiazepine scaffold is an useful framework on which to base BET bromodomain inhibitors.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

Experimental methods, synthesis, and characterization (PDF)

AUTHOR INFORMATION

Corresponding Author

* (C.B.) email: burns@wehi.edu.au

Present Addresses

†. Department of Cellular and Molecular Pharmacology, University of California, San Francisco, United States.

Author Contributions

‡These authors contributed equally.

Funding Sources

Our work is supported by the NHMRC (fellowship to PEC) and the Victorian State Government Operational Infrastructure Support, and the Australian Government NHMRC IRIISS.

Notes

The authors declare no competing financial interests.

ACKNOWLEDGMENT

We thank beamline staff at the Australian Synchrotron, where diffraction data were collected. Crystallization experiments were performed at the C3 Collaborative Crystallization Centre.

ABBREVIATIONS

BET, bromo- and extra C-terminal domain containing protein.

REFERENCES

- Belkina, A. C.; Denis, G. V. BET Domain Co-Regulators in Obesity, Inflammation and Cancer. *Nat. Rev. Cancer* 2012, 12 (7), 465–477.
- Filippakopoulos, P.; Knapp, S. Targeting Bromodomains: Epigenetic Readers of Lysine Acetylation. *Nat. Rev. Drug Discov.* 2014, 13 (5), 337–356.
- Picaud, S.; Wells, C.; Felletar, I.; Brotherton, D.; Martin, S.; Savitsky, P.; Diez-Dacal, B.; Philpott, M.; Bountra, C.; Lingard, H.; Fedorov, O.; Müller, S.; Brennan, P. E.; Knapp, S.; Filippakopoulos, P. RVX-208, an Inhibitor of BET Transcriptional Regulators with Selectivity for the Second Bromodomain. *Proc. Natl. Acad. Sci. U. S. A.* 2013, 110 (49), 19754–19759.
- Baud, M. G. J.; Lin-Shiao, E.; Cardote, T.; Tallant, C.; Pschibul, A.; Chan, K.-H.; Zengerle, M.; Garcia, J. R.; Kwan, T. T.-L.; Ferguson, F. M.; Ciulli, A. A Bump-and-Hole Approach to

Engineer Controlled Selectivity of BET Bromodomain Chemical Probes. *Science* 2014, 12 (7), 465–477.

(5) Filippakopoulos, P.; Picaud, S.; Mangos, M.; Keates, T.; Lambert, J. P.; Barseyte-Lovejoy, D.; Felletar, I.; Volkmer, R.; Müller, S.; Pawson, T.; Gingras, A. C.; Arrowsmith, C. H.; Knapp, S. Histone Recognition and Large-Scale Structural Analysis of the Human Bromodomain Family. *Cell* 2012, 149 (1), 214–231.

(6) Garnier, J.-M.; Sharp, P. P.; Burns, C. J. BET Bromodomain Inhibitors: A Patent Review. *Expert Opin. Ther. Pat.* 2013, 24 (2), 1–15.

(7) Müller, S.; Knapp, S. Discovery of BET Bromodomain Inhibitors and Their Role in Target Validation. *Medchemcomm* 2014, 5 (3), 288.

(8) Brand, M.; Measures, A. M.; Wilson, B. G.; Cortopassi, W. A.; Alexander, R.; Höss, M.; Hewings, D. S.; Rooney, T. P. C.; Paton, R. S.; Conway, S. J. Small Molecule Inhibitors of Bromodomain–Acetyl-Lysine Interactions. *ACS Chem. Biol.* 2015, 10 (1), 22–39.

(9) Zhang, G.; Smith, S. G.; Zhou, M. M. Discovery of Chemical Inhibitors of Human Bromodomains. *Chem. Rev.* 2015, 115 (21), 11625–11668.

(10) Liu, Z.; Wang, P.; Chen, H.; Wold, E. A.; Tian, B.; Brasier, A. R.; Zhou, J. Drug Discovery Targeting Bromodomain-Containing Protein 4. *J. Med. Chem.* 2017, 60 (11), 4533–4558.

(11) Gelato, K. A.; Shaikhibrahim, Z.; Ocker, M.; Haendler, B. Targeting Epigenetic Regulators for Cancer Therapy: Modulation of Bromodomain Proteins, Methyltransferases, Demethylases, and microRNAs. *Expert Opin. Ther. Targets* 2016, 20 (7), 783–799.

(12) Nicodeme, E.; Jeffrey, K. L.; Schaefer, U.; Beinke, S.; Dewell, S.; Chung, C.; Chandwani, R.; Marazzi, I.; Wilson, P.; Coste, H.; White, J.; Kirilovsky, J.; Rice, C. M.; Lora, J. M.; Prinjha, R. K.; Lee, K.; Tarakhovskiy, A. Suppression of Inflammation by a Synthetic Histone Mimic. *Nature* 2010, 468 (7327), 1119–1123.

(13) Mirguet, O.; Gosmini, R.; Toum, J.; Clément, C. A.; Barnathan, M.; Brusq, J. M.; Mordaunt, J. E.; Grimes, R. M.; Crowe, M.; Pineau, O.; Ajakane, M.; Daugan, A.; Jeffrey, P.; Cutler, L.; Haynes, A. C.; Smithers, N. N.; Chung, C. W.; Bamborough, P.; Uings, I. J.; Lewis, A.; Witherington, J.; Parr, N.; Prinjha, R. K.; Nicodème, E. Discovery of Epigenetic Regulator I-bet762: Lead Optimization to Afford a Clinical Candidate Inhibitor of the BET Bromodomains. *J. Med. Chem.* 2013, 56 (19), 7501–7515.

(14) Filippakopoulos, P.; Qi, J.; Picaud, S.; Shen, Y.; Smith, W. B.; Fedorov, O.; Morse, E. M.; Keates, T.; Hickman, T. T.; Felletar, I.; Philpott, M.; Munro, S.; McKeown, M. R.; Wang, Y.; Christie, A. L.; West, N.; Cameron, M. J.; Schwartz, B.; Heightman, T. D.; La Thangue, N.; French, C. A.; Wiest, O.; Kung, A. L.; Knapp, S.; Bradner, J. E. Selective Inhibition of BET Bromodomains. *Nature* 2010, 468 (7327), 1067–1073.

(15) Gehling, V. S.; Hewitt, M. C.; Vaswani, G. R.; et al.; Hewitt, M. C.; Vaswani, R. G.; Leblanc, Y.; Coité, A.; Nasveschuk, C. G.; Taylor, A. M.; Harmange, J. C.; Audia, J. E.; Pardo, E.; Joshi, S.; Sandy, P.; Mertz, J. A.; Sims, R. J.; Bergeron, L.; Bryant, B. M.; Bellon, S.; Poy, F.; Jayaram, H.; Sankaranarayanan, R.; Yellapantula, S.; Bangalore Srinivasamurthy, N.; Birudukota, S.; Albrecht, B. K. Discovery, Design, and Optimization of Isoxazole Azepine BET Inhibitors. *ACS Med. Chem. Lett.* 2013, 4 (9), 835–840.

(16) Taylor, A. M.; Vaswani, R. G.; Gehling, V. S.; Hewitt, M. C.; Leblanc, Y.; Audia, J. E.; Bellon, S.; Cummings, R. T.; Coité, A.; Harmange, J. C.; Jayaram, H.; Joshi, S.; Lora, J. M.; Mertz, J. A.; Neiss, A.; Pardo, E.; Nasveschuk, C. G.; Poy, F.; Sandy, P.; Setser, J. W.; Sims, R. J.; Tang, Y.; Albrecht, B. K.

Discovery of Benzotriazolo[4,3-d][1,4]diazepines as Orally Active Inhibitors of BET Bromodomains. *ACS Med. Chem. Lett.* 2016, 7 (2), 145–150.

(17) Filippakopoulos, P.; Picaud, S.; Fedorov, O.; Keller, M.; Wrobel, M.; Morgenstern, O.; Bracher, F.; Knapp, S. Benzodiazepines and Benzotriazepines as Protein Interaction Inhibitors Targeting Bromodomains of the BET Family. *Bioorganic Med. Chem.* 2012, 20 (6), 1878–1886.

(18) Sharp, P. P.; Garnier, J.-M.; Huang, D. C. S.; Burns, C. J. Evaluation of Functional Groups as Acetyl-Lysine Mimetics for BET Bromodomain Inhibition. *Medchemcomm* 2014, 5, 1834–1842.

(19) Majumdar, K. C.; Ray, K.; Ganai, S.; Ghosh, T. Catalyst-Free 1,3-Dipolar Cycloaddition: An Efficient Route for the Formation of the 1,2,3-Triazole-Fused Diazepinone Framework. *Synthesis (Stuttg.)* 2010, No. 5, 858–862.

(20) Hooyberghs, G.; De Coster, H.; Vachhani, D. D.; Ermolat'Ev, D. S.; Van Der Eycken, E. V. Synthesis of [1,2,3]-triazolo[1,5-a][1,4]benzodiazepines via an Unprecedented One-Pot Cu-Catalyzed Azidation-Cyclization Reaction. *Tetrahedron* 2013, 69 (21), 4331–4337.

(21) Sudhapriya, N.; Nandakumar, A.; Arun, Y.; Perumal, P. T.; Balachandran, C.; Emi, N. An Expedient Route to Highly Diversified [1,2,3]triazolo[1,5-a][1,4]benzodiazepines and Their Evaluation for Antimicrobial, Antiproliferative and in Silico Studies. *RSC Adv.* 2015, 5 (81), 66260–66270.

(22) Young, A. M.; Audus, K. L.; Proudfoot, J.; Yazdani, M. Tetrazole Compounds: The Effect of Structure and pH on Caco-2 Cell Permeability. *J. Pharm. Sci.* 2006, 95 (4), 717–725.

(23) Chapuy, B.; McKeown, M. R.; Lin, C. Y.; Monti, S.; Roemer, M. G.; Qi, J.; Rahl, P. B.; Sun, H. H.; Yeda, K. T.; Doench, J. G.; Reichert, E.; Kung, a L.; Rodig, S. J.; Young, R. a; Shipp, M. a; Bradner, J. E. Discovery and Characterization of Super-Enhancer-Associated Dependencies in Diffuse Large B Cell Lymphoma. *Cancer Cell* 2013, 24 (6), 777–790.

(24) Lovén, J.; Hoke, H. A.; Lin, C. Y.; Lau, A.; Orlando, D. A.; Vakoc, C. R.; Bradner, J. E.; Lee, T. I.; Young, R. A. Selective Inhibition of Tumor Oncogenes by Disruption of Super-Enhancers. *Cell* 2013, 153 (2), 320–334.

(25) Baker, E. K.; Taylor, S.; Gupte, A.; Sharp, P. P.; Walia, M.; Walsh, N. C.; Zannettino, A. C.; Chalk, A. M.; Burns, C. J.; Walkley, C. R. BET Inhibitors Induce Apoptosis through a MYC Independent Mechanism and Synergise with CDK Inhibitors to Kill Osteosarcoma Cells. *Sci. Rep.* 2015, 5 (May), 10120.

(26) Devaraj, S. G. T.; Fiskus, W.; Shah, B.; Qi, J.; Sun, B.; Iyer, S. P.; Sharma, S.; Bradner, J. E.; Bhalla, K. N. HEXIM1 Induction Is Mechanistically Involved in Mediating Anti-AML Activity of BET Protein Bromodomain Antagonist. *Leukemia* 2016, 30 (2), 504–508.

Insert Table of Contents artwork here

