

**Research Bank**

Journal article

**Paranoia and interpersonal functioning across the continuum from healthy to pathological – Network analysis**

**Hajdúk, Michal, Klein, Hans S., Harvey, Philip D., Penn, David L. and Pinkham, Amy E.**

This is the peer reviewed version of the following article:

Hajdúk, M., Klein, H. S., Harvey, P. D., Penn, D. L. and Pinkham, A. E. (2019). Paranoia and interpersonal functioning across the continuum from healthy to pathological – Network analysis. *British Journal of Clinical Psychology*, 58(1), pp. 19-34, which has been published in final form at <https://doi.org/10.1111/bjc.12199>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions. This article may not be enhanced, enriched or otherwise transformed into a derivative work, without express permission from Wiley or by statutory rights under applicable legislation. Copyright notices must not be removed, obscured or modified. The article must be linked to Wiley's version of record on Wiley Online Library and any embedding, framing or otherwise making available the article or pages thereof by third parties from platforms, services and websites other than Wiley Online Library must be prohibited



Published in final edited form as:

*Br J Clin Psychol.* 2019 March ; 58(1): 19–34. doi:10.1111/bjc.12199.

## Paranoia and interpersonal functioning across the continuum from healthy to pathological – Network Analysis

Michal Hajdúk, Ph.D.<sup>1,2</sup>, Hans S. Klein, MS.<sup>3</sup>, Philip D. Harvey, Ph.D.<sup>4,5</sup>, David L. Penn, Ph.D.<sup>6,7</sup>, and Amy E. Pinkham, Ph.D.<sup>3,8</sup>

<sup>1</sup>Department of Psychology, Faculty of Arts, Comenius University, Bratislava, Slovak Republic

<sup>2</sup>Department of Psychiatry, Faculty of Medicine, Comenius University, Bratislava, Slovak Republic

<sup>3</sup>School of Behavioral and Brain Sciences, The University of Texas at Dallas, Richardson, TX, USA

<sup>4</sup>Department of Psychiatry and Behavioral Sciences, University of Miami Miller School of Medicine, Miami, FL, USA

<sup>5</sup>Research Service, Miami VA Healthcare System, Miami, US

<sup>6</sup>Department of Psychology, University of North Carolina, Chapel Hill, NC, USA

<sup>7</sup>School of Psychology, Australian Catholic University, Melbourne, VIC, Australia

<sup>8</sup>Department of Psychiatry, University of Texas Southwestern Medical School, Dallas, TX, USA

### Abstract

**Objective:** Previous research has suggested that paranoia is associated with impaired social functioning in patients with schizophrenia and healthy individuals with high levels of paranoid ideation. The present study analyzed relationships between paranoia and interpersonal functioning across the paranoia continuum using network analysis.

**Method:** Levels of paranoid ideation and interpersonal functioning were measured in a non-clinical sample (N=853) and in patients with schizophrenia spectrum disorders (N=226). Network analyses were used to examine the nature of paranoia's relation to interpersonal functioning in both populations.

**Results:** Although the most central characteristic of paranoia in both samples was the feeling of being talked about behind one's back, across samples, individual characteristics were differentially related to various aspects of interpersonal functioning. Among clinical individuals, difficulties in

---

Correspondence should be addressed to Michal Hajdúk, Department of Psychology, Comenius University, Gondova 2, 814 99 Bratislava, Slovak Republic, Phone: +421 2 59 339 418, michal.hajduk@uniba.sk.

Declaration of Interest:

P.D.H serves as a consultant/advisory board member for Boehringer Ingelheim, Lundbeck, Otsuka Digital Health, Roche, Sanofi, Sunovion, and Takeda. M.H. reported receiving a fee from Lundbeck Slovakia as a speaker at an education grant conference. Other authors report no conflicts of interest.

Conflict of Interest:

We also added new information regarding the declaration of potential conflicts because the situation of one author (M.H.) changed after we submitted the manuscript. The following text has now been added.

„M.H. reported receiving a fee from Lundbeck as a speaker at an education grant conference. “

interpersonal functioning were related to perceived previous experiences of being treated poorly by others, whereas among the non-clinical sample, interpersonal functioning was related to negative beliefs about others.

**Conclusions:** The current results support previous findings linking paranoid ideation to interpersonal functioning in both clinical and non-clinical samples. Patterns of these relationships differed slightly across groups. Results in general support a continuum model of paranoia.

### Keywords

Paranoia; Interpersonal Functioning; Network analysis

---

## Introduction

Paranoia is a common symptom of mental disorders that is prominent in schizophrenia but also present in depression and bipolar disorder (Bentall et al., 2009). In psychiatric rating scales, paranoid ideation is usually measured via single items like suspiciousness, or hostility (Kay, Fiszbein, & Opler, 1987; Overall & Gorham, 1962). Recently however paranoia has been conceptualized not as a single construct, but rather as a hierarchy spanning from common emotional concerns like fear of rejection, social anxiety, and feelings of vulnerability to the most extreme forms of paranoia such as the persecutory delusions that can be seen in patients in acute stages of psychosis (Freeman et al., 2005). Data from the British National Survey of Psychiatric Morbidity (Bebbington et al., 2013) supports this view, demonstrating that paranoia has a multidimensional structure consisting of the following four factors: interpersonal sensitivities, mistrust, ideas of reference, and ideas of persecution.

Mirroring the hierarchical nature of paranoia, empirical studies have demonstrated that paranoia exists on a continuum spanning both psychopathology as well as the general population (Bebbington et al., 2013; Freeman et al., 2005; Freeman, 2016;). Indeed, conservative estimates indicate that 10–15% of individuals in the general population regularly experience paranoid thoughts (Freeman, 2007). While the degree of preoccupation, conviction, and associated distress tends to differentiate between non-clinical and clinical levels of persecutory ideation (Peters, Joseph, & Garety, 1999; Peters, Joseph, Day, & Garety, 2004), a growing body of literature suggests that paranoia is associated with impaired social functioning and reduced psychological well-being across this continuum (Martin & Penn, 2001; Olfson et al., 2002; Rössler et al., 2007). In the general population, overall higher levels of paranoid ideation are associated with greater social exclusion, lower social functioning, restricted social support, and less engagement in hobbies or activities of interest (Freeman et al., 2005, 2011). In clinical samples, distress associated with persecutory delusions mediates the association between these delusions and aggressive behavior towards others (van Dongen, Buck, & van Marle, 2012), and the presence of clinically significant levels of paranoid ideation appears to selectively exacerbate the social impairments that are seen in individuals with schizophrenia (Pinkham, Harvey, & Penn, 2016).

Despite these strong links between paranoia and social impairment, it is currently unclear which aspects of paranoia have the greatest impact on social functioning. Recent advances in psychopathological research have suggested that psychiatric disorders can be modeled as complex dynamic networks of interrelated symptoms (Kendler, Zachar, & Craver, 2011). This can be done via network analysis (NA), which differs from traditional psychometric approaches based on the assumption of latent variables that influence symptom expression (e.g. common - cause model). NA enables investigators to model symptom to symptom interactions (Borsboom, 2008; Borsboom & Cramer, 2013), and in extreme cases, associations between symptoms constitute disorders. The NA approach has been used to model networks of symptoms in psychotic disorder (Isvoranu et al., 2016), depression (Hajdúk, Heretik, & Pe e ák, 2017; van Borkulo et al., 2015), post-traumatic stress disorder (McNally et al., 2015), and substance abuse (Rhemtulla et al., 2016). Some studies have also utilized NA for modeling psychiatric comorbidity (Cramer, Waldorp, van der Maas, & Borsboom, 2010; McNally, Mair, Mugno, & Riemann, 2017), but it is also possible that NA can be used to examine links between multidimensional symptoms (i.e., paranoia) and outcomes (i.e., social functioning) (Jones, Heeren, & McNally, 2017).

The overall goal of the present study is therefore to analyze the relationships between elements of paranoid ideation and interpersonal functioning using a NA approach. The aims of the present study are following: (i) Identify the associations between specific dimensions of paranoia and various aspects of interpersonal functioning, (ii) Qualitatively assess the identified relationships in clinical and non-clinical samples to determine if the relationships are consistent across the paranoia continuum.

## Methods

To examine the continuum of paranoia, data were pooled across two larger studies conducted in our lab. The first study involved only non-clinical individuals and was administered via an online survey as part of a screening assessment for undergraduate research participants. Clinical participants were drawn from the Social Cognition Psychometric Evaluation project (SCOPE) (Pinkham et al., 2014; Pinkham, Penn, Green, & Harvey, 2016), in which data were collected via in person assessments. The same primary measure of paranoia was used across studies however, the measures of social functioning differed to allow informant reports of functioning for the clinical participants. Both studies were approved by the Institutional Ethical Review Board, and all participants provided informed consent.

## Participants

**Non-clinical sample**—The non-clinical sample was comprised of undergraduate students who completed the online screening assessment in exchange for course credit. After excluding participants who demonstrated careless responding (i.e., participants who did not complete the survey or gave the same item response across all questionnaires,  $n=87$ ), the final sample consisted of 853 students. Demographic characteristics are presented in Table 1.

**Clinical sample**—Clinical participants were individuals with diagnoses of schizophrenia or schizoaffective disorder. One hundred seventy nine patients were from the initial

psychometric study phase of SCOPE (Pinkham, Penn, et al., 2016), and 51 patients were from the modification phase (Buck, Pinkham, Harvey, & Penn, 2016; Cornacchio, Pinkham, Penn, & Harvey, 2017). Two patients from the initial psychometric study and two from the modification phase were dropped due to missing data on Specific Level of Functioning Scale. The final sample therefore consisted of 226 patients. Diagnoses were confirmed with the Mini International Neuropsychiatric Interview (Sheehan et al., 1998) and the Structured Clinical Interview for DSM Disorders - Psychosis Module (First, Spitzer, Gibbon, & Williams, 2012). Symptom severity was assessed with the Positive and Negative Syndrome Scale (PANSS: (Kay et al., 1987). Information regarding full inclusion and exclusion criteria for the clinical population can be found in Pinkham et al. (2016). Basic demographic characteristics are presented in Table 1.

## Measures

**Paranoia Scale**—Paranoid ideation was evaluated with the Paranoia Scale (PS), which consists of 20 items that measure subclinical levels of paranoid ideation (Fenigstein & Vanable, 1992). Participants were asked to read each statement and indicate how much each statement applies to him/her. Each item is scored on a 1–5 Likert scale with total scores ranging from 20 to 100 and higher scores indicating greater paranoid ideation. A recent factor analysis demonstrates that the scale consists of three factors: Self-depreciation, Mistrust Thoughts, and Persecutory Ideas (Carvalho et al., 2017). The PS is often used in non-clinical samples for measuring subclinical levels of paranoid ideation (Combs, 2004; Martin & Penn, 2001) but has also been used validly in clinical samples (Craig, 2004; Pinkham et al., 2012).

**Social Functioning Scale**—In the non-clinical sample, social functioning was assessed with the Social Functioning Scale (SFS: (Birchwood, Smith, Cochrane, Wetton, & Copestake, 1990)). The SFS is a self-report measure of seven domains of social adjustment. Three subscales (i.e., Communication rated 0 – 9, Social Engagement/Withdrawal rated 0 – 15, and Prosocial Activities rated 0 – 66) were utilized in the current analyses as these scales assess interpersonal social functioning. Other subscales measuring independence and vocational activities were not assessed here. Higher scores on the three subscales indicate better social functioning (e.g., more competent behavior or higher frequency behavior). The original scale has been demonstrated to be a reliable, valid, and sensitive instrument for patients with schizophrenia and has been successfully used in non – clinical samples (Iffland et al., 2015).

**Specific Level of Functioning**—Among patients, objective real-world social functioning was assessed using informant reports on the Specific Level of Functioning Scale (SLOF) (Schneider & Struening, 1983). The SLOF consists of 30 items assessing four domains: Interpersonal Relationships, Social Acceptability, Involvement in Activities, and Work Skills. An additional item assesses how well the informant knows the participant. Responses were gathered from informants who were high contact clinicians, family members, or close friends identified by the participants. Items are scored on 5-point Likert scale. Higher scores represent better functioning. To focus specifically on interpersonal aspects of social functioning, the present analyses utilized only items from the Interpersonal

Relationships (7 items) and Social Acceptability (6 items) subscales. Supporting this decision, previous work comparing paranoid to non-paranoid patients found that these subgroups of patients differed only on these two subscales but not Work Skills or Involvement in Activities (Pinkham, Harvey, et al., 2016).

### Network analysis – construction and analysis

We constructed independent networks for clinical and non-clinical samples, using the 20 items of the Paranoia Scale and measures of social functioning, SLOF and SFS, as “nodes” in the network. Specifically, the items from the SLOF interpersonal relationships and social acceptability subscales were used for our clinical sample, and the total scores for the SFS Communication, Social Engagement/Withdrawal, and Prosocial Activities subscales were used for our non-clinical sample. For the SFS, total subscale scores were used instead of individual item scores to improve interpretability and reliability of NA. Specifically, item scores on this scale are diverse, and we do not propose direct relationships to paranoid ideation for individual items. For example, Prosocial Activities consists of 22 individual prosocial behaviors (e.g. playing sports with others, visiting relatives, eating out in restaurants etc.), and Social Withdrawal contains an item asking at what time a person wakes in the morning. Using these items as individual nodes may be problematic due to an increased number of parameters being estimated, as well as decreasing interpretability of relationships between individual items and paranoid ideation. Thus, subscale scores were deemed more suitable for this exploratory analysis to ensure easily interpretable networks. Table 2 details the nodes and their abbreviations displayed in each of the networks’ figures.

Network structures were estimated with R-packages *qgraph* (Epskamp et al., 2012) and *bootnet* (Epskamp & Fried, 2017). Regularized Gaussian graphical models (GGM) were used to fit data. In the network, associations between nodes are displayed as edges, representing the partial correlations between these two nodes after controlling for all other relationships in the network, and the thickness of the edges indicate strength of these relationships. *Glasso* procedure was used for estimation of the network. Positive relationships are presented as solid lines, dashed lines represent negative relationships. To control for spurious connections within the network, least absolute shrinkage and selection (*lasso*) operator was used, identifying a stable, parsimonious and easily interpretable network. Lasso is a form of network regularization based on Extended Bayesian Information Criterion (Chen & Chen, 2008) and shrinks small regression coefficients, to zero. The remaining number of edges are used to explain the covariation structure in the data. In order to assist with visual interpretation of nodular significance, nodes with stronger and more numerous edges were moved to the center of the network using the Fruchterman - Reingold algorithm (Epskamp, Borsboom, & Fried, 2018).

Several indices are estimated in order to evaluate the importance of each node in the network. In the present analysis we used the following centrality estimates: i) Strength – indicates the magnitude of direct relationships with other nodes in network, ii) Betweenness – indicates the probability that a node lies along the shortest path between other nodes in the network, iii) Closeness – indicates the magnitude of indirect relationships with other nodes in the network.

Accuracy of networks were evaluated using an estimation of the accuracy of edge weights, utilizing a bootstrap with 95% confidence interval on edge weights. Furthermore, we investigated the stability of the centrality measures by sub-setting participants, i.e. re-estimating the networks after dropping participants. Estimates are considered stable if there is a strong association (correlation above 0.70) between original estimates and estimates from a subset of the sample. In both samples, estimation of accuracy and stability analyses were performed on 1000 bootstrap samples (Epskamp et al., 2018). Stability, expressed as the CS coefficient, should be at least 0.25 for centrality measures to be considered stable, and ideal values should be above 0.50 (Epskamp et al., 2018).

## Results

### Non – clinical sample

The stability coefficient for our primary measure of interest, Strength (CS (0.70) = 0.517) was over the recommended minimal threshold, as was the stability coefficient for Closeness (CS (0.70) = 0.284). The stability coefficient for Betweenness (CS (0.70) = 0.128) were below threshold, and therefore should be interpreted with care. Figures for network accuracy analysis for edge weight are in the online supplement of the manuscript. Inspection of the network reveals that the most central nodes in terms of strength in the non-clinical sample were within the PS, specifically items PS14 (*I am sure I have been talked about behind my back*), PS09 (*I often wonder what hidden reason another person may have for doing something nice for you*), and PS03 (*I believe that I have often been punished without cause*). The node with highest betweenness and closeness was item PS10 (*It is safer to trust no one*) indicating connections with a larger number of nodes in the network. Within network, the strongest association was found within the PS scale between nodes PS14 (*I am sure I have been talked about behind my back*) and P17 (*People have said insulting and unkind things about me*). The second strongest relationship was found between items PS04 (*Some people have tried to steal my ideas and takes credit for them*) and PS20 (*I have often found people jealous of my good ideas just because they had not thought of them first*). Additionally, the total scores for the three SFS subscales, Communication (COM), Social Engagement/Withdrawal (ENG), and Prosocial Activities (PRO), were strongly associated with each other. Across measures, the SFS Communication subscale was negatively associated with items PS06 (*No one really cares much what happens to you*) and PS10 (*It is safer to trust no one*). A higher level of Social Engagement was associated with lower scores on items related to self-depreciation, PS06 (*No one really cares much what happens to you*) and PS05 (*My parents and family find more fault with me than they should*). Social Engagement was also negatively associated with persecutory ideas as measured by item PS19 (*I am bothered by people outside, in cars, in stores, etc., watching me*). Estimated network and centrality estimates are presented in Figures 1 and 2.

### Clinical sample

As with the non-clinical sample, the stability coefficient for our primary measure of interest, strength, was over the recommended minimal threshold (CS (0.70) = 0.518). The stability coefficients for Closeness (CS (0.70) = 0.204) and Betweenness (CS (0.70) = 0) were below threshold. As our intent with this analysis is to describe the relationships between individual

items, and in line with recommendations from Epskamp et al. (2018), the presented analysis only interprets the strength centrality measure. Figures from network accuracy analysis for edge weight are available in the online supplement of the manuscript. The network of paranoia and interpersonal functioning for the clinical sample is presented in Figure 3. As expected, relationships between nodes within the measures are stronger than those between measures, and the two subscales of the SLOF formed separate sub-clusters in the network. The most central nodes in terms of strength indices in the whole network were: PS14, PS19, PS18, PS11. Centrality values for strength, closeness, and betweenness are presented in Figure 4. Within the nodes from PS, the strongest associations were found between nodes PS14 (*I am sure I have been talked about behind my back*) and PS17 (People have said insulting and unkind things about me). Additionally, two other pairs of strongly associated nodes were noted within the network, i.e. items PS1 (*Someone has it in for me*) and PS2 (*I sometimes feel as if I am being followed*), as well as items PS11 (I have often felt that strangers were looking at me critically) and PS19 (I am bothered by people outside, in cars, in stores etc. watching me). This relationship is consistent with previous factor analysis stating that these items load on a single, persecutory ideation factor (Carvalho et al., 2017). In terms of the whole network, the strongest associations were found within SLOF domains (IR01 and IR02; SA04 and SA05). Across measures, item SA01 (*Regularly argues with others*) from the SLOF social acceptability subscale was uniquely and negatively associated with PS17 (People have said insulting and unkind things about me). Additionally, the strongest negative association between the PS and the SLOF Interpersonal relationships subscale was between item IR07 (*Asks for help when needed*) and PS03 (I believe that I have often been punished without cause).

## Discussion

The current article provides the first network analysis of the relationship between paranoia and interpersonal functioning across the paranoia continuum. Overall, results supported associations between interpersonal functioning and paranoid symptoms both within the normative population and among individuals with severe mental illness. However, unique and distinct associations were observed between patients with schizophrenia and a healthy, non-clinical sample in both the relationships between different aspects of paranoia and how these aspects of paranoia are related to interpersonal functioning. A major benefit of network analysis is that high centrality nodes can be considered proxy indicators of clinical importance, and thus those nodes, or in this case specific aspects of paranoia, represent potential key targets of therapeutic intervention. These results therefore indicate that remediation of the particular elements of paranoia identified here could improve interpersonal functioning. Further, as aspects of paranoia appeared to be differentially related to functioning across samples, the current results suggest that targets of intervention should vary slightly pending the clinical status of the individual.

Consistent with previous factor analysis, the most central symptoms in our networks were aspects of persecutory ideations, such as feelings that others had talked about oneself behind one's back, being disappointed by others, and distressing feelings of being watched by others. Furthermore, feeling that others talked about oneself behind their back was most central in both clinical and non-clinical networks. This finding is consistent with previous



research noting that worries about being criticized or rejected in social situations is a central symptom of paranoia in general populations (Bell & O'Driscoll, 2018). In our clinical sample, item PS17 (People have said insulting and unkind things about me) had the highest value of betweenness, indicating that this sensitivity for criticism could play an important role in triggering and maintaining paranoid ideation.

Differences between samples were also evident in the relationships between paranoia and social functioning. In the clinical sample, lower quality of interpersonal relationships was associated with more feelings of being punished without cause (ideas of persecution). Additionally, beliefs that others are saying unkind or insulting things (persecutory ideation) were associated with less socially acceptable behavior (arguing with others). In our non-clinical sample, social engagement and quality of interpersonal communication were associated with interpersonal sensitivity and mistrust. Some of these differences may be partially due to methodological differences between samples. As self-report measures of social functioning for patients with schizophrenia may not accurately estimate true levels of functioning (Sabbag et al., 2011; Ermel et al., 2017), our analyses utilized ratings of interpersonal functioning as assessed by high contact informants who know the patient well. Network analyses may differ when using only self-reported measures of functioning, as self-report measures may inflate the degree of relationships simply due to the fact that the same person is making ratings (i.e., shared method variance). Therefore, using an impartial assessment of interpersonal functioning may lead to fewer connections, but should result in a less biased network.

Not surprisingly, interpersonal functioning in our non-clinical sample seems to be predominantly associated with lower paranoia severity in contrast to our clinical sample. However, this demonstrates that levels of subclinical paranoid ideation may have a meaningful impact on daily functioning. Socially stressful situations are associated with increased negative emotions, predicting the onset of paranoia in psychosis (Myin-Germeys, van Os, Schwartz, Stone, & Delespaul, 2001; Veling, Pot-Kolder, Counotte, van Os, & van der Gaag, 2016), and in subclinical individuals, social conflict may lead to increased paranoid ideation, or vice versa. In our clinical sample, lower frequency and quality of social contact may be explained as a safety behavior that patients utilize to reduce the perceived threat of others (Freeman et al., 2007). Future studies are needed to examine the causal mechanisms of social conflict and isolation in seeking social support and how these, along with other confounding symptoms of schizophrenia such as negative symptoms, play a role in paranoid ideation and social functioning.

### **Limitations and future directions**

The male-female ratio, age, and proportion of each race differed across samples, which partially limits direct comparisons of networks. The non-clinical sample was also more diverse than our clinical population. As being a member of a minority group and migration are well known risk factors of psychotic disorder (Morgan, Charalambides, Hutchinson, & Murray, 2010), future studies should match clinical and non-clinical groups on demographic factors in order to perform direct comparisons.

Another limitation of this study is our measure of paranoia. Despite the fact that the Paranoia Scale is widely used in research, the 20 items do not sufficiently cover all relevant aspects of paranoid ideation. As such, the networks presented do not account for other dimensions of paranoid ideation, namely degree of conviction and level of associated distress. These dimensions may play a vital role in social functioning and analyzing these aspects may provide a more comprehensive understanding of how paranoid ideation is related to interpersonal functioning across the continuum.

Furthermore, our study utilized different measures of interpersonal functioning in the clinical and non-clinical samples, which further limits direct comparison of estimated networks. In our clinical sample, we used SLOF items as assessed by independent raters, which we feel is more appropriate to reveal fine grained relationships in a clinical population. However, this measure may not be sensitive enough to measure variability in a non-clinical sample. Thus, for our non-clinical sample, we utilized the SFS, which allowed for more variability, but limited the interpretability of individual items within the network analysis. Although SLOF items assess generalized behaviors such as whether a participant engages in activity without prompting and with self-sufficiency in daily tasks, the individual items on the SFS measure distinct activities, such as how often a participant goes to the movies or attends evening classes. This limits the ability to interpret a network analysis on an item by item basis with the SFS, and subscale totals must be utilized. This reduced our ability to assess associations between paranoia and specific behaviors in our non-clinical sample.

Although the aim of the current study was to observe the network structure of paranoid ideation and social functioning in each of our two groups, and not to fully examine the structure of paranoia itself across groups, future research should compare clinical samples with matched healthy controls to better examine whether the structures of these paranoia networks are the same in across the continuum.

Longitudinal studies are also needed to evaluate how positive or negative social events cause changes in levels of paranoid ideation. Experience sampling methods may reveal whether changes in network are only in terms of overall connectivity or in the structure of paranoia. Future studies should also address the impact of other important variables outside social functioning such as positive and negative emotions and negative symptoms.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgements:

This work was supported by the National Institute of Mental Health at the National Institutes of Health (R01 MH093432 to P.H.D., D.L.P., and A.E.P.). M.H was supported by Fulbright Visiting Scholar Grant.

## References


- Bebbington PE, McBride O, Steel C, Kuipers E, Radovanovi M, Brugha T, ... Freeman D (2013). The structure of paranoia in the general population. *The British Journal of Psychiatry*, 202(6), 419–427. 10.1192/bjp.bp.112.119032 [PubMed: 23661767]
- Bell V, & O'Driscoll C (2018). The network structure of paranoia in the general population. *Social Psychiatry and Psychiatric Epidemiology*, 1–8. 10.1007/s00127-018-1487-0
- Bentall RP, Rowse G, Shryane N, Kinderman P, Howard R, Blackwood N, ... Corcoran R (2009). The cognitive and affective structure of paranoid delusions: a transdiagnostic investigation of patients with schizophrenia spectrum disorders and depression. *Archives of General Psychiatry*, 66(3), 236–247. Retrieved from <http://archneur.jamanetwork.com/article.aspx?articleid=483012> [PubMed: 19255373]
- Birchwood M, Smith J, Cochrane R, Wetton S, & Copestake S (1990). The Social Functioning Scale. The development and validation of a new scale of social adjustment for use in family intervention programmes with schizophrenic patients. *The British Journal of Psychiatry: The Journal of Mental Science*, 157, 853–859. [PubMed: 2289094]
- Borsboom D (2008). Psychometric perspectives on diagnostic systems. *Journal of Clinical Psychology*, 64(9), 1089–1108. 10.1002/jclp.20503 [PubMed: 18683856]
- Borsboom D, & Cramer AOJ (2013). Network Analysis: An Integrative Approach to the Structure of Psychopathology. *Annual Review of Clinical Psychology*, 9(1), 91–121. 10.1146/annurev-clinpsy-050212-185608
- Buck BE, Pinkham AE, Harvey PD, & Penn DL (2016). Revisiting the validity of measures of social cognitive bias in schizophrenia: Additional results from the Social Cognition Psychometric Evaluation (SCOPE) study. *British Journal of Clinical Psychology*, 55(4), 441–454. 10.1111/bjc.12113 [PubMed: 27168196]
- Carvalho CB, Sousa M, da Motta C, Pinto-Gouveia J, Caldeira SN, Peixoto EB, ... Fenigstein A (2017). Paranoia in the general population: A revised version of the General Paranoia Scale for adults. *Clinical Psychologist*, 21(2), 125–134. 10.1111/cp.12065
- Chen J, & Chen Z (2008). Extended Bayesian information criteria for model selection with large model spaces. *Biometrika*, 95(3), 759–771. 10.1093/biomet/asn034
- Combs D (2004). The role of subclinical paranoia on social perception and behavior. *Schizophrenia Research*, 69(1), 93–104. 10.1016/S0920-9964(03)00051-3 [PubMed: 15145475]
- Cornacchio D, Pinkham AE, Penn DL, & Harvey PD (2017). Self-assessment of social cognitive ability in individuals with schizophrenia: Appraising task difficulty and allocation of effort. *Schizophrenia Research*, 179, 85–90. 10.1016/j.schres.2016.09.033 [PubMed: 27693281]
- Craig J (2004). Persecutory beliefs, attributions and theory of mind: comparison of patients with paranoid delusions, Asperger's syndrome and healthy controls. *Schizophrenia Research*, 69(1), 29–33. 10.1016/S0920-9964(03)00154-3 [PubMed: 15145468]
- Cramer AOJ, Waldorp LJ, van der Maas HLJ, & Borsboom D (2010). Comorbidity: A network perspective. *Behavioral and Brain Sciences*, 33(2–3), 137–150. 10.1017/S0140525X09991567 [PubMed: 20584369]
- Epskamp S, Borsboom D, & Fried EI (2018). Estimating psychological networks and their accuracy: A tutorial paper. *Behavior Research Methods*, 50(1), 1–18. 10.3758/s13428-017-0862-1 [PubMed: 29340969]
- Epskamp S, Cramer AO, Waldorp LJ, Schmittmann VD, Borsboom D, & others. (2012). qgraph: Network visualizations of relationships in psychometric data. *Journal of Statistical Software*, 48(4), 1–18. Retrieved from <http://www.jstatsoft.org/htaccess.php?volume=48&type=i&issue=04&paper=true>
- Epskamp S, & Fried EI (2017). bootnet: Bootstrap Methods for Various Network Estimation Routines Retrieved from <https://cran.r-project.org/web/packages/bootnet/index.html>
- Ernel J, Carter CS, Gold JM, MacDonald AW, Daniel Ragland J, Silverstein SM, ... Barch DM (2017). Self versus informant reports on the specific levels of functioning scale: Relationships to depression and cognition in schizophrenia and schizoaffective disorder. *Schizophrenia Research: Cognition*, 9, 1–7. 10.1016/j.scog.2017.04.001 [PubMed: 28740827]

- Fenigstein A, & Venable PA (1992). Paranoia and self-consciousness. *Journal of Personality and Social Psychology*, 62(1), 129–138. Retrieved from <http://psycnet.apa.org/journals/psp/62/1/129/> [PubMed: 1538311]
- First MB, Spitzer RL, Gibbon M, & Williams JB (2012). *Structured Clinical Interview for DSM-IV® Axis I Disorders (SCID-I)*. American Psychiatric Pub
- Freeman D (2007). Suspicious minds: The psychology of persecutory delusions. *Clinical Psychology Review*, 27(4), 425–457. 10.1016/j.cpr.2006.10.004 [PubMed: 17258852]
- Freeman D (2016). Persecutory delusions: a cognitive perspective on understanding and treatment. *The Lancet Psychiatry*, 3(7), 685–692. 10.1016/S2215-0366(16)00066-3 [PubMed: 27371990]
- Freeman D, Garety PA, Bebbington PE, Smith B, Rollinson R, Fowler D, ... Dunn G (2005). Psychological investigation of the structure of paranoia in a non-clinical population. *The British Journal of Psychiatry*, 186(5), 427–435. Retrieved from <http://bjp.rcpsych.org/content/186/5/427.short> [PubMed: 15863749]
- Freeman D, Garety PA, Kuipers E, Fowler D, Bebbington PE, & Dunn G (2007). Acting on persecutory delusions: The importance of safety seeking. *Behaviour Research and Therapy*, 45(1), 89–99. 10.1016/j.brat.2006.01.014 [PubMed: 16530161]
- Freeman D, McManus S, Brugha T, Meltzer H, Jenkins R, & Bebbington P (2011). Concomitants of paranoia in the general population. *Psychological Medicine*, 41(5), 923–936. 10.1017/S0033291710001546 [PubMed: 20735884]
- Hajdúk M, Heretik A, & Pešek J (2017). Nové možnosti výskumu psychopatológie: sieťová analýza symptómov depresie. *Psychiatrie*, 21(1), 12–17.
- Iffland JR, Lockhofen D, Gruppe H, Gallhofer B, Sammer G, & Hanewald B (2015). Validation of the German Version of the Social Functioning Scale (SFS) for Schizophrenia. *PLoS ONE*, 10(4). 10.1371/journal.pone.0121807
- Isvoranu A-M, Borkulo C. D. van, Boyette L-L, Wigman JTW, Vinkers CH, Borsboom D, & Investigators G (2016). A Network Approach to Psychosis: Pathways Between Childhood Trauma and Psychotic Symptoms. *Schizophrenia Bulletin*, 43(1), 187–196. 10.1093/schbul/sbw055 [PubMed: 27165690]
- Jones PJ, Heeren A, & McNally RJ (2017). Commentary: A network theory of mental disorders. *Frontiers in Psychology*, 8 10.3389/fpsyg.2017.01305
- Kay SR, Fiszbein A, & Opler LA (1987). The positive and negative syndrome scale (PANSS) for schizophrenia. *Schizophrenia Bulletin*, 13(2), 261–276. [PubMed: 3616518]
- Kendler KS, Zachar P, & Craver C (2011). What kinds of things are psychiatric disorders? *Psychological Medicine*, 41(6), 1143–1150. 10.1017/S0033291710001844 [PubMed: 20860872]
- Martin JA, & Penn DL (2001). Social cognition and subclinical paranoid ideation. *The British Journal of Clinical Psychology / the British Psychological Society*, 40(Pt 3), 261–265.
- McNally RJ, Mair P, Mugno BL, & Riemann BC (2017). Co-morbid obsessive-compulsive disorder and depression: a Bayesian network approach. *Psychological Medicine*, 47(7), 1204–1214. 10.1017/S0033291716003287 [PubMed: 28052778]
- McNally RJ, Robinaugh DJ, Wu GW, Wang L, Deserno MK, & Borsboom D (2015). Mental disorders as causal systems a network approach to posttraumatic stress disorder. *Clinical Psychological Science*, 3(6), 836–849. Retrieved from <http://cpx.sagepub.com/content/3/6/836.short>
- Morgan C, Charalambides M, Hutchinson G, & Murray RM (2010). Migration, Ethnicity, and Psychosis: Toward a Sociodevelopmental Model. *Schizophrenia Bulletin*, 36(4), 655–664. 10.1093/schbul/sbq051 [PubMed: 20513653]
- Myin-Germeys I, Os J. van, Schwartz JE, Stone AA, & Delespaul PA (2001). Emotional Reactivity to Daily Life Stress in Psychosis. *Archives of General Psychiatry*, 58(12), 1137–1144. 10.1001/archpsyc.58.12.1137 [PubMed: 11735842]
- Olfson M, Lewis-Fernández R, Weissman MM, Feder A, Gameraoff MJ, Pilowsky D, & Fuentes M (2002). Psychotic symptoms in an urban general medicine practice. *The American Journal of Psychiatry*, 159(8), 1412–1419. 10.1176/appi.ajp.159.8.1412 [PubMed: 12153836]
- Overall JE, & Gorham DR (1962). The brief psychiatric rating scale. *Psychological Reports*, 10(3), 799–812. Retrieved from <http://www.amsciepub.com/doi/abs/10.2466/pr0.1962.10.3.799>


- Peters E, Joseph S, Day S, & Garety P (2004). Measuring delusional ideation: the 21-item Peters et al. Delusions Inventory (PDI). *Schizophrenia Bulletin*, 30(4), 1005–1022. [PubMed: 15954204]
- Peters ER, Joseph SA, & Garety PA (1999). Measurement of delusional ideation in the normal population: introducing the PDI (Peters et al. Delusions Inventory). *Schizophrenia Bulletin*, 25(3), 553–576. Retrieved from <http://schizophreniabulletin.oxfordjournals.org/content/25/3/553.short> [PubMed: 10478789]
- Pinkham AE, Harvey PD, & Penn DL (2016). Paranoid individuals with schizophrenia show greater social cognitive bias and worse social functioning than non-paranoid individuals with schizophrenia. *Schizophrenia Research: Cognition*, 3, 33–38. 10.1016/j.scog.2015.11.002 [PubMed: 27990352]
- Pinkham AE, Penn DL, Green MF, Buck B, Healey K, & Harvey PD (2014). The social cognition psychometric evaluation study: results of the expert survey and RAND panel. *Schizophrenia Bulletin*, 40(4), 813–823. 10.1093/schbul/sbt081 [PubMed: 23728248]
- Pinkham AE, Penn DL, Green MF, & Harvey PD (2016). Social Cognition Psychometric Evaluation: Results of the Initial Psychometric Study. *Schizophrenia Bulletin*, 42(2), 494–504. 10.1093/schbul/sbv056 [PubMed: 25943125]
- Pinkham AE, Sasson NJ, Beaton D, Abdi H, Kohler CG, & Penn DL (2012). Qualitatively distinct factors contribute to elevated rates of paranoia in autism and schizophrenia. *Journal of Abnormal Psychology*, 121(3), 767–777. 10.1037/a0028510 [PubMed: 22686868]
- Rhemtulla M, Fried EI, Aggen SH, Tuerlinckx F, Kendler KS, & Borsboom D (2016). Network analysis of substance abuse and dependence symptoms. *Drug and Alcohol Dependence*, 161, 230–237. 10.1016/j.drugalcdep.2016.02.005 [PubMed: 26898186]
- Rössler W, Riecher-Rössler A, Angst J, Murray R, Gamma A, Eich D, ... Gross VA (2007). Psychotic experiences in the general population: a twenty-year prospective community study. *Schizophrenia Research*, 92(1–3), 1–14. 10.1016/j.schres.2007.01.002 [PubMed: 17363221]
- Sabbag S, Twamley EM, Vella L, Heaton RK, Patterson TL, & Harvey PD (2011). Assessing Everyday Functioning in Schizophrenia: Not all Informants Seem Equally Informative. *Schizophrenia Research*, 131(1–3), 250–255. 10.1016/j.schres.2011.05.003 [PubMed: 21620682]
- Schneider LC, & Struening EL (1983). SLOF: a behavioral rating scale for assessing the mentally ill. *Social Work Research & Abstracts*, 19(3), 9–21. [PubMed: 10264257]
- Sheehan DV, Lecrubier Y, Harnett K, Amorim P, Janavs J, Weiller E, ... Dunbar GC (1998). The Mini-International Neuropsychiatric Interview (M.I.N.I): The development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *Journal of Clinical Psychiatry*, 59(Suppl 20), 22–33.
- van Borkulo C, Boschloo L, Borsboom D, Penninx BH, Waldorp LJ, & Schoevers RA. (2015). Association of symptom network structure with the course of depression. *JAMA Psychiatry*, 72(12), 1219–1226. 10.1001/jamapsychiatry.2015.2079 [PubMed: 26561400]
- van Dongen JDM, Buck NML, & van Marle HJC (2012). Delusional distress partly explains the relation between persecutory ideations and inpatient aggression on the ward. *Psychiatry Research*, 200(2), 779–783. 10.1016/j.psychres.2012.05.016 [PubMed: 22809853]
- Veling W, Pot-Kolder R, Counotte J, van Os J, & van der Gaag M (2016). Environmental Social Stress, Paranoia and Psychosis Liability: A Virtual Reality Study. *Schizophrenia Bulletin*, 42(6), 1363–1371. 10.1093/schbul/sbw031 [PubMed: 27038469]

### Practitioner Points

- Network analyses were used to identify central aspects of persecutory ideation in both clinical and non-clinical samples. Qualitative assessment of clinical and non-clinical networks revealed similar central symptoms and supported a continuum model of paranoia.
- Central aspects of paranoia, i.e. feeling that others have talked about oneself behind one's back, being disappointed by others, and having distressing feelings of being watched by others, were associated with deficits in interpersonal functioning in both samples.
- Central aspects of paranoia may be beneficial targets for psychosocial interventions aimed at reducing paranoid ideation and improving interpersonal functioning.
- Demographic characteristics for the current study differed between samples which may limit generalization of findings.
- Future research is needed to explore temporal associations and moment-to-moment dynamics between paranoid ideation and problems in interpersonal functioning.




**Figure 1.**  
Estimated network on non-clinical sample.



**Figure 2. Centrality estimates for non-clinical sample network.**  
 Strength indicates the magnitude of direct relationships with other nodes in network;  
 Betweenness indicates the probability that a node lies along the shortest path between other  
 nodes in the network, Closeness indicates the magnitude of indirect relationships with other  
 nodes in the network.







**Figure 4. Centrality estimates for non-clinical sample network.**

Strength indicates the magnitude of direct relationships with other nodes in network;

Betweenness indicates the probability that a node lies along the shortest path between other nodes in the network, Closeness indicates the magnitude of indirect relationships with other nodes in the network.

**Table 1.**

## Demographic characteristics of samples

Characteristic	Non-clinical sample		Clinical sample	
	N	%	N	%
Gender				
Male	198	23.2	144	63.7
Female	652	76.4	82	36.3
Race				
Caucasian	347	40.7	94	41.6
African American	64	7.5	119	52.7
Asian	337	39.5		
Other	105	12.3	13	5.7
Diagnosis	-	-		
Schizophrenia			118	52.2
Schizoaffective disorder			108	47.8
	M	SD	M	SD
Age	21.36	4.36	42.25	11.96
PANSS	-	-		
Positive			16.15	5.59
Negative			13.51	4.93
General			30.92	7.94
Paranoia scale	39.35	12.39	47.46	19.22
SFS			-	-
Social engagement	10.85	2.26		
Communication	8.12	1.10		
Prosocial	27.90	10.86		
SLOF	-	-		
Interpersonal relationships			3.32	0.90
Social Acceptability			4.33	0.58

M – Mean, SD – Standard deviation

**Table 2:**

## Explanation of nodes abbreviations

P01	Someone has it in for me
P02	I sometimes feel as if I'm being followed
P03	I believe that I have often been punished without cause
P04	Some people have tried to steal my ideas and take credit for them
P05	My parents and family find more fault with me than they should
P06	No one really cares much what happens to you
P07	I am sure I get a raw deal from life
P08	Most people will use somewhat unfair means to gain profit or advantage, rather than lose it
P09	I often wonder what hidden reason another person may have for doing something nice for you
P10	It is safer to trust no one
P11	I have often felt that strangers were looking at me critically
P12	Most people make friends because friends are likely to be useful to them
P13	Someone has been trying to influence my mind
P14	I am sure I have been talked about behind my back
P15	Most people inwardly dislike putting themselves out to help other people
P16	I tend to be on my guard with people who are somewhat more friendly than expected
P17	People have said insulting and unkind things
P18	People often disappoint me
P19	I am bothered by people outside, in cars, in stores, etc., watching me
P20	I have often found people jealous of my good ideas just because they had not thought of them first
IR01	Accept contact with others
IR02	Initiates contact with others
IR03	Communicates effectively
IR04	Engages in activities without prompting
IR05	Participates in groups
IR06	Forms and maintains friendships
IR07	Asks for help when needed
SA01	Regularly argues with others
SA02	Has physical fights with others
SA03	Destroys property
SA04	Physically abuses self
SA05	Is fearful, crying, clinging
SA06	Takes property from others without permission
COM	SFS – Communication
PRO	SFS – Prosocial behavior

ENG	SFS – Social Engagement
-----	-------------------------

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript