

Research Bank

Journal article

Lower limb muscle size after anterior cruciate ligament injury : A systematic review and meta-analysis

Dutaillis, Benjamin, Maniar, Nirav, Opar, David A., Hickey, Jack T. and Timmins, Ryan G.

Accepted manuscript. Link to publisher's version <https://doi.org/10.1007/s40279-020-01419-0>

Dutaillis, Benjamin, Maniar, Nirav, Opar, David A., Hickey, Jack T. and Timmins, Ryan G.. (2021) Lower limb muscle size after anterior cruciate ligament injury : A systematic review and meta-analysis. Sports Medicine. 51(6), pp. 1209-1226.
<https://doi.org/10.1007/s40279-020-01419-0>

1 **Title**

2 Lower Limb Muscle Size After Anterior Cruciate Ligament Injury: A Systematic Review and Meta-Analysis.

3 **Short title**

4 Muscle Size in ACL injured limbs.

5 **Authors**

6 Benjamin Dutaillis 1, Nirav Maniar 1, David A Opar 1,2, Jack T Hickey 1, Ryan G Timmins 1,2.

7 **ORCID ID's:**

8 Benjamin Dutaillis: 0000-0002-8747-3796, Nirav Maniar: 0000-0002-6180-6003, David Opar: 0000-0002-8354-
9 6353, Jack Hickey: 0000-0003-2764-698X, Ryan Timmins: 0000-0003-4964-1848.

10 **Affiliations**

11 1. School of Behavioural and Health Sciences, Australian Catholic University, Fitzroy, Victoria, Australia.

12 2. Sports Performance, Recovery, Injury and New Technologies (SPRINT) Research Centre, Australian Catholic
13 University, Fitzroy, Victoria, Australia.

14 **Corresponding Author**

15 Benjamin Dutaillis

16 benji.dutaillis@outlook.com

17 +61 432 155 749

18 Australian Catholic University

19 School of Behavioural and Health Sciences

20 17 Young Street, Fitzroy, Victoria, Australia, 3065

21

Abstract

Background: Anterior cruciate ligament (ACL) injury is known to have a number of deleterious effects on lower limb muscle function. Alterations in muscle size are one such effect that has implications towards reductions in strength and functioning of the lower limbs. However, a comprehensive analysis of alterations in muscle size has yet to be undertaken.

Objective: To systematically review the evidence investigating lower limb muscle size in ACL injured limbs.

Design: Systematic review

Data sources: Database searches of Medline, SPORTDiscus, Embase, Cinahl and Web of Science as well as citation tracking and manual reference list searching.

Eligibility criteria for selecting studies: Individuals with ACL deficient or reconstructed limbs with an assessment of lower limb muscle size and control limb data (contralateral or uninjured control group)

Methods: Risk of bias assessment was completed on included studies. Data was extracted and where possible meta-analyses performed. Best evidence synthesis was also undertaken.

Results: 49 articles were included in this review, with 37 articles included in the meta-analyses. 66 separate meta-analyses were performed using various measures of lower limb muscle size. Across all measures, ACL deficient limbs showed lesser quadriceps femoris muscle size (d range= -0.35 to -0.40), whereas ACL reconstructed limbs showed lesser muscle size in the quadriceps femoris (d range= -0.41 to -0.69), vastus medialis (d= -0.25), vastus lateralis (d= -0.31, hamstrings (d= -0.28), semitendinosus (d range= -1.02 to -1.14) and gracilis (d range= -0.78 to -0.99) when compared to uninjured limbs.

Conclusion: This review highlights the effect ACL injury has on lower limb muscle size. Regardless of whether an individual chooses a conservative or surgical approach, the quadriceps of the injured limb appear to have lesser muscle size compared to an uninjured limb. When undertaking reconstructive surgery with a semitendinosus/gracilis tendon graft, the harvested muscle shows lesser muscle size compared to the uninjured limb.

Author contributions

Concept and design (BD, NM, DO and RT), running of search strategy (BD), inclusion/exclusion criteria assessment (BD and JH), quality assessment (BD and RT), data extraction (BD), data analysis and interpretation (BD, NM, DO, JH and RT), manuscript preparation (BD, NM, DO, JH and RT).

Declarations

Funding

The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests.

55 Benjamin Dutailis, Nirav Maniar, David Opar, Jack Hickey and Ryan Timmins declare that they have no conflicts
56 of interest relevant to the content of this review.

57 **Availability of data and material**

58 Not applicable.

59 **Code availability**

60 Not applicable.

61 **Ethics approval**

62 Not applicable.

63 **Consent to participate**

64 Not applicable.

65 **Consent for publication**

66 Not applicable.

67 **Key points:**

- 68
- Anterior cruciate ligament (ACL) deficient limbs possess lower quadriceps femoris cross sectional area
69 and muscle volume than the contralateral uninjured limb;
 - Limbs with a prior ACL reconstruction have evidence of reduced muscle size in the quadriceps femoris,
70 vastus medialis, vastus lateralis, the hamstrings, semitendinosus and gracilis compared to the
71 contralateral uninjured limb;
 - In the prior ACL reconstructed limbs, the evidence of reduced muscle size in the quadriceps femoris is
72 greatest within the first 30 weeks post-surgery;
 - In the prior ACL reconstructed limbs with semitendinosus or semitendinosus-gracilis grafts, there is a
73 large effect and evidence of reduced muscle size in the semitendinosus up to 348 weeks post-surgery.
74
75
76

77

1 Introduction

Anterior cruciate ligament (ACL) rupture, a debilitating injury typically treated with surgical reconstruction to re-establish structural integrity of the knee, is characterised by a lengthy rehabilitation period [1, 2]. Rates of ACL reconstructions are increasing in the USA, Australia, England and New Zealand [3-7], suggesting an increase worldwide. For example, in Australia the rates of ACL reconstruction have risen in the last 15 years and are now the highest in the world per capita at 77.4 per 100 000 persons [7, 6]. The estimated annual cost of ACL surgery to the Australian economy is over \$72 million (AUD)[7], not including rehabilitation, income replacement, societal costs and long-term disability associated with the injury. Recent work has shown only 55-83% of people returning to pre-injury levels of sport [1, 8] and 23% of patients passing return to play criteria [9]. While it is unknown whether the primary injury, surgery, or a combination of both are responsible, an estimated 50% of ACL reconstructed individuals present with radiographic evidence of knee joint osteoarthritis within 10 years following surgery [10, 11].

The long-term negative impacts of ACL injury and reconstructive surgery on knee function are multifactorial. Prior work has identified persistent deficits in objective markers of lower limb muscle function following rehabilitation, including reduced muscle strength [12, 13], muscle activation [14-17], muscle fibre force production [18] and muscle cross sectional area (CSA) [19-21]. Reductions in quadriceps muscle volume is commonly noted in both ACL deficient [22, 23] and reconstructed [24-26] individuals. Additionally, reduced semitendinosus and gracilis volume [19, 27] is seen following harvesting of the respective tendons for reconstructive surgery. The reductions in lower limb muscle size seen after ACL injury and surgery may be linked to the decreased levels of strength in ACL injured limbs [28, 29, 19].

Reductions in quadriceps strength is one factor that leads to alterations in biomechanics and consequently knee joint loading commonly seen following ACL injury. These factors may subsequently influence the development of knee joint osteoarthritis [30-33] and risk of secondary injury [34, 17]. Recent work has also highlighted the influence of other knee and non-knee spanning muscles in contributing to knee joint loading [35-37]. Alterations in the size and strength of other lower limb muscles therefore may contribute to the development of knee joint osteoarthritis and secondary ACL rupture following ACL injury. Combined, these maladaptation's suggest that adequate and timely recovery of lower limb muscle size is imperative in restoring health, function and performance in ACL injured individuals.

While a number of systematic reviews have investigated reductions in muscle strength [12, 38], only one has investigated changes in muscle size following ACL injury [39], which focused solely on the quadriceps, without a meta-analysis. Therefore, we aimed to review and meta-analyse the evidence base related to changes in the size of all lower limb muscles in ACL injured populations to better understand the effect of injury and surgery on these individuals, and guide clinical prognoses.

2 Method

2.1 Study Design

This review was registered on PROSPERO (ID: CRD42019129262) to reduce the risk of reporting bias and minimise research wastage.

This review is compliant with the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) guidelines [40]. A comprehensive systematic literature search of Medline, SPORTDiscus, Embase, Cinahl and Web of Science was conducted. The search terms (Table 1) were chosen in order to identify original research articles that fit the aims of the review. Where possible relevant MeSH and subject headings were included in the search strategies. The search captured all studies from inception to 17/03/2020 with retrieved references being imported into EndNote X8 (Thomson Reuters, New York City, NY, USA) and all subsequent screening followed the PRISMA guidelines [40].

2.2 Study Selection

A pre-determined set of inclusion and exclusion criteria were used to screen retrieved articles (Table 2). The lead author (BD) screened the titles and abstracts for relevance. Articles deemed appropriate underwent full text review which was conducted by two authors (BD and JH) for inclusion in the review. Any disputes were discussed and resolved through consultation with a third author (RT).

2.3 Study Quality Assessment

Methodological quality assessment of included articles was performed using a modified version of the Downs and Black Checklist [41] by two authors (BD and RT). The original checklist contains 27 items, however, a number of these are only relevant to intervention studies and as this review included mainly retrospective studies, items 4, 8, 13, 14, 15, 19, 20, 22, 23 and 24 were deemed inappropriate to assess study quality and were therefore removed. An additional item (29) was added to assess whether rehabilitation among participants was controlled for and reported in each study [42] (see Electronic Supplementary Material Table S1).

2.4 Data Extraction and Analysis

Data extraction included the population (ACL deficient or reconstructed), sample size, control comparisons (whether the uninjured contralateral limb and/or a healthy control group), time since injury or surgery, graft type used in reconstruction, imaging method used (magnetic resonance imaging, computed tomography or ultrasound), size measurement (e.g muscle CSA, volume or thickness) and site of measurement (if relevant). For all extracted data, group mean and standard deviations (SD) for all reported muscle(s) and muscle groups were collated. Where articles reported standard error (SE), SD was calculated using; $SD = SE \times \sqrt{N}$ due to the statistical analysis applied (SD = standard deviation, SE = standard error, N = sample size).

For articles that reported measures at multiple time points, data from each point was extracted to allow for subgroup and regression analysis. However, articles that reported pre-surgical measures were not included in the ACL deficient analysis [43, 25, 20, 44-46], as clear time points were not given as to when the pre-surgical measure was taken, and thus may have contaminated results. Where data were not available or reported as median rather than mean, corresponding authors were contacted for data in mean and standard deviation.

Where sufficient data were available, meta-analyses was conducted using the ‘metafor’ [47] and ‘meta’ [48] packages in R (R Development Core Team. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing, 2019. Due to the differing methods and techniques used to calculate the obtained measures of muscle size, standardised mean differences (Cohen’s d) and 95% confidence intervals were used to facilitate comparisons of studies. A random-effects model with a restricted maximum

likelihood (REML) method was used to estimate the overall effect within each measure and subgroup, with the p value set at 0.05. The size of the effect was interpreted as either small ($d = 0.20$ to 0.49), moderate ($d = 0.50$ to 0.79) and large ($d \geq 0.8$)[49]. Where a statistically significant overall effect size was seen within a measure, pooled time-point subgroups and meta-regressions were used to estimate the effect of time since surgery within the reconstructed limbs. Where the meta-regression relationship was logarithmic rather than linear, analysis was run with log transformed 'time since surgery' data. Due to the lack of reporting in articles investigating ACL deficient limbs, similar subgroup analyses and meta-regressions were not able to be run in this population.

In some cases, multiple groups from the same article may appear within a single meta-analysis. This was due to the reporting style and data in these articles being different populations (e.g. male and female, experiment and control). The decision was made to preserve the original mean and SD values and included them in the meta-analysis as separate cohorts. Where this occurs, article names in the forest plots are followed by brackets indicating the subgroup from the article.

Where meta-analysis was unable to be run due the inability to obtain data from the corresponding author, a best evidence synthesis was employed [50]. The level of evidence was ranked according to the following criteria;

- Strong: two or more studies of a high quality and generally consistent findings ($\geq 75\%$ of studies showing consistent results).
- Moderate: one high-quality study and/or two or more low quality studies and generally consistent findings ($\geq 75\%$ of studies showing consistent results).
- Limited: one low-quality study.
- Conflicting: inconsistent findings ($< 75\%$ of studies showing consistent results).
- None: no supportive findings in the literature.

3 Results

3.1 Search Results

The initial search yielded 11635 articles (Cinahl = 1187, Embase = 2693, Medline = 3129, SPORTDiscus = 1079 and Web of Science = 3547). After duplicate removal, title/abstract screening and full text review, a total of 49 articles were deemed eligible for inclusion in this review (Figure 1).

3.2 Study Quality Assessment

Electronic Supplementary Material Table S2 shows the results of the quality assessment. Study quality ranged from 7 to 20 out of 20, with 27 articles (55%) deemed high quality.

3.3 Meta-analysis

Of the 49 articles included in this review, 37 [51-57, 43, 25, 58-63, 19, 26, 20, 64-66, 22, 67, 68, 27, 69, 44, 21, 70, 45, 71, 72, 29, 46, 23, 73, 74] were included in the meta-analyses. The data has been grouped in to five main comparisons:

1. Muscle CSA of ACL deficient limbs compared to the contralateral uninjured limb (Table 3);
2. Muscle volume of ACL deficient limbs compared to the contralateral uninjured limb (Table 4);

3. Muscle CSA of ACL reconstructed limbs compared to the contralateral uninjured limb (Table 5);
4. Muscle volume of ACL reconstructed limbs compared to the contralateral uninjured limb (Table 6) and
5. Muscle volume of ACL reconstructed limbs compared to a healthy control group (Electronic Supplementary Material Table S3).

Overall, there were 66 meta-analyses run, with 11 showing statistically significant effects. These 11 indicated reduced muscle size in specific muscles and muscle groups within the ACL deficient or reconstructed limbs when compared to the contralateral uninjured limbs. No statistically significant effects were found when comparing the ACL reconstructed limb to a healthy control group.

3.4 ACL Deficient Populations

A summary of the results of the meta-analyses comparing the muscle CSA and volume of ACL deficient limbs to the contralateral uninjured limbs are found in tables 3 and 4, respectively. Of these results, 12 measures of muscle CSA and 11 measures of muscle volume were analysed, with 2 showing statistically significant effect sizes.

3.4.1 Quadriceps Femoris

When comparing the ACL deficient limbs to the contralateral uninjured limbs, there was a moderate effect for both quadriceps femoris muscle CSA ($d = -0.35$; 95% CI -0.59 to -0.11; I² 0%, Figure 2a) and volume ($d = -0.40$; 95% CI -0.78 to -0.02; I² 18%, Figure 2b) indicating reduced muscle size in deficient limb. All other muscles and groups showed no statistically significant effects (Table 3 and 4).

3.5 ACL Reconstructed Populations

A summary of the results of the meta-analyses comparing muscle CSA and volume of the ACL reconstructed limbs to the contralateral uninjured limbs can be found in tables 5 and 6, respectively. Of those included, 16 measures of muscle CSA and 12 measures of muscle volume were analysed, with 9 showing statistically significant effect sizes.

3.5.1 Quadriceps Femoris

There was a moderate effect for both quadriceps femoris CSA and volume, indicating reduced muscle size in the reconstructed limbs when compared to the contralateral uninjured limb (Figure 3). For quadriceps femoris CSA data, time subgrouping analysis showed large effects between limbs at 6 to 9 weeks and 26 to 30 weeks, whereas moderate effects were seen between limbs at both 52 to 86 weeks and 156 to 289 weeks (Figure 3a). Studies included in the quadriceps femoris CSA meta-analysis had participants with a mix of harvest sites used for ACL reconstruction (patella tendon, iliotibial band and semitendinosus-gracilis tendon). For quadriceps femoris volume data, time subgrouping showed a large effect between limbs at 4 to 12 weeks (Figure 3b). Studies included in the quadriceps femoris volume meta-analyses had participants with a mix of harvest sites used for ACL reconstruction (patella tendon, semitendinosus tendon and semitendinosus-gracilis tendon). Meta-regression analysis found no significant effect for time since surgery on quadriceps femoris CSA, however, a significant positive effect was seen for volume in the ACL reconstructed limbs indicating differences between limbs decreased over time. (Electronic Supplementary Material Figure S1; intercept -1.245, $p = 0.0002$; coefficient 0.517, $p = 0.008$).

3.5.2 Vastus Medialis

A small effect was found for vastus medialis volume, indicating reduced muscle size in the reconstructed limb when compared to the contralateral uninjured limb (Figure 4a). Time subgrouping showed a moderate effect between limbs at 45 to 86 weeks (Figure 4a). Studies included in the meta-analyses had participants with a mix of harvest sites used for ACL reconstruction (patella tendon, semitendinosus tendon and semitendinosus-gracilis tendon). Meta-regression analysis showed no significant effect for time since surgery on vastus medialis volume in the ACL reconstructed limb (Electronic Supplementary Material Table S4).

3.5.3 Vastus Lateralis

Similar to the vastus medialis, a small effect was found for vastus lateralis volume, indicating reduced muscle size in the reconstructed limb when compared to the contralateral uninjured limb (Figure 4b). Time subgrouping showed no effects between limbs at any subgroup (Figure 4b). Studies included in the meta-analyses had participants with a mix of harvest sites used for ACL reconstruction (patella tendon, semitendinosus tendon and semitendinosus-gracilis tendon). Meta-regression analysis showed no significant effect for time since surgery on vastus lateralis volume in the ACL reconstructed limb (Electronic Supplementary Material Table S4).

3.5.4 Hamstrings

A small effect was found for total hamstring muscle volume indicating reduced muscle size in the reconstructed limb when compared to the contralateral uninjured limb (Figure 5). Time subgrouping analysis showed no significant effects between limbs for any time subgroup (Figure 5). Studies included in the hamstring meta-analysis had participants with either semitendinosus or semitendinosus-gracilis harvested for ACL reconstruction. Meta-regression analysis also showed no significant effect for time since surgery on total hamstring muscle volume in the ACL reconstructed limbs (Electronic Supplementary Material Table S4).

3.5.5 Semitendinosus

A large effect was found for semitendinosus muscle CSA and volume, indicating reduced muscle size in the reconstructed limb when compared to the contralateral uninjured limb (Figure 6). For semitendinosus CSA, time subgrouping analysis showed a large effect at both 52 to 53 weeks and 156 to 348 weeks post-surgery, whilst a moderate effect between limbs was seen at 26 to 30 weeks (Figure 6a). For semitendinosus muscle volume, time subgrouping showed a large effect between limbs at 26 weeks, 100 to 156 and 212 to 348 weeks post-surgery (Figure 6b). All studies included in the meta-analyses for semitendinosus CSA and volume had participants with semitendinosus tendon harvested for ACL reconstruction. Meta-regression analysis showed no effect for time since surgery on semitendinosus muscle CSA or volume in the ACL reconstructed limbs (Electronic Supplementary Material Table S4).

3.5.6 Gracilis

Large and moderate effects for gracilis muscle CSA and volume were found, indicating reduced muscle size in the reconstructed limb when compared to the contralateral uninjured limb (Figure 7). For gracilis CSA data, time subgrouping showed large effects between limbs at 52 to 53 weeks and 104 to 348 weeks post-surgery (Figure 7a). For gracilis muscle volume data, time subgrouping analysis showed a moderate effect at 156 to 348 weeks

post-surgery (Figure 7b). All studies included in the meta-analysis for gracilis CSA and volume had participants with gracilis tendon harvested for ACL reconstruction. Meta-regression analysis showed no effect for time since surgery on gracilis CSA or volume in the ACL reconstructed limbs (Electronic Supplementary Material Table S4).

3.6 Best Evidence Synthesis

Of the 49 articles included in this review, 12 were not in the above meta-analyses for the following reasons:

- Data not able to be obtained (n=8) [24, 75-81]
- Unique measures of ultrasound muscle size that were not grouped for meta-analysis (n=4) [82-85].

For these articles a best evidence synthesis (BES) was undertaken (Table 7). Due to the low number of studies, all of the muscle size measures were combined to obtain a single outcome for each muscle and group.

3.6.1 Quadriceps – ACL Reconstructed Limb Compared to Contralateral Uninjured Limb

When comparing the ACL reconstructed limbs to the contralateral uninjured limbs there was strong evidence to suggest reduced muscle size in the reconstructed limbs quadriceps femoris, vastus lateralis and rectus femoris (Table 7). There was also moderate evidence to suggest reduced muscle size in the vastus medialis and vastus intermedius. Similar to the meta-analyses for these muscles, studies included participants with mixed graft types (patella tendon, quadriceps tendon, iliotibial tract, semitendinosus tendon and semitendinosus-gracilis tendon).

3.6.2 Knee Flexors – ACL Reconstructed Limb Compared to Contralateral Uninjured Limb

There was moderate and strong evidence to suggest reduced muscle size in the reconstructed limbs semitendinosus and gracilis muscles of the ACL reconstructed limbs, respectively, when compared to the contralateral uninjured limbs. There was also limited and strong evidence to suggest no change in semimembranosus and biceps femoris ultrasound derived muscle size, respectively (Table 7). Similar to the meta-analysis on these measures, studies included or provided subgroup results for participants with the semitendinosus and/or gracilis tendon harvested for reconstruction.

3.6.3 Gluteus maximus – ACL Reconstructed Limb Compared to Contralateral Uninjured Limb

When comparing the ACL reconstructed limbs to the contralateral uninjured limbs there was moderate evidence to suggest reduced muscle size in the gluteus maximus of the reconstructed limb (Table 7). The single study that presented this data included participants with mixed graft types (patella tendon and semitendinosus-gracilis tendon).

3.6.4 Hamstrings - ACL Reconstructed Limb Compared to Healthy Control Group

When comparing the ACL reconstructed limb to a healthy control group, there was strong evidence to suggest an increase in semitendinosus size in the reconstructed limb (Table 7), however, studies included participants with mixed graft types (patella tendon and semitendinosus tendon). There was also moderate evidence to suggest reduced muscle size in the biceps femoris of the reconstructed limb from a single study that included participants with patella tendon grafts. Finally, there is moderate evidence to suggest there is no difference in

semimembranosus size between the groups with the one study using this measure including participants with patella tendon grafts.

4 Discussion

4.1 Statement of Main Findings

The main findings of this systematic review and meta-analyses are:

1) ACL deficient limbs (without subsequent surgical repair) have lower quadriceps femoris CSA and volume than the contralateral uninjured limb.

2) Regardless of graft site, ACL reconstructed limbs have lower quadriceps femoris CSA and deficits in quadriceps femoris, vastus medialis and vastus lateralis volume compared to the contralateral uninjured limb.

3) ACL reconstructed limbs have lower semitendinosus and gracilis CSA and muscle volume as well as lower total hamstring muscle volume compared to the contralateral uninjured limbs, when the respective tendons are harvested for reconstruction.

4.2 ACL Deficient Limbs

Suffering an ACL rupture causes knee joint instability, which is commonly treated with a surgical reconstruction. However, there is evidence to suggest approximately 25% of ACL injured individuals have successful two year outcomes and comparable knee function to non-injured individuals, without reconstructive surgery [86]. Conservative management of ACL injury carries some benefits over a surgical approach, namely avoiding the trauma and extensive healing time-frame associated with reconstructive surgery, as well as graft site morbidity. However, the results from the current meta-analysis highlight the atrophic effects the initial injury has on selected lower limb muscles. In ACL deficient limbs, quadriceps femoris muscle size appears significantly impacted from injury, showing reductions in both CSA and volume compared to the contralateral uninjured limb. It remains unclear if this reduction in quadriceps femoris size influences an individual's ability to successfully undergo conservative management.

4.3 ACL Reconstructed Populations

Similar to the ACL deficient limbs, reconstructed limbs showed significantly reduced quadriceps femoris muscle size. This is in agreement with the results from a recent meta-analysis that investigated changes in quadriceps CSA and volume following ACL reconstruction [39]. It is beyond the scope of this meta-analysis to assess any differences between graft types, but it appears all grafts (semitendinosus/gracilis and patella tendon) result in significantly reduced quadriceps size. This raises an important question of whether graft choice impacts any reduction in quadriceps size following ACL reconstruction. The meta-regression results suggested that quadriceps femoris CSA did not change as a function of time, but volume did appear to improve showing a reduction in effect size between limbs. This finding combined with the results from post-operative subgrouping in CSA and volume suggests that this reduction in muscle size may be most pronounced during early recovery, slowly returning to near that of the uninjured limb over the first two years following surgery.

Further breakdown of the individual muscles within the quadriceps group show that reduced muscle size is most likely concentrated in the vasti muscles, in particular vastus medialis and vastus lateralis which showed significantly reduced volume compared to the uninjured limb. Rectus femoris CSA and volume data showed no significant differences within the meta-analyses. However, the best evidence synthesis suggested strong evidence towards reduced rectus femoris muscle size in the ACL reconstructed limb when compared to the uninjured limb. These findings suggest that a percentage of ACL reconstructed individuals present with reduced rectus femoris size, though, it may not present as commonly as reductions in the vasti muscles. The reason for this remains unclear, however, it may be in part due to the differing action of the rectus femoris. Unlike the vasti muscles, the rectus femoris is a hip flexor, potentially exposing it to a unique stimulus (relative to the vasti) in the post-operative period and thus protecting it against atrophy.

In the ACL reconstructed limbs there was evidence of reduced total hamstring muscle volume when compared to the contralateral uninjured limb. This seems to be solely attributed to a large reduction in the semitendinosus muscle size, when its tendon is harvested for reconstruction. Similar to the quadriceps femoris CSA findings, semitendinosus meta-regression analysis showed no significant impact of time since surgery on semitendinosus CSA and volume. However, subgrouping for time suggests that reductions in semitendinosus muscle size occurs within the first few months following surgery but, unlike the quadriceps femoris findings, does not recover, with the latest available data (348 weeks post-surgery) showing large deficits in the reconstructed limb [66, 27].

Similarly, the gracilis of the ACL reconstructed limbs have significantly reduced CSA and volume when compared to the contralateral uninjured limb when the gracilis tendon is harvested for reconstruction. This is also strongly supported within the best evidence synthesis. Whilst the meta-regression-analyses showed no significant impact of time since surgery on gracilis CSA and volume, when subgrouping for time, the findings suggest that that a reduction in muscle size is apparent within the first few months following surgery and does not recover up to 348 weeks later.

4.4 Clinical Implications

Reductions in quadriceps muscle size is potentially impacted through two key mechanisms: 1) disuse atrophy associated with the period of unloading post-injury and surgery [87, 88]; 2) Neuromuscular inhibition, which lowers neural drive and fibre recruitment, leading to a lessened stimulus for adaptation, which is shown to significantly impact the quadriceps post-injury and surgery [28, 26, 23]. The quadriceps contribute the majority of force to knee joint compressive loading [35], as well supporting, braking and redirecting the centre of mass during sports specific movements such as side step cutting [89]. Quadriceps dysfunction therefore may impact the development of both knee joint osteoarthritis, and potential for secondary ACL injury. Addressing the atrophy of the quadriceps muscle group in ACL deficient and reconstructed limbs, through exercise based interventions should be of high importance and started as early as possible. However, care needs to be taken when aiming to restore quadriceps muscle size following reconstruction as time is needed for biological healing post-surgery, as well as allowing for the incorporation of grafted tissue into the ACL. Exposure to eccentrically biased [75] and blood flow restricted resistance training [90, 91] have been shown to be safe and effective approaches to improve muscle size in the early post-operative period. Subsequently both of these approaches represent an opportunity to

address the potential quadriceps atrophy in the early period post-surgery and prepare the individual for the higher resistance training loads in the later parts of the program which will also promote optimum muscle mass gains.

Restoration of semitendinosus and gracilis size following tendon harvest for ACL reconstruction may be limited, as the structural integrity of the muscle-tendon unit is severely disrupted due to the surgical intervention. This is supported by our results with the longest period post-surgery that was available (348 weeks) still showing significant deficits in the ACL reconstructed limb. The choice of graft remains a key consideration for surgeons to factor in when initially choosing a harvest site for reconstruction, as it is possible that the reduction in size and therefore the force producing capacity of the semitendinosus and gracilis may be permanent. Although work appears to be underway looking to address these deficits [92], to date, the authors are not aware of any published exercise interventions showing a restoration of semitendinosus and gracilis muscle size following tendon harvest for ACL reconstruction. Future research should continue to investigate if exercise based interventions in these muscles might be a useful approach to offset the extent of what seems inevitable atrophy.

4.5 Quality Assessment

Overall, only 55% of articles included in this review were deemed to be of a high quality. There were two areas of poorer performance in the quality assessment that may have the potential to significantly impact the findings of included articles. These were: 1) there was a lack of control in post-operative rehabilitation, with only 55% of all articles included in this review controlling for rehabilitation and 2) there were low percentages of external validity, with only 22% reporting the source of the population and how they were selected, and only 8% reporting the proportion of participants asked who agreed to participate. The low percentage scoring in these areas suggests that a number of articles included in this review may be unintentionally biasing their results by not controlling for these factors.

4.6 Limitations

One of the main limitations of this systematic review was an inability to split any meta-analyses based on the graft type of included participants. Advantageously, the studies included in the meta-analyses undertaken on hamstring volume, as well as semitendinosus and gracilis volume and CSA, were made up of studies which only included participants with semitendinosus/gracilis tendon harvests. The lack of data within the quadriceps specific studies limited the ability to split by graft types, without large reductions in the power of the meta-analyses. Furthermore, some studies included mixed cohorts with multiple graft types. The lack of data within some meta-analyses was a further limitation, therefore certain results may have suffered from sparse data bias. Additionally, visual inspection of funnels plots for meta-analysis with over ten included articles showed evidence of potential publication bias (Electronic Supplementary Material Figure S2). However, we reported all analysis to provide transparency of our methods and limit biased reporting. Another limitation was the time-point subgrouping of data from individual studies. Grouping was done via the mean ‘time since surgery’ values reported, however, studies often included populations from a larger range (e.g. six to twelve months). Additionally, the vast majority of comparisons made were between the injured limb and the contralateral uninjured limb. As such, the true reduction in size of muscles may be masked, as changes in strength, activation and functional performance are known to occur bilaterally in ACL injured individuals [38, 93] and the same may be true of CSA and volume.

There was also a strong focus from the majority of articles towards quadriceps, hamstrings and gracilis size. No included articles investigating ACL deficient populations included measures of muscle size from the lower leg or proximal hip. Only two articles investigating ACL reconstructed individuals included measures of gastrocnemius size, and only one article included measures of gluteus maximus size. Emerging research has highlighted the importance of the muscles of the shank and hip in opposing anterior shear and valgus knee forces [36, 37, 89] as well as contributing to knee joint compressive loading [35]. If these muscles groups are shown to be reduced in size following ACL injury, this may pose serious implications towards both the development of knee joint osteoarthritis and subsequent injury. Future work should look to investigate the effect of ACL injury and surgery on these muscles, specifically the gluteals, gastrocnemius and soleus.

5 Conclusion

This review highlights the overall differences in lower limb muscle size when comparing the ACL injured limb to the uninjured contralateral limb, in ACL reconstructed and deficient populations. Regardless of whether an individual chooses a conservative or surgical rehabilitation approach, the quadriceps femoris of the injured limb appear to show significantly reduced muscle size in the short-term, with the potential to recover to levels matching their contralateral limb in the long-term. However, if an individual undergoes reconstructive surgery with a semitendinosus and/or gracilis tendon graft, the harvested muscle display long term deficits in muscle size that may not be fully reversible. These findings suggest the need to focus the hypertrophic plans of rehabilitation around regaining quadriceps femoris muscle size, whilst considering the potential for possible permanent reductions in semitendinosus and gracilis muscle size if these tendons are harvested for reconstruction.

Acknowledgments

We would like to thank Jessica Dickson, Senior Librarian at Australian Catholic University, Melbourne, Australia, for help preparing the literature search strategy.

Table and Figure Captions

Table 1: Key search terms. Boolean term OR was used within categories, AND was used between categories.

Table 2: Inclusion and exclusion criteria applied to retrieved articles.

Table 3: Meta-analysis results for muscle cross sectional area of the ACL deficient limb compared to the contralateral uninjured limb.

Table 4: Meta-analysis results for muscle volume of the ACL deficient limb compared to the contralateral uninjured limb.

Table 5: Meta-analysis results for muscle cross sectional area of the ACL reconstructed limb compared to the contralateral uninjured limb.

Table 6: Meta-analysis results for muscle volume of the ACL reconstructed limb compared to the contralateral uninjured limb.

Table 7: Results of the best evidence synthesis for ACL reconstructed populations.

Figure 1: Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) flowchart outlining study selection process.

Figure 2: Results of the meta-analysis for a) quadriceps femoris muscle cross sectional area, and b) quadriceps femoris muscle volume, in the ACL deficient limb compared to the contralateral uninjured limb. Negative effect size indicates the ACL injured limb measure is less than the contralateral uninjured limb.

Figure 3: Results of the meta-analysis for a) quadriceps femoris cross sectional area, and b) quadriceps femoris muscle volume, in the ACL reconstructed limb compared to the contralateral uninjured limb. Negative effect size indicates the ACL injured limb measure is less than the contralateral uninjured limb.

Figure 4: Results of the meta-analysis for a) vastus medialis, and b) vastus lateralis muscle volume, in the ACL reconstructed limb compared to the contralateral uninjured limb. Negative effect size indicates the ACL injured limb measure is less than the contralateral uninjured limb.

Figure 5: Results of the meta-analysis for total hamstring muscle volume, in the ACL reconstructed limb compared to the contralateral uninjured limb. Negative effect size indicates the ACL injured limb measure is less than the contralateral uninjured limb.

Figure 6: Results of the meta-analysis for, a) semitendinosus muscle cross sectional area, and b) semitendinosus muscle volume, in the ACL reconstructed limb compared to the contralateral uninjured limb. Negative effect size indicates the ACL injured limb measure is less than the contralateral uninjured limb.

Figure 7: Results of the meta-analysis for, a) gracilis muscle cross sectional area, and b) gracilis muscle volume in the ACL reconstructed limb compared to the contralateral uninjured limb. Negative effect size indicates the ACL injured limb measure is less than the contralateral uninjured limb.

Supplementary Content Captions

Electronic Supplementary Material Table S1: Modified quality assessment tool derived from Downs and Black [41].

Electronic Supplementary Material Table S2: Quality assessment scores of included studies.

Electronic Supplementary Material Table S3: Meta-analysis results for muscle volume of the ACL reconstructed limb compared to a healthy control group.

Electronic Supplementary Material Table S4: Meta-regression results comparing between the ACL reconstructed limb and the contralateral uninjured limb.

Electronic Supplementary Material Figure S1: Meta-regression plot comparing between the ACL reconstructed limb and the contralateral uninjured limb for quadriceps femoris muscle volume.

467 Electronic Supplementary Material Figure S2: Funnel plots assessing publication bias for meta-analyses with
468 >10 included studies. a) Quadriceps femoris cross sectional area, and b) semitendinosus cross sectional area. Trim-
469 fill (metafor) was used to estimate missing studies, however, both plots returned 0 inputted studies.

470

References

1. Ardern CL, Taylor NF, Feller JA, Webster KE. Fifty-five per cent return to competitive sport following anterior cruciate ligament reconstruction surgery: an updated systematic review and meta-analysis including aspects of physical functioning and contextual factors. *Br J Sports Med.* 2014;48(21):1543. doi:10.1136/bjsports-2013-093398.
2. Welling W, Benjaminse A, Seil R, Lemmink K, Zaffagnini S, Gokeler A. Low rates of patients meeting return to sport criteria 9 months after anterior cruciate ligament reconstruction: a prospective longitudinal study. *Knee Surg Sports Traumatol Arthrosc.* 2018;1-9. doi:10.1007/s00167-018-4916-4.
3. Sanders TL, Maradit Kremers H, Bryan AJ, Larson DR, Dahm DL, Levy BA, et al. Incidence of anterior cruciate ligament tears and reconstruction: a 21-year population-based study. *Am J Sports Med.* 2016;44(6):1502-7. doi:10.1177/0363546516629944.
4. Sutherland K, Clatworthy M, Fulcher M, Chang K, Young SW. Marked increase in the incidence of anterior cruciate ligament reconstructions in young females in New Zealand. *ANZ J Surg.* 2019;89(9):1151-5. doi:10.1111/ans.15404.
5. Abram SGF, Price AJ, Judge A, Beard DJ. Anterior cruciate ligament (ACL) reconstruction and meniscal repair rates have both increased in the past 20 years in England: Hospital statistics from 1997 to 2017. *Br J Sports Med.* 2020;54(5):286-91. doi:10.1136/bjsports-2018-100195.
6. Zbrojkiewicz D, Vertullo C, Grayson JE. Increasing rates of anterior cruciate ligament reconstruction in young Australians, 2000-2015. *Med J Aust.* 2018;208(8):354-8. doi:10.5694/mja17.00974.
7. Janssen KW, Orchard JW, Driscoll TR, van Mechelen W. High incidence and costs for anterior cruciate ligament reconstructions performed in Australia from 2003-2004 to 2007-2008: time for an anterior cruciate ligament register by Scandinavian model? *Scand J Med Sci Sports.* 2012;22(4):495-501. doi:10.1111/j.1600-0838.2010.01253.x.
8. Webster KE, Feller JA, Kimp AJ, Whitehead TS. Low Rates of Return to Preinjury Sport After Bilateral Anterior Cruciate Ligament Reconstruction. *Am J Sports Med.* 2018;0363546518813901. doi:10.1177/0363546518813901.
9. Webster KE, Hewett TE. What is the evidence for and validity of return-to-sport testing after anterior cruciate ligament reconstruction surgery? A systematic review and meta-analysis. *Sports Med.* 2019;49(6):917-29. doi:10.1007/s40279-019-01093-x.
10. Culvenor AG, Cook JL, Collins NJ, Crossley KM. Is patellofemoral joint osteoarthritis an under-recognised outcome of anterior cruciate ligament reconstruction? A narrative literature review. *Br J Sports Med.* 2013;47(2):66-70. doi:10.1136/bjsports-2012-091490.
11. Lohmander LS, Englund PM, Dahl LL, Roos EM. The long-term consequence of anterior cruciate ligament and meniscus injuries: osteoarthritis. *Am J Sports Med.* 2007;35(10):1756-69. doi:10.1177/0363546507307396.

505 12. Kim HJ, Lee JH, Ahn SE, Park MJ, Lee DH. Influence of anterior cruciate ligament tear on thigh muscle
506 strength and hamstring-to-quadriceps ratio: A meta-analysis. *PLoS One*. 2016;11(1).
507 doi:10.1371/journal.pone.0146234.

508 13. Larsen JB, Farup J, Lind M, Dalgas U. Muscle strength and functional performance is markedly impaired at
509 the recommended time point for sport return after anterior cruciate ligament reconstruction in recreational athletes.
510 *Hum Mov Sci*. 2015;39:73-87. doi:https://doi.org/10.1016/j.humov.2014.10.008.

511 14. Kuenze CM, Hertel J, Weltman A, Diduch D, Saliba SA, Hart JM. Persistent neuromuscular and corticomotor
512 quadriceps asymmetry after anterior cruciate ligament reconstruction. *J Athl Train*. 2015;50(3):303-12.
513 doi:10.4085/1062-6050-49.5.06.

514 15. Lepley AS, Gribble PA, Thomas AC, Tevald MA, Sohn DH, Pietrosimone BG. Quadriceps neural alterations
515 in anterior cruciate ligament reconstructed patients: a 6-month longitudinal investigation. *Scand J Med Sci Sports*.
516 2015;25(6):828-39. doi:10.1111/sms.12435.

517 16. Lepley LK, Palmieri-Smith RM. Pre-operative quadriceps activation is related to post-operative activation,
518 not strength, in patients post-ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc*. 2016;24(1):236-46.
519 doi:10.1007/s00167-014-3371-0.

520 17. Ward SH, Blackburn JT, Padua DA, Stanley LE, Harkey MS, Luc-Harkey BA, et al. Quadriceps
521 neuromuscular function and jump-landing sagittal-plane knee biomechanics after anterior cruciate ligament
522 reconstruction. *J Athl Train*. 2018;53(2):135-43. doi:10.4085/1062-6050-306-16.

523 18. Gumucio JP, Sugg KB, Enselman ERS, Konja AC, Eckhardt LR, Bedi A, et al. Anterior cruciate ligament tear
524 induces a sustained loss of muscle fiber force production. *Muscle Nerve*. 2018;58(1):145-8.
525 doi:10.1002/mus.26075.

526 19. Konrath JM, Vertullo CJ, Kennedy BA, Bush HS, Barrett RS, Lloyd DG. Morphologic characteristics and
527 strength of the hamstring muscles remain altered at 2 years after use of a hamstring tendon graft in anterior cruciate
528 ligament reconstruction. *Am J Sports Med*. 2016;44(10):2589-98.

529 20. Lindström M, Str, berg S, Wredmark T, Felländer-Tsai L, Henriksson M. Functional and muscle morphometric
530 effects of ACL reconstruction. A prospective CT study with 1 year follow-up. *Scand J Med Sci Sports*.
531 2013;23(4):431-42.

532 21. Nomura Y, Kuramochi R, Fukubayashi T. Evaluation of hamstring muscle strength and morphology after
533 anterior cruciate ligament reconstruction. *Scand J Med Sci Sports*. 2015;25(3):301-7.

534 22. Macleod TD, Snyder-Mackler L, Buchanan TS. Differences in neuromuscular control and quadriceps
535 morphology between potential copers and noncopers following anterior cruciate ligament injury. *J Orthop Sports*
536 *Phys Ther*. 2014;44(2):76-84.

537 23. Williams GN, Buchanan TS, Barrance PJ, Axe MJ, Snyder-Mackler L. Quadriceps weakness, atrophy, and
538 activation failure in predicted noncopers after anterior cruciate ligament injury. *Am J Sports Med*. 2005;33(3):402-
539 7.

540 24. Friedmann-Bette B, Profit F, Gwechenberger T, Weiberg N, Parstorfer M, Weber M-A, et al. Strength Training
541 Effects on Muscular Regeneration after ACL Reconstruction. *Med Sci Sports Exerc.* 2018;50(6):1152-61.

542 25. Grapar Žargi T, Drobnič M, Vauhnik R, Koder J, Kacin A. Factors predicting quadriceps femoris muscle
543 atrophy during the first 12weeks following anterior cruciate ligament reconstruction. *Knee.* 2017;24(2):319-28.

544 26. Lepley AS, Grooms DR, Burl, JP, Davi SM, Kinsella-Shaw JM et al. Quadriceps muscle function following
545 anterior cruciate ligament reconstruction: systemic differences in neural and morphological characteristics. *Exp*
546 *Brain Res.* 2019.

547 27. Messer DJ, Shield AJ, Williams MD, Timmins RG, Bourne MN. Hamstring muscle activation and
548 morphology are significantly altered 1-6 years after anterior cruciate ligament reconstruction with semitendinosus
549 graft. *Knee Surg Sports Traumatol Arthrosc.* 2020;28(3):733-41. doi:10.1007/s00167-019-05374-w.

550 28. Kuenze CM, Blemker SS, Hart JM. Quadriceps function relates to muscle size following ACL reconstruction.
551 *J Orthop Res.* 2016;34(9):1656-62.

552 29. Thomas AC, Wojtys EM, Br, on C, Palmieri-Smith RM. Muscle atrophy contributes to quadriceps weakness
553 after anterior cruciate ligament reconstruction. *J Sci Med Sport.* 2016;19(1):7-11.

554 30. Saxby DJ, Bryant AL, Modenese L, Gerus P, Killen BA, Konrath J, et al. Tibiofemoral contact forces in the
555 anterior cruciate ligament-reconstructed knee. *Med Sci Sports Exerc.* 2016;48(11):2195-206.
556 doi:10.1249/mss.0000000000001021.

557 31. Wellsandt E, Gardinier ES, Manal K, Axe MJ, Buchanan TS, Snyder-Mackler L. Decreased knee joint loading
558 associated with early knee osteoarthritis after anterior cruciate ligament injury. *Am J Sports Med.* 2016;44(1):143-
559 51. doi:10.1177/0363546515608475.

560 32. Pietrosimone B, Pfeiffer SJ, Harkey MS, Wallace K, Hunt C, Blackburn JT, et al. Quadriceps weakness
561 associates with greater T1rho relaxation time in the medial femoral articular cartilage 6 months following anterior
562 cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2018. doi:10.1007/s00167-018-5290-y.

563 33. Øiestad BE, Juhl CB, Eitzen I, Thorlund JB. Knee extensor muscle weakness is a risk factor for development
564 of knee osteoarthritis. A systematic review and meta-analysis. *Osteoarthritis Cartilage.* 2015;23(2):171-7.
565 doi:10.1016/j.joca.2014.10.008.

566 34. Ithurburn MP, Paterno MV, Ford KR, Hewett TE, Schmitt LC. Young athletes with quadriceps femoris
567 strength asymmetry at return to sport after anterior cruciate ligament reconstruction demonstrate asymmetric
568 single-leg drop-landing mechanics. *Am J Sports Med.* 2015;43(11):2727-37. doi:10.1177/0363546515602016.

569 35. Maniar N, Bryant AL, Sritharan P, Schache AG, Opar DA. Muscle contributions to medial and lateral
570 tibiofemoral compressive loads during sidestep cutting. *J Biomech.* 2020;101.
571 doi:10.1016/j.jbiomech.2020.109641.

572 36. Maniar N, Schache AG, Pizzolato C, Opar DA. Muscle contributions to tibiofemoral shear forces and valgus
573 and rotational joint moments during single leg drop landing. *Scand J Med Sci Sports.* 2020;30(9):1664-74.
574 doi:10.1111/sms.13711.

575 37. Maniar N, Schache AG, Sritharan P, Opar DA. Non-knee-spanning muscles contribute to tibiofemoral shear
576 as well as valgus and rotational joint reaction moments during unanticipated sidestep cutting. *Sci Rep*.
577 2018;8(1):2501. doi:10.1038/s41598-017-19098-9.

578 38. Lisee C, Lepley AS, Birchmeier T, O'Hagan K, Kuenze C. Quadriceps strength and volitional activation after
579 anterior cruciate ligament reconstruction: a systematic review and meta-analysis. *Sports Health*.
580 2019;1941738118822739. doi:10.1177/1941738118822739.

581 39. Birchmeier T, Lisee C, Kane K, Brazier B, Triplett A, Kuenze C. Quadriceps muscle size following acl injury
582 and reconstruction: a systematic review. *J Orth Res*. 2020;38(3):598-608. doi:10.1002/jor.24489.

583 40. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-
584 analyses: the PRISMA statement. *PLoS Med*. 2009;6(7):e1000097. doi:10.1371/journal.pmed.1000097.

585 41. Downs SH, Black N. The feasibility of creating a checklist for the assessment of the methodological quality
586 both of randomised and non-randomised studies of health care interventions. *J Epidemiol Community Health*.
587 1998;52(6):377-84. doi:10.1136/jech.52.6.377.

588 42. Maniar N, Shield AJ, Williams MD, Timmins RG, Opar DA. Hamstring strength and flexibility after
589 hamstring strain injury: A systematic review and meta-analysis. *Br J Sports Med*. 2016;50(15):909-20.
590 doi:10.1136/bjsports-2015-095311.

591 43. Grapar Zargi T, Drobic M, Jkoder J, Strazar K, Kacin A. The effects of preconditioning with ischemic
592 exercise on quadriceps femoris muscle atrophy following anterior cruciate ligament reconstruction: a quasi-
593 randomized controlled trial. *Eur J Phys Rehabil Med*. 2016;52(3):310-20.

594 44. Noehren B, Andersen A, Hardy P, Johnson DL, Irel ML, et al. Cellular and morphological alterations in the
595 vastus lateralis muscle as the result of acl injury and reconstruction. *J Bone Joint Surg Am*. 2016;98(18):1541-7.

596 45. Setuain I, Izquierdo M, Idoate F, Bik, i E, Gorostiaga EM, et al. Differential effects of 2 rehabilitation programs
597 following anterior cruciate ligament reconstruction. *J Sport Rehabil*. 2017;26(6):544-55.

598 46. Wigerstad-Lossing I, Grimby G, Jonsson T, Morelli B, Peterson L, Renström P. Effects of electrical muscle
599 stimulation combined with voluntary contractions after knee ligament surgery. *Med Sci Sport Exerc*.
600 1988;20(1):93-8.

601 47. Viechtbauer W. Conducting meta-analyses in R with the metafor package. 2010. 2010;36(3):48.
602 doi:10.18637/jss.v036.i03.

603 48. Schwarzer G. Meta: general package for meta-analysis. 2015.

604 49. Cohen J. A power primer. *Psychol Bull*. 1992;112(1):155-9. doi:10.1037//0033-2909.112.1.155.

605 50. Slavin RE. Best evidence synthesis: an intelligent alternative to meta-analysis. *J Clin Epidemiol* 1995;48(1):9-
606 18. doi:10.1016/0895-4356(94)00097-a.

607 51. Arangio GA, Chen C, Kalady M, Reed JF, 3rd. Thigh muscle size and strength after anterior cruciate ligament
608 reconstruction and rehabilitation. *J Orthop Sports Phys Ther*. 1997;26(5):238-43.

609 52. Arvidsson I, Arvidsson H, Eriksson E, Jansson E. Prevention of quadriceps wasting after immobilization: an
610 evaluation of the effect of electrical stimulation. *Orthopedics*. 1986;9(11):1519-28.

611 53. Burks RT, Crim J, Fink BP, Boylan DN, Greis PE. The effects of semitendinosus and gracilis harvest in
612 anterior cruciate ligament reconstruction. *Arthroscopy*. 2005;21(10):1177-85.

613 54. Eriksson K, Hamberg P, Jansson E, Larsson H, Shalabi A, Wredmark T. Semitendinosus muscle in anterior
614 cruciate ligament surgery: Morphology and function. *Arthroscopy*. 2001;17(8):808-17.

615 55. Flück M, Vieceilli C, Bapst AM, Kasper S, Valdivieso P, Franchi MV, et al. Knee extensors muscle plasticity
616 over a 5-years rehabilitation process after open knee surgery. *Front Physiol*. 2018;9:1343-.

617 56. Gandolfi M, Ricci M, Sambugaro E, Valè N, Dimitrova E, Meschieri A, et al. Changes in the sensorimotor
618 system and semitendinosus muscle morphometry after arthroscopic anterior cruciate ligament reconstruction: a
619 prospective cohort study with 1-year follow-up. *Knee Surg Sports Traumatol Arthrosc*. 2018;26(12):3770-9.
620 doi:10.1007/s00167-018-5020-5.

621 57. Garcia SA, Curran MT, Palmieri-Smith RM. Longitudinal assessment of quadriceps muscle morphology
622 before and after anterior cruciate ligament reconstruction and its associations with patient-reported outcomes.
623 *Sports Health*. 2020:1941738119898210-. doi:10.1177/1941738119898210.

624 58. Janssen RPA, van der Velden MJF, Pasmans HLM, Sala HAGM. Regeneration of hamstring tendons after
625 anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc*. 2013;21(4):898-905.

626 59. Kariya Y, Itoh M, Nakamura T, Yagi K, Kurosawa H. Magnetic resonance imaging and spectroscopy of thigh
627 muscles in cruciate ligament insufficiency. *Acta Orthop Scand*. 1989;60(3):322-5.

628 60. Konishi Y, Fukubayashi T. Relationship between muscle volume and muscle torque of the hamstrings after
629 anterior cruciate ligament reconstruction. *J Sci Med Sport*. 2010;13(1):101-5.

630 61. Konishi Y, Ikeda K, Nishino A, Sunaga M, Aihara Y, Fukubayashi T. Relationship between quadriceps
631 femoris muscle volume and muscle torque after anterior cruciate ligament repair. *Scand J Med Sci Sports*.
632 2007;17(6):656-61.

633 62. Konishi Y, Kinugasa R, Oda T, Tsukazaki S, Fukubayashi T. Relationship between muscle volume and muscle
634 torque of the hamstrings after anterior cruciate ligament lesion. *Knee Surg Sports Traumatol Arthrosc*.
635 2012;20(11):2270-4.

636 63. Konishi Y, Oda T, Tsukazaki S, Kinugasa R, Fukubayashi T. Relationship between quadriceps femoris muscle
637 volume and muscle torque at least 18 months after anterior cruciate ligament reconstruction. *Scand J Med Sci*
638 *Sports*. 2012;22(6):791-6.

639 64. Lopresti C, Kirkendall DT, Street GM, Dudley AW. Quadriceps insufficiency following repair of the anterior
640 cruciate ligament. *J Orthop Sports Phys Ther*. 1988;9(7):245-9.

641 65. Lorentzon R, Elmqvist LG, Sjöström M, Fagerlund M, Fuglmeyer AR. Thigh musculature in relation to
642 chronic anterior cruciate ligament tear: muscle size, morphology, and mechanical output before reconstruction.
643 *Am J Sports Med.* 1989;17(3):423-9.

644 66. Macleod TD, Snyder-Mackler L, Axe MJ, Buchanan TS. Early regeneration determines long-term graft site
645 morphology and function after reconstruction of the anterior cruciate ligament with semitendinosus-gracilis
646 autograft: a case series. *Int J Sports Phys Ther* . 2013;8(3):256-68.

647 67. Marcon M, Ciritsis B, Laux C, Nanz D, Fischer MA, Andreisek G, et al. Quantitative and qualitative MR-
648 imaging assessment of vastus medialis muscle volume loss in asymptomatic patients after anterior cruciate
649 ligament reconstruction. *J Magn Reson Imaging.* 2015;42(2):515-25.

650 68. Marcon M, Ciritsis B, Laux C, Nanz D, Nguyen-Kim TDL, Fischer MA et al. Cross-sectional area
651 measurements versus volumetric assessment of the quadriceps femoris muscle in patients with anterior cruciate
652 ligament reconstructions. *Eur Radiol.* 2014;25(2):290-8.

653 69. Nishino A, Sanada A, Kanehisa H, Fukubayashi T. Knee-flexion torque and morphology of the semitendinosus
654 after ACL reconstruction. *Med Sci Sports Exerc.* 2006;38(11):1895-900.

655 70. Reeves ND, Maganaris CN, Maffulli N, Rittweger J. Human patellar tendon stiffness is restored following
656 graft harvest for anterior cruciate ligament surgery. *J Biomech.* 2009;42(7):797-803.

657 71. Simonian PT, Harrison SD, Cooley VJ, Escabedo EM, Deneka DA, Larson RV. Assessment of morbidity of
658 semitendinosus and gracilis tendon harvest for ACL reconstruction. *Am J Knee Surg.* 1997;10(2):54-9.

659 72. Strandberg S, Lindström M, Wretling M-L, Aspelin P, Shalabi A. Muscle morphometric effect of anterior
660 cruciate ligament injury measured by computed tomography: aspects on using non-injured leg as control. *BMC*
661 *Musculoskelet Disord.* 2013;14:150-.

662 73. Williams GN, Snyder-Mackler L, Barrance PJ, Axe MJ, Buchanan TS. Muscle and tendon morphology after
663 reconstruction of the anterior cruciate ligament with autologous semitendinosus-gracilis graft. *J Bone Joint Surg*
664 *Am.* 2004;86(9):1936-46.

665 74. Williams GN, Snyder-Mackler L, Barrance PJ, Buchanan TS. Quadriceps femoris muscle morphology and
666 function after ACL injury: a differential response in copers versus non-copers. *J Biomech.* 2005;38(4):685-93.

667 75. Gerber JP, Marcus RL, Dibble L, E., Greis PE, Burks RT, et al. Effects of early progressive eccentric exercise
668 on muscle structure after anterior cruciate ligament reconstruction. *J Bone Joint Surg Am.* 2007;89(3):559-70.

669 76. Hunnicutt JL, Gregory CM, McLeod MM, Woolf SK, Chapin RW, Slone HS. Quadriceps recovery after
670 anterior cruciate ligament reconstruction with quadriceps tendon versus patellar tendon autografts. *Orthop J Sports*
671 *Med.* 2019;7(4):1-7. doi:10.1177/2325967119839786.

672 77. Hunnicutt JL, McLeod MM, Slone HS, Gregory CM. Quadriceps muscle strength, size, and activation and
673 physical function after anterior cruciate ligament reconstruction. *J Athl Train.* 2020. doi:10.4085/1062-6050-516-
674 18.

78. Irie K, Tomatsu T. Atrophy of semitendinosus and gracilis and flexor mechanism function after hamstring tendon harvest for anterior cruciate ligament reconstruction. *Orthopedics*. 2002;25(5):491-5.
79. Kellis E, Karagiannidis E, Patsika G. Patellar tendon and hamstring moment-arms and cross-sectional area in patients with anterior cruciate ligament reconstruction and controls. *Comput Methods Biomech Biomed Engin*. 2015;18(10):1083-9.
80. Snow BJ, Wilcox JJ, Burks RT, Greis PE. Evaluation of muscle size and fatty infiltration with MRI nine to eleven years following hamstring harvest for ACL reconstruction. *J Bone Joint Surg Am*. 2012;94(14):1274-82.
81. Takahashi K, Hayashi M, Fujii T, Kawamura K, Ozaki T. Early rehabilitation with weight-bearing standing-shaking-board exercise in combination with electrical muscle stimulation after anterior cruciate ligament reconstruction. *Acta Med Okayama*. 2012;66(3):231-7.
82. Karagiannidis E, Kellis E, Galanis N, Vasilios B. Semitendinosus muscle architecture during maximum isometric contractions in individuals with anterior cruciate ligament reconstruction and controls. *Muscles Ligaments Tendons J*. 2017;7(1):147-51.
83. Kilgas MA, Lytle LLM, Drum SN, Elmer SJ. Exercise with blood flow restriction to improve quadriceps function long after acl reconstruction. *International Journal of Sports Medicine*. 2019;40(10):650-6. doi:10.1055/a-0961-1434.
84. Longo UG, Rizzello G, Frnaceschi F, Campi S, Maffulli N, Denaro V. The architecture of the ipsilateral quadriceps two years after successful anterior cruciate ligament reconstruction with bone-patellar tendon-bone autograft. *Knee*. 2014;21(3):721-5.
85. Timmins RG, Bourne MN, Shield AJ, Williams MD, Lorenzen C, Opar DA. Biceps femoris architecture and strength in athletes with a previous anterior cruciate ligament reconstruction. *Med Sci Sports Exerc*. 2016;48(3):337-45.
86. Grindem H, Wellsandt E, Failla M, Snyder-Mackler L, Risberg MA. Anterior cruciate ligament injury-who succeeds without reconstructive surgery? The Delaware-Oslo acl cohort study. *Orthop J Sports Med*. 2018;6(5):2325967118774255. doi:10.1177/2325967118774255.
87. Bodine SC. Disuse-induced muscle wasting. *Int J Biochem Cell Biol*. 2013;45(10):2200-8. doi:10.1016/j.biocel.2013.06.011.
88. Brooks N, Myburgh K. Skeletal muscle wasting with disuse atrophy is multi-dimensional: the response and interaction of myonuclei, satellite cells and signaling pathways. *Front Physiol*. 2014;5(99). doi:10.3389/fphys.2014.00099.
89. Maniar N, Schache AG, Cole MH, Opar DA. Lower-limb muscle function during sidestep cutting. *J Biomech*. 2018. doi:https://doi.org/10.1016/j.jbiomech.2018.10.021
90. Hughes L, Rosenblatt B, Haddad F, Gissane C, McCarthy D, Clarke T, et al. Comparing the effectiveness of blood flow restriction and traditional heavy load resistance training in the post-surgery rehabilitation of anterior

709 cruciate ligament reconstruction patients: a UK national health service randomised controlled trial. *Sports Med.*
710 2019;49(11):1787-805.

711 91. Ohta H, Kurosawa H, Ikeda H, Iwase Y, Satou N, Nakamura S. Low-load resistance muscular training with
712 moderate restriction of blood flow after anterior cruciate ligament reconstruction. *Acta Orthop Scand.*
713 2003;74(1):62-8.

714 92. Bregenhof B, Jørgensen U, Aagaard P, Nissen N, Creaby MW, Thorlund JB et al. The effect of targeted
715 exercise on knee-muscle function in patients with persistent hamstring deficiency following ACL reconstruction
716 – study protocol for a randomized controlled trial. *Trials.* 2018;19(1):75. doi:10.1186/s13063-018-2448-3.

717 93. Patterson BE, Crossley KM, Perraton LG, Kumar AS, King MG, Heerey JJ et al. Limb symmetry index on a
718 functional test battery improves between one and five years after anterior cruciate ligament reconstruction,
719 primarily due to worsening contralateral limb function. *Phys Ther Sport.* 2020;44:67-74.
720 doi:<https://doi.org/10.1016/j.ptsp.2020.04.031>.