#### SPORTS MEDICINE AND BIOMECHANICS

OPEN ACCESS Check for updates

# The application of the Lyapunov Exponent to analyse human performance: A systematic review

Lachlan Winter (D<sup>a,b</sup>, Paul Taylor<sup>c</sup>, Clint Bellenger<sup>a,b</sup>, Paul Grimshaw<sup>d,e</sup> and Robert G. Crowther<sup>a,b,f</sup>

<sup>a</sup>UniSA Allied Health and Human Performance, University of South Australia, Adelaide, South Australia, Australia; <sup>b</sup>Alliance for Research in Exercise, Nutrition & Activity (ARENA), University of South Australia, Adelaide, South Australia, Australia; <sup>c</sup>School of Behavioural and Health Sciences, Australian Catholic University, North Sydney, New South Wales, Australia; <sup>d</sup>College of Health and Life Sciences, Hamad Bin Khalifa University, Doha, Qatar; <sup>e</sup>Faculty of Sciences, Engineering and Technology, Computer and Mathematical Sciences, University of Adelaide, Adelaide, South Australia; <sup>f</sup>School of Behavioural and Health Sciences, Australian Catholic University, Melbourne, Victoria, Australia

#### ABSTRACT

Variability is a normal component of human movement, allowing one to adapt to environmental perturbations. It can be analysed from linear or non-linear perspectives. The Lyapunov Exponent (LyE) is a commonly used non-linear technique, which quantifies local dynamic stability. It has been applied primarily to walking gait and appears to be limited application in other movements. Therefore, this systematic review aims to summarise research methodologies applying the LyE to movements, excluding walking gait. Four databases were searched using keywords related to movement variability, dynamic stability, LyE and divergence exponent. Articles written in English, using the LyE to analyse movements, excluding walking gait were included for analysis. 31 papers were included for data extraction. Quality appraisal was conducted and information related to the movement, data capture method, data type, apparatus, sampling rate, body segment/joint, number of strides/steps, state space reconstruction, algorithm, filtering, surrogation and time normalisation were extracted. LyE values were reported in supplementary materials (Appendix 2). Running was the most prevalent non-walking gait movement assessed. Methodologies to calculate the LyE differed in various aspects resulting in different LyE values being generated. Additionally, test-retest reliability, was only conducted in one study, which should be addressed in future.

## Introduction

Movement variability has previously been viewed as measurement noise which should be eliminated (Stergiou et al., 2004). As such successful skill execution was previously characterised by a lack of movement variability, since less variability in outcome measures like error and accuracy represents better performance (Davids et al., 2006). Research has since challenged this notion, positing that a low outcome variability does not necessarily coincide with low technique variability (Preatoni et al., 2013).

Dynamic systems theory (DST) is a human movement theory that corroborates the idea that variability is a functional component of skill, suggesting that movement patterns are created from the collaborative organisation of the neuromuscular system based on the interaction between task (e.g., goals and rules), environmental (e.g., weather, spectators) and organism (individual anthropometry and morphology) constraints (Bernstein, 1967; Davids et al., 2006; Dingwell & Cusumano, 2000; Hamill et al., 1999). Alterations in these constraints may alter movement patterns. DST acknowledges the inherent variability that exists in human movement, recognising that movement variations occur naturally, and allow individuals to adapt to new situations (Davids et al., 2006; Magill & Anderson, 2017; Stergiou & Decker, 2011). For example, individuals alter their walking gait in response to different environmental factors. For example, to compensate for the moving belt of a treadmill, individuals implement a more cautious walking gait by having smaller step lengths and slower self-selected speeds when walking on a treadmill compared to overground walking (Yang & King, 2016). Stergiou and Decker (2011) suggest that a healthy bandwidth of variability exists, where individuals can successfully adapt to novel situations. Beyond this bandwidth, individuals are too variable, resulting in instability, whereas too little variability results in a rigid system, that appears robotic; both of which are unhealthy and do not allow one to successfully manage perturbations.

Variability is also present in sporting movements. For example, basketballers alter their shooting movement pattern to make a basket in response to different environmental factors like the distance from the basket or the presence of defenders (Slegers et al., 2021). Additionally, whether movement variability is desired, depends on the task and its specific context (Davids et al., 2006; Stergiou & Decker, 2011). For example, reducing shoulder joint angular kinematic variability is associated with improved pitch location control in baseball pitching (Glanzer et al., 2021), but in baseball hitting, elite players exhibit

© 2024 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

#### **ARTICLE HISTORY**

Received 11 September 2023 Accepted 15 January 2024

## KEYWORDS

Dynamic stability; movement variability: Lyapunov Exponent



CONTACT Lachlan Winter 🖾 Lachlan.Winter@mymail.unisa.edu.au 🗈 UniSA Allied Health and Human Performance, University of South Australia, Campus Central -City East, GPO Box 2471, Adelaide, South Australia SA 5001, Australia

The present affiliation of Robert G. Crowther is School of Science and Technology, University of New England, Armidale, NSW, Australia

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (http://creativecommons.org/licenses/by-nc-nd/4.0/), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way. The terms on which this article has been published allow the posting of the Accepted Manuscript in a repository by the author(s) or with their consent.

greater timing variability in their swing than novices to adapt to different pitch types and locations (Gray, 2020).

However, how variability is quantified influences the interpretation of the results. Variability has typically been guantified with simple statistical measures like the coefficient of variation and standard deviation, which assess variability across multiple trials after generating an average (Davids et al., 2006; James, 2004). However, these measures only quantify the magnitude of variability, and assume a "typical" variability exists, without reference to how movement patterns exhibit variability over time (Caballero et al., 2014; James, 2004). Additionally, they assume variability is a deviation from an average, representing error (Caballero et al., 2014). Furthermore, by taking only a single measure from a continuous variable, such as kinematics from a walking gait, a large amount of information is discarded, resulting in the loss of potentially useful information (Preatoni et al., 2013). As such, linear measures of variability do not evaluate the structure of variability. Another limitation of linear measures is that they can only measure one of the temporal or spatial aspects of movement, whilst neglecting the other component (Longo et al., 2018).

Consequently, DST has led to a rise in the use of nonlinear analysis methods, which can characterise the structure of variability because they acknowledge the deterministic origin (i.e., different constraints) and the influence of constraint alteration on movement patterns (Hamill et al., 1999; Stergiou et al., 2004; Van Emmerik et al., 2004). As such, nonlinear analysis tools recognise that variability is inherent and important to functional human movement (Estep et al., 2017). Broadly, nonlinear tools analyse the repeated cycles of movement over time and attempt to evaluate the dynamics causing the changes that occur between cycles (Preatoni et al., 2013).

Various non-linear analysis techniques exist (and debate exists amongst researchers about what constitutes a nonlinear tool), each examining a unique aspect of data and thus comparisons should only be made between the same analysis techniques. As such, the tool being used is dictated by the question being answered (Benguigui et al., 2015). Defining the specific type of variability measure being applied is critical and the technique applied should be dictated by the question as different techniques analyse different aspects of movement, whether that be stability or variability (Cowin et al., 2022), which despite being related, represent different concepts (Longo et al., 2018). The specific relationship between variability (linear measure) and stability (non-linear measure) is context dependent, as typically an increase in the magnitude of variability represents decreased stability, but instances exist where movements are stable but exhibit variability (Smith et al., 2010).

One technique that analyses stability is the LyE, which is derived from chaos theory, a theory examining dynamic systems, which states that a system is influenced by its initial conditions (Straussfogel & von Schilling, 2009). Specifically, the LyE analyses the local dynamic stability of a system – the degree of sensitivity to small perturbations that exists in a system – by examining how neighbouring trajectories diverge from one another amongst data points at multiple time instances (Buzzi et al., 2003; Chang et al., 2010; Mehdizadeh, 2019; Toebes et al., 2012). The rate of convergence and divergence of trajectories indicate the dynamic stability of a system

(Reynard & Terrier, 2015). A positive LyE value indicates greater variance/divergence in trajectories, a more unstable system and an inability to diminish local perturbations, whereas a negative LyE signifies trajectories converging more over time and a locally stable system (Mehdizadeh, 2018; Stenum et al., 2014; Stergiou et al., 2004). In order to calculate LyE, the state space of the dynamical system must be reconstructed, which requires determining an appropriate embedding dimension, the number of successive points in the dynamical system and time delay, an integer determining how many data points are included for analysis (Matilla-García et al., 2021) Two methods exist to analyse the LyE, the Rosenstein and Wolf methods, each examining movement trajectories but possess differences in their calculation (Wurdeman, 2018a).

Like many non-linear analysis techniques, the LyE has primarily been used to analyse walking gait and the factors that impact it, such as ageing (Terrier & Reynard, 2015), disease/injury (Beurskens et al., 2014; Reynard et al., 2014), and performing a secondary task (Sejdić et al., 2013). This has been prioritised due to the risk associated with falls and the impact a fall has on an individual's economic, physical and psychological health, of which instability during gait is a contributor to this risk (Mehdizadeh, 2018). The prominence of LyE walking gait literature resulted in Mehdizadeh (2018) conducting a systematic review of its application to walking gait. In this review, it was concluded that the different methodology researchers employ result in different LyE exponents being calculated. These methodological differences included experimentation methods like the method of capture (kinematic modelling or accelerometry), body segments/joints assessed, and data analysis methods such as different state space reconstructions (embedding dimension and time delay) or the LyE algorithm applied. Despite the LyE being suited to analyse perturbations during cyclic movements, the application of the LyE to other cyclic movements like cycling and running appears to be much scarcer in comparison to walking gait. However, to properly determine the application of LyE to other forms of movement, a systematic review is required, which to the author's knowledge had previously not been conducted.

As such, this review aims to address the following questions: What cyclic movements, excluding walking, have LyE been applied to? How has LyE been applied to analyse cyclic movement other than walking? Are there differences in the methodologies used?

Summarising the current literature using LyE to analyse human movement (excluding walking gait) will inform researchers of the prevalence of LyE application to other cyclic movements and potentially help guide the future use of LyE.

#### Methods

The structure of the review follows the PRISMA guidelines (Page et al., 2021). The protocol was developed using the Joanna Briggs Institute (JBI) Manual for Evidence Synthesis systematic review development recommendations (Aromataris & Munn, 2020; Peters et al., 2020) and was registered on Open Science Framework on 20 January 2022 to ensure methodology transparency (0.17605/OSF.IO/7BKND). All alterations from the protocol are documented in the relevant sections.

## **Eligibility criteria**

Studies were selected if they adhered to the following criteria.

Inclusion criteria: Full-text peer-reviewed literature/journal articles that are primary original research studies; Englishlanguage papers or papers available in English; sporting movements assessed at the individual level; LyE applied.

Exclusion criteria: Secondary analysis of an intervention study, systematic reviews and meta-analyses; no full-text or English text available; literature that is not peer reviewed including but not limited to grey literature, websites, blogs, conference papers/ abstracts; walking gait research studies; modelling and simulation studies; individual movement not assessed; LyE has not been applied to analyse biomechanical data.

#### Search strategy

A literature search of titles, abstracts and key-words was performed using Medline, EmCare, Embase Scopus and SPORTDiscus, as well as Web of Science databases on 20 February 2022 initially and updated on 1 June 2023 (Figure 1) using the search terms in Table 1.

#### **Data collation/management**

References from each database were exported to Endnote (Endnote 20.1, Clarivate analytics, London, United Kingdom) in separate folders to maintain records of each database. The Endnote library was exported in its entirety to Covidence (Covidence systematic review software, Veritas Health Innovation, Melbourne, Australia) for title and abstract screening. Following database searching (both initial and the re-run), 1672 studies were identified (218 on the re-run). A total of 608 duplicates were removed in Covidence prior to tile and abstract screening, leaving 1213 articles to screen (1096 originally, 117 after the re-run).

#### Screening in covidence

Title and abstract screening was performed by the primary investigator (L.W.), were articles that were clearly not eligible were removed; any that were unclear were included. Following title and abstract screening, as well as reference list searching, 159 articles were eligible for full-text screening. Full-text screening was performed independently by the primary investigator and



Figure 1. Flow chart of the database searching and screening process.

Line	Search terms combined with AND
1	(lyapunov exponent* OR divergence exponent OR dynamic stability OR movement variability).tw,kf.
2	Sports/
3	(sport* OR bicycl* OR swim* OR run* OR row* OR jump* OR basketball* OR throw* OR handball* OR cycling OR danc* OR athlet*).tw,kf.
4	Bicycling/
5	Swimming/
6	Dancing/
7	running/
8	athletic performance/
9	2 OR 3 OR 4 OR 5 OR 6 OR 7 OR 8
10	1 and 9

Table 2. Quality analysis form used in the systematic review.

- Q1 Are there clearly defined research questions?
- Q2 Was a power analysis conducted to determine the sample size required? Optional
- Q3 Are the participant/athlete demographics defined?
- Q4 Is the inclusion/exclusion criteria of the study clearly defined?
- Q5 Were the participants exposed to a familiarisation protocol before data collection? Optional
- Q6 From the information provided on the experimental protocol used, could the research be replicated?
- Q7 Were the methods used for Lyapunov Exponent clearly explained?
- Q8 Was the data used for the Lyapunov Exponent clearly defined?
- Q9 Was test-retest reliability performed or referenced?

Questions were scored as follows: 0 = no description; 1 = limited description; 2 = good description.

a second reviewer (P.T.). Disagreements on the relevance of an article were settled by a third reviewer (R.G.C.). The level of agreement between the researchers was reported using the Kappa statistic to determine inter-rater reliability where 0 is no agreement, 0.01–0.20 is none to slight, 0.21–0.40 is fair agreement, 0.41–0.60 is moderate, 0.6 1–0.80 is substantial and 0.81–0.99 is almost perfect agreement and 1 is perfect agreement. Figure 1 details the combined screening process (both initial and re-run) in flow chart form, utilising the PRISMA guidelines (Page et al., 2021).

#### **Data extraction**

Data was extracted by the primary investigator (LW) Table 3. Similar to Mehdizadeh (2018), the data extracted included the author, study aim/objective, participant number and characteristics (age, height and mass), data capture method, movement, apparatus, sampling rate, segment/joint assessed, data type, surrogation, filtering, embedding dimension (values, calculation and individualisation), time delay (values, calculation and individualisation), number of strides, time-series length, time normalisation, the algorithm applied and LyE values.

#### **Quality assessment**

Critical appraisal was performed to determine the quality of literature that exists in applying the LyE to other forms of movement. This was conducted by both reviewers (L.W. and P.T). The methodological quality assessment tool used was based off a tool implemented by Brown et al. (2014) and tailored to meet the requirements of this review. The quality assessment criteria were 0, 1 or 2, representing clearly did not answer question, maybe answered the question or inadequate information provided and clearly answered the question, respectively. Table 2 lists the questions that were used to analyse the quality of each article.

As such, the highest possible score is either 14, 16 or 18, depending on whether question 2 and/or question 5 is relevant to the study being critically appraised. Quality assessment was expressed as a percentage of the maximum score. A third reviewer (R.G.C.) resolved data extraction disputes if consensus was not reached.

## Results

Overall, 159 studies were identified and 31 satisfied the inclusion/exclusion criteria and were retained for analysis (Figure 1). The inter-rater reliability of the full test screening process was  $\kappa$  = 0.59. Inter-rater reliability was low because initially there was a mis-communication between the reviewers, whereby one reviewer was excluding all gait studies including running.

#### Study characteristics

The included studies contained between 1 and 41 participants with a mean sample size of  $18.1 \pm 8.7$ . The mean participant age was  $25.3 \pm 5.0$  yrs and ranged from 14.0 to 37.1 yrs. The mean participant height was  $1.75 \pm 0.04$  m, while the mean participant body mass was  $69.4 \pm 4.9$  kg. Age, height and mass were not reported in 2 <sup>15, 24</sup>, 5 <sup>11, 15, 24, 27–28</sup> and 6 <sup>10–11, 15, 24, 27–28</sup> studies, respectively. Table 3 reports the study characteristics of studies employing the LyE to analyse movement variability. Studies utilising the LyE have analysed paddling/kayaking twice<sup>10, 20</sup>, jumping once<sup>14</sup>, basketball dribbling once<sup>28</sup>, weighted raise once<sup>27</sup>, skiing once<sup>4</sup>, cycling twice<sup>1, 24</sup> and

LyE.	
the	
investigating	
studies	
of	
. Characteristics	
3.	
Table	

ALG	NS	ROS	NS	ROS	ROS	ROS	ROS	ROS	≥	ROS	ROS	ROS	tinued)
NL	<b>≻</b>	≻	≻	≻	≻	≻	≻	≻	NS	≻	≻	≻	(Con
No. of strides	30	100	50	30	278	120 s	279	100	216	70	150	3 int. of 100	
£	10 MoCap 4 EMG	10	20	26	22–26	19–24	21–27	11 Sternum/ Sacrum 8 Tibia 10 Foot	Reported in supp.	13 arms 16 hands 22 trunk	Foot 5–6; Pelvis 8– 11 Thorax 10–12	$\frac{X = 7.6}{Y = 12.2}$ $Z = 11.8$	
D III N/N	z	z	≻	NS	≻	≻	≻	z	≻	z	≻	z	
calc	AC	AMI	AMI	AMI	AMI	AMI	AMI	AMI	AMI	AMI	AMI	AMI	
ED	4 MoCap 5 EMG	Ŋ	5–6	Ŋ	m	m	ĸ	9 sternum/ sacrum 6 Tibia	7 Ankle 6 Hip and Knee	12 hands 9 upper arms and trunk	6 Foot + Pelvis 6–9 Thorax	6	
ED (VN)	z	z	≻	NS	~	≻	NS	Z	z	z	~	z	
ED calc.	FNN	FNN	FNN	FNN	FNN	FNN	FNN	NN	FNN	FNN	Z N L	FNN	
Filtering (Y/N)	Y – 4 BW 10Hz	z	Y – 4 BW 8Hz	z	Y 4 BW 20 Hz	Y – 4 BW 20Hz	Y - 4 BW - 20Hz	z	Y – 2 BW 15 Hz	NS	NS	NS	
Surrogation (Y/N)	NS	NS	NS	۶	NS	NS	NS	N	NS	NS	SN	NS	
Data Type	Ang. Velo/ Disp + MA	Marker Motion	٩ſ	Ang. Disp.	Vert. Disp.	Vert. Disp.	Vert. Disp	Ang. Velo.	AL	Ang. Velo.	Ang. Velo.	Ang. Velo	
Segment/ Joint	Hip, Knee, Ankle	Trunk	Hip, Knee, Ankle	Leg/Arm	Trunk	Trunk	Trunk	Sternum/ Sacrum/ Tibia/Foot	Ankle/Knee/ Hip	Trunk/ Hands/ Upper arm	Thorax/ Pelvis/ Foot	Shank	
Ч	100 (MoCap)/ 1000 EMG	100	100	100	190	170	190	128	100	100	100	256	
Apparatus	Ergometer	Treadmill	Treadmill	Treadmill	Treadmill	Treadmill	Treadmill	Treadmill/ Overground	Treadmill	Ergometer	Overground	Treadmill	
Movement	Cycling	Running	Running	Skiing	Running	Running	Running	Running	Running	Kayaking	Running	Running	
DCM	MoCap/ EMG	MoCap	MoCap	MoCap	MoCap.	MoCap	MoCap	SI	MoCap	SI	NWI	IMU	
Study	Abbasi, Zamanian and Svoboda (2019)	Arshi, Mehdizadeh and Davids (2015)	Ataabadi et al. (2021)	Cignetti, Schena and Rouard (2009)	Ekizos, Santuz and Arampatzis (2017)	Ekizos, Santuz and Arampatzis (2018)	Ekizos et al. (2018)	Fohrmann et al. (2022)	Frank, Prentice and Callaghan (2019)	Hamacher et al. (2018)	Hoenig et al. (20192019)	Hollander, Hamacher and Zech (2021)	
Ref. No.	-	7	m	4	Ŋ	9	7	ω	6	10	1	12	

ALG	NS	ROS	¥	ROS	ROS	ROS	ROS	ROS	ROS	NS	<b>&gt;</b> >	ROS + W	×	NS	NS	ROS	tinued)
NL	NS	≻	z	≻	≻	≻	≻	≻	NS	NS	NS NS	≻	≻	NS	NS	≻	(Con
No. of strides	5 min	30	8-10	150	100	100	100	50 (approx.)	30	100	NS NS	80	3 min	10	42	287 (max)	
£	0.192 s	12.7	52-56	9–18 sagittal 13–25 frontal 13–24 horizontal	10	10	10	20	18–24	NS	10 21–33	22 A-P 15 Vert 29 M-L	NS	NS	NS	36-40	
TD (V/N)	NS	≻	NS	>	z	z	z	z	≻	NS	NS N/A	~	≻	NS	NS	≻	
TD calc	AMI	AC	AMI	AMI	AMI	AMI	AMI	AMI	AMI	AC	AMI	AMI	AMI	NS	NS	AMI	
E	ъ	5.12	ŝ	8 sagittal 7 frontal + horizontal	4	5	5	4	4	NS	4 4	J.	NS	NS	NS	NS	
ED (V/N)	NS	≻	NS	z	z	z	z	NS	z	NS	NS N/A	≻	≻	z	NS	NS	
ED calc.	FNN	FNN	FNN	Z Z Z	FNN	FNN	FNN	FNN	FNN	Delay embedding vectors	FNN FNN	NN	FNN	NS	NS	FNN	
Filtering (Y/N)	Y – 2 BW 6Hz	Y – BW 6 Hz	NS	z	Z	Z	Z	NS	Y – BW 10 Hz	NS	NS NS	NS	z	NS	NS	Y – 4 BW 50Hz	
urrogation (Y/N)	NS	NS	NS	SN	NS	NS	NS	NS	NS	NS	NS Y	NS	Ns	NS	NS	NS	
S Data Type	Vert. Disp.	AL	JA/COM pos.	Ang. Velo.	Marker Motion	Marker Motion	X, Y, Marker Velocity	Vertical traiectorv	AL	Stride frequency	Posture Vector JA	Marker pos.	JA/Sacrum Pos.	EEG	Marker Coordinates	Vertical Coordinates	
Segment/ Joint	Head, Ankle	Ankle, Knee, Hin	Hip, Knee, Sacrum	Trunk	ankle, knee, hip, sacrum	Trunk	Ankle	Wrist	Ankle, Knee, Hin, Trunk	N/A	Whole Body Knee	Sacrum	Hip, Knee, Pelvis	Brain	Wrist, Elbow, Shoulder	Trunk	
뷮	125	240	300	100	200	100	100	120	100	1000	250 100	120	120	1000	120	100	
Apparatus	Treadmill	Overground	Treadmill	Treadmill	Treadmill	Treadmill	Treadmill	Ergometer	Treadmill	Treadmill	Treadmill Ergometer	Treadmill	Treadmill	Dumbbell	N/A	Treadmill	
Movement	Running	Jumping	Running	Running	Running	Running	Running	Paddling	Running	Running	Running Cycling	Running	Running	Weighted Raise	Dribbling	Running	
DCM	MoCap	MoCap	MoCap	NM	MoCap	MoCap	MoCap	MoCap	MoCap	MoCap	MoCap MoCap	MoCap	MoCap	EEG	MoCap	MoCap	
Study	Jordan et al.	Larson et al.	Look et al. (2013)	Mahaki et al. (2020)	Mehdizadeh, Arshi and Davids (2014)	Mehdizadeh, Arshi and Davids (2014)	Mehdizadeh, Arshi and Davids (2016)	Nessler et al. (2015)	Ogaya et al.	Padulo et al. (2023)	Promsri (2022) Quintana-Duque and Saupe (2013)	Raffalt et al., (2019)	Raffalt et al., (2020)	Rahatabad et al.	Robalo et al. (2021)	Santuz et al. (2018)	
Ref. No.	13	14	15	16	17	18	19	20	21	22	23 24	25	26	27	28	29	

		I ALG	×	ROS	200
		Ŧ	Ň	≻	4   4   4
	No. of	strides	2 min	120	
		D	18	7 A-P 10 M-L	9 Vert
β	ind.	(N/λ)	z	z	
	β	calc	AMI	AMI	TN     
		ED	Ŋ	9	icited
⊞	Ind.	(N/N)	z	z	
		ED calc.	FNN	FNN	unclop comi
		Filtering (Y/N)	z	Y 2 BW 20/2 Hz (IMU), 15Hz	(EMG)
	Surrogation	(N/N)	NS	NS	- and dish
		Data Type	٩ſ	ACC/EMG	
	Segment/	Joint	Hip, Knee	Trunk/LB muscles	
		Ηz	250	100	>.\o+c
		Apparatus	Treadmill	Treadmill	Lotter (complete
		Movement	Running	Running	- H - H - H - H - H - H - H - H - H - H
		DCM	MoCap	IMU	
		Study	Strongman and Morrison	Walsh (2021)*	former: DCM - dat
	Ref.	No.	30	31	DEC

Table 3. (Continued).

T Velo = angular velocity; Ang. Disp = angular displacement; MA = muscle activity; BW = Butterworth filter; FNN = global false nearest neighbours method; AC = auto-correlation function; EMG = electromyography; AMI = average mutual information function; ROS = Rosenstein algorithm; Vert. disp. = vertical displacement; s = seconds; IS = inertial sensors; JA = joint angle; W = Wolf; IMU = inertial measurement unit; COM = Centre of mass; pos. = position; K = Kantz; min = minutes; ACC = acceleration; A-P = anterior-posterior, M-L = medio-lateral, vert = vertical; Int. = Interval; LB = lower body. predominately running, which occurred applied 23 times<sup>2-3,5-</sup> 9,11-13, 14-19, 21-23, 25-26, 29-31

## Methodological quality assessment

The methodological quality assessment is reported in Appendix 1. The mean quality of the 31 studies was 67.6%. The mean rating for sample size power analysis was 0.21 as it was conducted in three studies <sup>3,23, 28</sup> (10%) to justify their sample size. The mean rating for the description of participant demographics and explanation of the inclusion/exclusion criteria was 1.65 and 1.73, respectively. Whether participants were exposed to a familiarisation protocol had a mean rating of 0.73. Test replicability had a mean rating of 1.81. Whether the methods used to calculate the LyE were clearly explained and if the data used to calculate the LyE were clearly defined had a mean rating of 1.81 and 1.87, respectively. Test–retest reliability was performed or referenced in 10<sup>3,4,6,8,12,16,21,22,24,29</sup> studies (32%), possessing a mean rating of 0.48.

#### LyE protocol

Table 3 also reports the LyE methodologies that each study employed. LyE has been applied to inertial measurement units/ sensors (IMUs) six times,<sup>8,10–12,16,31</sup>, electroencephalography (EEG) once<sup>27</sup> and electromyography (EMG), twice<sup>1,31</sup>. Predominately, data was captured via motion capture (segment and joint angular kinematics) and was done so 24 times <sup>1–7, 9, 13–15, 17–26, 28–30</sup>.

Sampling rate ranged in studies using motion capture from 100 to 1000 Hz, with 100 Hz being the most commonly used sampling rate, used 10 times<sup>1-4, 9, 18–20, 21, 24, 29</sup>. Similarly, the IMU sampling rate ranged between 100 and 256 Hz, whereby 100 Hz was the most commonly used sampling rate, used four times<sup>10–11,16,31</sup>. The sampling rate in the EEG<sup>27</sup> and EMG<sup>1,31</sup> (in both instances) studies was 1000 Hz.

Surrogation was only performed in two studies<sup>4, 24</sup>. Filtering occurred in  $11^{1,3,5-7,9,13,14,21,29,31}$  studies where a Butterworth filter was applied in all instances. Cut-off-frequencies ranged between 2.5 and 50 Hz, with 20 Hz being the most common (4 times). A 2<sup>nd</sup> and 4<sup>th</sup> order filter was applied 3<sup>9, 13,31</sup> and 6<sup>1,3,5-7,29</sup> times, respectively, and two studies did not report this information<sup>14,21</sup>.

A range of segments/joints were analysed. In paddling, the upper extremity was analysed in both studies<sup>10,20</sup> and one also analysed the upper body (trunk)<sup>10</sup>. The repeated jumping study<sup>14</sup> assessed the lower body (ankle, knee and hip). In the two cycling studies<sup>1, 24</sup>, the lower body was assessed. The upper extremity was analysed in the basketball dribbling study<sup>28</sup>. The upper extremity and lower body were assessed in the skiing study<sup>4</sup>. In running, the lower body was assessed in 12<sup>3, 8,9,11–14,17,19,21,26,30</sup> instances, the upper body in 11<sup>2, 5–8, 11, 16,18,21,29,31</sup>, the pelvis in 6<sup>8, 11,14,17,25–26</sup>, and the head<sup>13</sup> and whole body<sup>23</sup> in one instance each. Stride frequency (which is not associated with a joint) was assessed in one case<sup>22</sup>. The brain was assessed in the weighted raise (EEG) study<sup>27</sup>.

The embedding dimension must be defined before the LyE is calculated. The global false nearest neighbours were the most

commonly applied algorithm, applied in 28 cases<sup>1–21,23–26,29–31</sup>. Delay embedding vectors were implemented once<sup>22</sup> and how the embedding dimension was constructed was not reported in two cases<sup>27, 28</sup>. The embedding dimension was individualised in 7 instances<sup>3,5–6,11,14,25–26</sup>, fixed (i.e., constant between participants) in 14<sup>1–2,8–1012,16–19,21,27,30–31</sup> cases and whether individualisation occurred was not stated in 9 instances<sup>4,7,13,15,20,22–23,28–29</sup>. In the case study, whether individualisation between participants was not applicable<sup>27</sup>. Embedding dimension values ranged between 3 and 12, with five being the most common value, similar to walking gait (Mehdizadeh, 2018) used in eight instances<sup>2–4,13,18–19,25,30</sup>. Embedding dimension values were not reported in five studies<sup>22, 26–29</sup>.

Time delay like the embedding dimension must be defined before the LyE is calculated (Matilla-García et al., 2021). Within the non-walking gait movement literature, the average mutual information function is the most commonly applied algorithm to calculate it, used in 25 instances<sup>2-13,15-21,23,25-26,29-31</sup>. The autocorrelation function was applied in four instances<sup>1,14,22,24</sup>, and how the time delay was calculated was not reported in two instances<sup>27-28</sup>. Time delay was individualised in 12 instances, 3,5-7,9,11,14,16,21,25-26,29 but it was either not reported whether it was individualised or not individualised in  $7^{4,13,15,22-}$ <sup>23,27-28</sup> and  $11^{1-2,8,10,12,17-20,30-31}$  studies, respectively. Time delay individualisation was not relevant in the case study<sup>27</sup>. Time delay was reported in two ways, as either a fixed number (26 instances<sup>1-12,14-21,23-25,29-31</sup>) or in seconds (2 instances)<sup>4,13</sup>. One study reported both seconds and a fixed number<sup>4</sup>. Time delays ranged between 5 and 56, with the most common occurrence being 10 (7 instances)<sup>1-2,8,17-19,23,31</sup>, similar to the walking gait as reported in (Mehdizadeh, 2018). Time delay values were not reported in four studies<sup>22, 26–28</sup>.

The number of strides/cycles used varied, ranging between 8<sup>15</sup> and 287<sup>29</sup> steps in running, 50<sup>20</sup> and 70<sup>10</sup> in kayaking, 42 in basketball dribbling<sup>28</sup> and 30 cycles in cycling<sup>1</sup> and skiing<sup>4</sup> and 10 in the weighted raise<sup>27</sup>. The number of cycles/strides was not reported in two studies<sup>23–24</sup> or written as an interval of time in four studies<sup>6,13,26,30</sup>. Time-series length was reported in 16 studies<sup>1–8,10–12,16,18–19,29,31</sup> and not reported in 15 instances<sup>9</sup>, <sup>13–15,17,20–28,30</sup>, Data was time normalised in 21 instances<sup>1–8,10–12,1416–19,25–26,29,31</sup>, was not time normalised in one instance<sup>15</sup>, and it was not stated whether time normalisation occurred in 9 instances<sup>9,13,21–24,27–28,30</sup>.

The Rosenstein algorithm was the most widely adopted algorithm, applied in 19 studies<sup>2,4–8,10–12,14,16–21,25,29,31</sup>. Wolf's and Kantz algorithm was applied in  $6^{9,23–26,30}$  and  $1^{15}$  study, respectively. However, three papers did not specifically mention what algorithm they applied but referenced Rosenstein and Kantz in  $2^{3, 13}$  and  $1^{22}$  study, respectively. The algorithm used was not specified in three studies<sup>1,27–28</sup>.

Due to the large range of LyE values that have been produced in the studies included in this review, even when analysing the same task, LyE values were not reported in text, and are instead reported in Appendix 2. This was particularly evident in the 23 studies that analysed running. As no study replication has occurred, it renders comparison of LyE values implausible.

## Discussion

The aim of this review was to determine the breadth of current research applying the LyE to movements other than walking gait and determine if methodological differences exist between studies. Within the 31 studies that were included for the analysis, it was confirmed that methodological differences do exist within the literature applying the LyE culminating in a range of LyE values being produced. Additionally, running is the most commonly assessed non-walking gait movement when applying the LyE, utilised in 23 of the 31 included studies. Furthermore, research has commonly taken place using motion capture systems.

#### LyE protocol

#### Environment

Due to the large number of data points required when calculating the LyE, research analysing other activities has predominately been conducted on treadmills and/or ergometers. Similar to walking gait, it has been found that treadmill running produces a more stable running pattern than overground running (Fohrmann et al., 2022), which is likely due to the constraints of the environment. It has been demonstrated in various activities that removing a task from its original domain can alter movement expression (Pinder et al., 2011). For example, in cycling (Wilkinson & Lichtwark, 2021), concluded that ergometer alter the lateral dynamics of a bicycle, as they do not permit side-to-side movement. As such, future research should be conducted in ecologically valid domains to best capture the movement pattern that occurs during performance in realistic situations (Cowin et al., 2022). To do this, data must be captured with measurement tools that allow for greater ecological validity. One such tool is IMUs, portable devices that consist of a triaxial accelerometer, gyroscope and magnetometer (Schall et al., 2016). Despite their improved ecological validity and reduced cost, making them a more plausible and practical option, research has predominantly been done via motion capture. Twenty-four studies collected data via motion capture compared to six that captured data via IMUs in this review. Therefore, further research analysing the LyE should be applied using IMUs to improve the ecological validity of research applying the LyE to human movement.

## Filtering

The purpose of filtering data is to remove unwanted noise and leave the relevant information (de Cheveigné & Nelken, 2019). However, within LyE research, it is recommended that filtering does not occur (or occurs with a high cut-off frequency) as it can affect the LyE calculation by potentially removing "true" fluctuations that occur, thus changing the dynamics of the system (Raffalt et al., 2020). Despite this, filtering occurred in 11 of the studies analysed, with varying cut-off frequencies, which is not best practice. Future LyE studies should avoid filtering data prior to calculating the LyE due to its potential effect on the results.

#### Sampling rate

An appropriate sampling rate is important as too small a sampling rate will result in information not being captured and too high a sampling rate will result in too much information being captured (Fallahtafti et al., 2021). Whilst studies have analysed the effect of data length (Hussain et al., 2020) and time normalisation (Raffalt et al., 2019) on the calculation of the LyE, to the author's knowledge, no study has analysed the effect of sampling rate on LyE calculation on kinematic data. This could be done in future investigations to ensure more similar methodologies occur between research applying the LyE.

### Surrogation

Surrogation is a technique applied in non-linear analysis to determine if "true" non-linearity exists within a dynamic system (Faes et al., 2009). Specifically, it involves generating a null hypothesis, which assumes the system is linear, after which, the original data is transformed and a random data set is generated that retains some of the properties of the original time series (Lancaster et al., 2018). Within the LyE analysis, if the LyE of the original data differs significantly from the surrogate data, then it can be inferred that the original data is chaotic/deterministic (Stergiou et al., 2004). However, only two studies had performed a surrogate analysis to determine the non-linearity of the system; most researchers have assumed that non-linearity exists within their data set, without first establishing it. As such, future research applying the LyE to movement data (or any data assessing a dynamical system) should first run a surrogation analysis to determine if the data set has a chaotic origin. This is important because data that is completely random will produce a positive LyE just as chaotic data produce a positive LyE. Without a surrogate analysis, the difference between the two cannot be detected (Stergiou et al., 2004).

#### State space reconstruction

To calculate the LyE, the state space (the vector area where the dynamic system is defined) must be defined and involves determining the embedding dimension and time delay. There are several approaches to determine the embedding dimension and time delay, the most commonly used methods are the global false nearest neighbour and average mutual information function, respectively, consistent with walking gait literature (Mehdizadeh, 2018; Wurdeman, 2018b). However, different methods have also been reported to reconstruct the state space. Studies with different methodologies should not be compared as different state space reconstructions will affect the LyE calculation (Mehdizadeh, 2018; van Schooten et al., 2013). It is thought that each individual is a unique dynamic system due to possessing their own individual constraints (Davids et al., 2006). As such, it can be inferred that individual state space reconstruction should occur. Additionally, Raffalt et al. (2019) determined that the effectiveness of LyE calculation, regardless of which algorithm was applied, improved when individualising the time delay and embedding dimension. Individualisation of the time delay and

embedding dimension occurred in 12 and 7 occasions, respectively. Individualising the state space should occur in future research. More alarmingly, both how it was calculated, and values of the embedding dimension and time delay were not reported in three and four occasions for the embedding dimension and two and five occasions for the time delay, respectively. Failing to report this makes study replication difficult, particularly as these values will impact the LyE calculation because state space reconstruction is a precursor to determining the LyE (Amirpourabasi et al., 2020; Wurdeman, 2018b).

### **Time series length**

The number of strides/cycles influences LyE calculation and as such should be reported (Bruijn et al. 2009). This did not occur in two studies, making repeatability of the study implausible. Furthermore, due to the impact of the number of cycles/strides on the calculation of the LyE, studies should implement a fixed number of strides/cycles to compare the LyE between different conditions (Mehdizadeh, 2018). The number of strides analysed will affect time-series length which impacts the calculation of the LyE (Mehdizadeh, 2018). Similarly, the LyE calculation is also affected by whether time normalisation occurs prior to it, hence it should be reported if it took place (Raffalt et al., 2019; Stenum et al., 2014). However, in 10 instances, studies either did not time normalise their data or state whether time normalisation occurred.

### Algorithm

Consistent with the walking gait literature (Mehdizadeh, 2018), the most commonly applied algorithm is the Rosenstein algorithm. However, six studies either did not report or did not explicitly state the algorithm they used, which is problematic. Both the Wolf and Rosenstein algorithms calculate the LyE in different ways, which leads to different results (Raffalt et al., 2019). For example, when using the same lower limb data set, Cignetti et al. (2012)demonstrated that that different results were produced from the Rosenstein and Wolf algorithms, where the Rosenstein algorithm underestimated the LyE, and the Wolf algorithm overestimated it. As such, due to the different results they may produce, it is best practice to report the algorithm applied for transparency.

A broad range of LyE values exist, even for the same activity. Similar to Mehdizadeh (2018), the broad range of LyE reported is due to the broad range of methods being employed to calculate the LyE. However, another cause is the different research questions that have been answered. Because of this, it is difficult to generate normative values and determine what constitutes an acceptable LyE value for a given activity. As such, it is important that comparisons of the LyE only be made to other studies that are attempting to answer a similar question and employ the same or very similar methods (i.e., similar participant characteristics, state space reconstruction and capture method).

#### **Quality assessment**

The mean quality of the 31 studies was 67.6%. This quality score was generated through items 1 (clear research guestions), and 6-8 (was the experimental protocol clearly detailed?, where the methods used for the LyE clearly detailed? and was the data used to calculate the LyE clearly defined?) which suggest that the majority of the studies are repeatable. However, a power analysis was performed in 3 of 28 eligible studies (11%) of studies. Within biomechanics, an insufficient sample size is an issue that continues to persist (Oliveira & Pirscoveanu, 2021), likely stemming through difficulties with recruitment. However, power analyses are required to determine the required sample size needed to see whether the results obtained are significant or not (Kemal, 2020). As such, this should be addressed in future research. Test-retest reliability determines the repeatability of a measurement and involves repeating the same measurement more than once (Hopkins, 2000). As such, it determines the consistency of the measurement. However, only 10 studies (32%) referenced reliability, resulting in a mean score of 0.48. However, only one study performed test re-test reliability which future studies should address this to better determine if the observed variability changes as calculated by the LyE are "real" or not.

## Conclusion

Similar to walking gait (Mehdizadeh, 2018), discrepancy in calculating the LyE to analyse dynamic stability exists in experimental design (i.e., the question the researchers are trying to answer), data pre-processing (i.e., filtering and sampling rate) and the LyE calculation method (i.e., algorithm and state space reconstruction). This renders comparison of LyE values (even when comparing the same task) implausible. Additionally, there exist limitations in the current research applying LyE, namely the lack of research conducting a surrogation analysis prior to LyE calculation to determine whether the data does have a chaotic origin, and the lack of ecological validity in the current research (which could be addressed better through the use of devices like IMUs). Furthermore, there is a scarcity of test-retest reliability analysis that has been conducted which is necessary to determine if changes in the LyE are "real" or not. Addressing these limitations and others that have been presented will improve the application of the LyE.

#### **Disclosure statement**

No potential conflict of interest was reported by the author(s).

## Funding

The author(s) reported that there is no funding associated with the work featured in this article.

## ORCID

Lachlan Winter (D) http://orcid.org/0000-0003-0136-4546

#### References

- Abbasi, A., Zamanian, M., & Svoboda, Z. (2019). Nonlinear approach to study the acute effects of static and dynamic stretching on local dynamic stability in lower extremity joint kinematics and muscular activity during pedalling. *Human Movement Science*, 66, 440–448. https://doi.org/10. 1016/j.humov.2019.05.025
- Aghaie Ataabadi, P., Sarvestan, J., Alaei, F., Yazdanbakhsh, F., & Abbasi, A. (2021). Linear and non-linear analysis of lower limb joints angle variability during running at different speeds. *Acta Gymnica*, *51*. https://doi. org/10.5507/ag.2021.023
- Amirpourabasi, A., Chow, J. Y., Vicinanza, D., & Williams, G. K. R. (2020). Influence of calculation parameters on nonlinear dynamics measures. 38th International Society of Biomechanics in Sport Conference, International Society of Biomechanics. https://commons.nmu.edu/cgi/ viewcontent.cgi?article=2071&context=isbs
- Aromataris, E., & Munn, Z. E. (2020), July 16. JBI manual for evidence synthesis. JBI. https://synthesismanual.jbi.global/
- Arshi, A Reza., Mehdizadeh, S, and Davids, K. (2015). Quantifying foot placement variability and dynamic stability of movement to assess control mechanisms during forward and lateral running. *Journal of Biomechanics*, 48(15), 4020–4025. https://doi.org/10.1016/j.jbiomech. 2015.09.046
- Benguigui, N., Komar, J., Seifert, L., Thouvarecq, R., & Benguigui, N. (2015). What variability tells us about motor expertise: Measurements and perspectives from a complex system approach. *Movement & Sport Sciences - Science & Motricité*, 89(89), 65–77. https://doi.org/10.1051/ sm/2015020
- Bernstein, N. A. (1967). The Co-ordination and regulation of movements. Pergamon Press.
- Beurskens, R., Wilken, J. M., & Dingwell, J. B. (2014). Dynamic stability of superior vs. inferior body segments in individuals with transtibial amputation walking in destabilizing environments. *Journal of Biomechanics*, 47 (12), 3072–79. https://doi.org/10.1016/j.jbiomech.2014.06.041
- Brown, S. R., Brughelli, M., & Hume, P. A. (2014). Knee mechanics during planned and unplanned sidestepping: A systematic review and meta-analysis. *Sports Medicine*, 44(11), 1573–88. https://doi.org/10. 1007/s40279-014-0225-3
- Bruijn, S. M., van Dieën, J. H., Meijer, O. G., & Beek, P. J. (2009). Statistical precision and sensitivity of measures of dynamic gait stability. *Journal of Neuroscience Methods*, *178*(2), 327–333. https://doi.org/10.1016/j.jneu meth.2008.12.015
- Buzzi, U. H., Stergiou, N., Kurz, M. J., Hageman, P. A., & Heidel, J. (2003). Nonlinear dynamics indicates aging affects variability during gait. *Clinical Biomechanics*, 18(5), 435–43. https://doi.org/10.1016/S0268-0033(03)00029-9
- Caballero, C., Barbado, D., & Moreno, F. J. (2014). Non-linear tools and methodological concerns measuring human movement variability: An overview. *European Journal of Human Movement*, 32(1), 61–81.
- Chang, M. D., Sejdić, E., Wright, V., & Chau, T. (2010). Measures of dynamic stability: Detecting differences between walking overground and on a compliant surface. *Human Movement Science*, 29(6), 977–86. https:// doi.org/10.1016/j.humov.2010.04.009
- Cignetti, F., Decker, L. M., & Stergiou, N. (2012). Sensitivity of the Wolf's and Rosenstein's algorithms to evaluate local dynamic stability from small gait data sets. *Annals of Biomedical Engineering*, *40*(5), 1122–30. https:// doi.org/10.1007/s10439-011-0474-3
- Cignetti, F., Schena, F., and Rouard, A. (2009). Effects of fatigue on intercycle variability in cross-country skiing. *Journal of Biomechanics*, 42(10), 1452–1459. https://doi.org/10.1016/j.jbiomech.2009.04.012
- Cowin, J., Nimphius, S., Fell, J., Culhane, P., & Schmidt, M. (2022). A proposed framework to describe movement variability within sporting tasks: A scoping review. *Sports Medicine - Open*, 8(1), 85–109. https://doi.org/ 10.1186/s40798-022-00473-4
- Davids, K., Bennett, S., & Newell, K. M. (2006). Movement system variability. Human Kinetics.

- de Cheveigné, A., & Nelken, I. (2019). Filters: When, why, and how (not) to use them. *Neuron*, 102(2), 280–93. https://doi.org/10.1016/j.neuron.2019. 02.039
- Dingwell, J. B., & Cusumano, J. P. (2000). Nonlinear time series analysis of normal and pathological human walking. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 10(4), 848–63. https://doi.org/10.1063/1. 1324008
- Ekizos, A., Santuz, A., & Arampatzis, A. (2017). Transition from shod to barefoot alters dynamic stability during running. *Gait & Posture*, *56*, 31– 36. https://doi.org/10.1016/j.gaitpost.2017.04.035
- Ekizos, A., Santuz, A., & Arampatzis, A. (2018). Short- and long-term effects of altered point of ground reaction force application on human running energetics. *Journal of Experimental Biology*, https://doi.org/10.1242/jeb. 176719
- Ekizos A., Santuz A., Schroll A., & Arampatzis A. (2018). The Maximum Lyapunov Exponent During Walking and Running: Reliability Assessment of Different Marker-SetsTable\_1.docx. Frontiers in Physiology, 9. https://doi.org/10.3389/fphys.2018.0110110.3389/fphys. 2018.01101.s001
- Estep, A., Morrison, S., Caswell, S., Ambegaonkar, J., & Cortes, N. (2017). Differences in pattern of variability for lower extremity kinematics between walking and running. *Gait & Posture*, 60, 111–115. https://doi. org/10.1016/j.gaitpost.2017.11.018
- Faes, L., Zhao, H., Chon, K. H., & Nollo, G. (2009). Time-varying surrogate data to assess nonlinearity in nonstationary time series: Application to heart rate variability. *IEEE Transactions to Biomedical Engineering*, 56(3), 685–695. https://doi.org/10.1109/TBME.2008.2009358
- Fallahtafti, F., Wurdeman, S. R., & Yentes, J. M. (2021). Sampling rate influences the regularity analysis of temporal domain measures of walking more than spatial domain measures. *Gait & Posture*, 88, 216–20. https://doi.org/10.1016/j.gaitpost.2021.05.031
- Fohrmann, D., Hamacher, D., Sanchez-Alvarado, A., Potthast, W., Mai, P., Willwacher, S., & Hollander, K. (2022). Reliability of running stability during treadmill and overground running. *Sensors*, 23(1), 347–58. https://doi.org/10.3390/s23010347
- Frank, N. S., Prentice, S. D., & Callaghan, J. P. (2019). Local dynamic stability of the lower extremity in novice and trained runners while running intraditional and minimal footwear. *Gait & Posture*, 68, 50–54. https:// doi.org/10.1016/j.gaitpost.2018.10.034
- Glanzer, J. A., Diffendaffer, A. Z., Slowik, J. S., Drogosz, M., Lo, N. J., & Fleisig, G. S. (2021). The relationship between variability in baseball pitching kinematics and consistency in pitch location. *Sports Biomechanics*, *20*(7), 879–86. https://doi.org/10.1080/14763141.2019. 1642378
- Gray, R. (2020). Changes in movement coordination associated with skill acquisition in baseball batting: Freezing/freeing degrees of freedom and functional variability. *Frontiers in Psychology*, *11*(1295), 1–18. https://doi. org/10.3389/fpsyg.2020.01295
- Hamacher, D., Krebs, T., Meyer, G., & Zech, A. (2018). Does local dynamic stability of kayak paddling technique affect the sports performance? A pilot study. *European Journal of Sport Science*, *18*(4), 491–496. https://doi. org/10.1080/17461391.2018.1435726
- Hamill, J., van Emmerik, R. E. A., Heiderscheit, B. C., & Li, L. (1999). A dynamical systems approach to lower extremity running injuries. *Clinical Biomechanics*, 14(5), 297–308. https://doi.org/10.1016/S0268-0033(98)90092-4
- Hoenig T., Hamacher D., Braumann K., Zech A., & Hollander K. (2019). Analysis of running stability during 5000 m running. *European Journal* of Sport Science, 19(4), 413–421. https://doi.org/10.1080/17461391.2018. 1519040
- Hollander, K., Hamacher, D., & Zech, A. (2021). Running barefoot leads to lower running stability compared to shod running - results from a randomized controlled study. *Scientific reports*, 11(1). https://doi.org/10. 1038/s41598-021-83056-9
- Hopkins, W. G. (2000). Measures of reliability in sports medicine and science. Sports Medicine, 30(1), 1–15. https://doi.org/10.2165/00007256-200030010-00001

- Hussain, V. S., Spano, M. L., & Lockhart, T. E. (2020). Effect of data length on time delay and embedding dimension for calculating the Lyapunov exponent in walking: Effect of data length on time delay and embedding dimension for calculating the Lyapunov exponent in walking. *Journal of the Royal Society Interface*, 17(168), 20200311–19. https://doi.org/10. 1098/rsif.2020.0311
- James, C. R. (2004). Considerations of movement variability in biomechanics research. In N. Stergiou (Ed.), *Innovative analyses of human movement* (Vol. 1, pp. 29–62). Human Kinetics.
- Jordan, K., Challis, J. H., Cusumano, J. P., & Newell, K. M. (2009). Stability and the time-dependent structure of gait variability in walking and running. *Human Movement Science*, 28(1), 113–128. https://doi.org/10.1016/j. humov.2008.09.001
- Kemal, O. (2020). Power analysis and sample size, when and why? *Turkish Archives of Otorhinolaryngology*, *58*(1), 3–4. https://doi.org/10.5152/tao. 2020.0330
- Lancaster, G., latsenko, D., Pidde, A., Ticcinelli, V., & Stefanovska, A. (2018). Surrogate data for hypothesis testing of physical systems. *Physics Reports*, 748, 1–60. https://doi.org/10.1016/j.physrep.2018.06.001
- Larson, J., Perkins, E., Oldfather, T., Zabala, M., & Sakakibara, M. (2021). Local dynamic stability of the lower-limb as a means of post-hoc injury classification. *PLoS ONE*, *16*(6), e0252839. https://doi.org/10.1371/journal. pone.025283910.1371/journal.pone.0252839.g00310.1371/journal. pone.0252839.g00210.1371/journal.pone.0252839.g00310.1371/journal.pone.0252839.g00510.371/journal.pone.0252839.g0051000000000
- Longo, A., Federolf, P., Haid, T., & Meulenbroek, R. (2018). Effects of a cognitive dual task on variability and local dynamic stability in sustained repetitive arm movements using principal component analysis: A pilot study. *Experimental Brain Research*, 236(6), 1611–19. https://doi. org/10.1007/s00221-018-5241-3
- Look N., Arellano, C. J., Grabowski, A. M., McDermott, W. J., Kram R., & Bradley, E. (2013). Dynamic stability of running: The effects of speed and leg amputations on the maximal Lyapunov exponent. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, *23*(4). https://doi.org/10. 1063/1.4837095
- Magill, R., & Anderson, D. (2017). *Motor learning and control: Concepts and applications* (11th ed.). McGraw Hill Education.
- Mahaki, M., Mimar, R., Sadeghi, H., Khaleghi Tazji, M., & Vieira, M. F. (2020). The effects of general fatigue induced by incremental exercise test and active recovery modes on energy cost, gait variability and stability in male soccer players. *Journal of Biomechanics*, *106*, 109823. https://doi. org/10.1016/j.jbiomech.2020.109823
- Matilla-García, M., Morales, I., Rodríguez, J. M., & Marín, M. R. (2021). Selection of embedding dimension and delay time in phase space reconstruction via symbolic dynamics. *Entropy*, 23(2), 1–13. https://doi. org/10.3390/e23020221
- Mehdizadeh, S. (2018). The largest lyapunov exponent of gait in young and elderly individuals: A systematic review. *Gait & Posture, 60,* 241–50. https://doi.org/10.1016/j.gaitpost.2017.12.016
- Mehdizadeh, S. (2019). A robust method to estimate the largest lyapunov exponent of noisy signals: A revision to the Rosenstein's algorithm. *Journal of Biomechanics*, 85, 84–91. https://doi.org/10.1016/j.jbiomech. 2019.01.013
- Mehdizadeh, S., Arshi, A. R., & Davids, K. (2014). Effect of speed on local dynamic stability of locomotion under different task constraints in running. *European Journal of Sport Science*, 14(8) 791–798. https://doi.org/ 10.1080/17461391.2014.905986
- Mehdizadeh, S., Arshi, A. R., & Davids, K. (2014). Quantification of stability in an agility drill using linear and nonlinear measures of variability. Acta of Bioengineering and biomechanics, 16(3), 59–67.
- Mehdizadeh, S., Arshi, A. R., & Davids, K. (2016). Constraints on dynamic stability during forward, backward and lateral locomotion in skilled football players. *European Journal of Sport Science*, 16(2), 190–198. https://doi.org/10.1080/17461391.2014.995233
- Nessler, J. A., Silvas, M., Carpenter, S., Newcomer, S. C., & Gao, Z. (2015). Wearing a Wetsuit Alters Upper Extremity Motion during Simulated Surfboard Paddling. *PLoS ONE*, *10*(11), e0142325. https://doi.org/10. 1371/journal.pone.014232510.1371/journal.pone.0142325.g00110. 1371/journal.pone.0142325.g00210.1371/journal.pone.0142325.

g00310.1371/journal.pone.0142325.g00410.1371/journal.pone. 0142325.g00510.1371/journal.pone.0142325.t00110.1371/journal.pone. 0142325.t00210.1371/journal.pone.0142325.t00310.1371/journal.pone. 0142325.t004

- Ogaya, S., Suzuki, M., Yoshioka, C., Nakamura, Y., Kita, S., & Watanabe, K. (2021). The effects of trunk endurance training on running kinematics and its variability in novice female runners. *Sports Biomechanics*, 1–12. https://doi.org/10.1080/14763141.2021.1906938
- Oliveira, A. S., & Pirscoveanu, C. I. (2021). Implications of sample size and acquired number of steps to investigate running biomechanics. *Scientific Reports*, *11*(1), 3083–98. https://doi.org/10.1038/s41598-021-82876-z
- Padulo, J., Ayalon, M., Barbieri, F. A., Di Capua, R., Doria, C., Ardigò, L. P., & Dello Iacono, A. (2023). Effects of Gradient and Speed on Uphill Running Gait Variability. Sports Health: A Multidisciplinary Approach, 15(1) 67–73. https://doi.org/10.1177/19417381211067721
- Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D., Shamseer, L., Tetzlaff, J. M., Akl, E. A., Brennan, S. E., Chou, R., Glanville, J., Grimshaw, J. M., Hróbjartsson, A., Lalu, M. M., Li, T., Loder, E. W., Mayo-Wilson, E., & Moher, D. (2021). The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *PloS Medicine*, *18*(3), 1003583–98. https://doi.org/10.1371/journal.pmed.1003583
- Peters, M. D. J., Marnie, C., Tricco, A. C., Pollock, D., Munn, Z., Alexander, L., McInerney, P., Godfrey, C. M., & Khalil, H. (2020). Updated methodological guidance for the conduct of scoping reviews. *JBI Evidence Synthesis*, *18*(10), 2119–26. https://doi.org/10.11124/JBIES-20-00167
- Pinder, R. A., Davids, K., Renshaw, I., & Araújo, D. (2011). Representative learning design and functionality of research and practice in sport. *Journal of Sport and Exercise Psychology*, 33(1), 146–55. https://doi.org/ 10.1123/jsep.33.1.146
- Preatoni, E., Hamill, J., Harrison, A., Hayes, K., Emmerik, R., Wilson, C., & Rodano, R. (2013). Movement variability and skills monitoring in sports. *Sports Biomechanics*, *12*(2), 69–92. https://doi.org/10.1080/14763141.2012. 738700
- Promsri, A. (2022). Sex Difference in Running Stability Analyzed Based on a Whole-Body Movement: A Pilot Study. Sports, 10(9), 138. https://doi.org/ 10.3390/sports10090138
- Quintana-Dupue, JC, and Saupe, D. (2013). Evidence of chaos in indoor pedaling motion using non-linear methods. In D.M. Peters & P. O'Donoghue (Eds.), *Performance Analysis of Sport IX* (pp. 203–14). Routledge.
- Raffalt, P. C., Kent, J. A., Wurdeman, S. R., & Stergiou, N. (2019). Selection procedures for the largest lyapunov exponent in gait biomechanics. *Annals of Biomedical Engineering*, 47(4), 913–23. https://doi.org/10. 1007/s10439-019-02216-1
- Raffalt, P. C., Senderling, B., & Stergiou, N. (2020). Filtering affects the calculation of the largest lyapunov exponent. *Computers in Biology and Medicine*, 122(1103786), 103786–103789. https://doi.org/10.1016/j.comp biomed.2020.103786
- Rahatabad, F., Rangraz, P., Dalir, M., & Nasrabadi, A. (2021). The relation between chaotic feature of surface eeg and muscle force: Case study report. *Journal of Medical Signals and Sensors*, 11(4), 229. https://doi.org/ 10.4103/jmss.JMSS\_47\_20
- Reynard, F., & Terrier, P. (2015). Role of visual input in the control of dynamic balance: Variability and instability of gait in treadmill walking while blindfolded. *Experimental Brain Research*, 233(4), 1031–40. https://doi. org/10.1007/s00221-014-4177-5
- Reynard, F., Vuadens, P., Deriaz, O., Terrier, P., & Paul, F. (2014). Could local dynamic stability serve as an early predictor of falls in patients with moderate neurological gait disorders? A reliability and comparison study in healthy individuals and in patients with paresis of the lower extremities. *PLoS One*, *9*(6), e100550–e59. https://doi.org/10.1371/journal.pone. 0100550
- Robalo, R. A., Diniz, A. M., Fernandes, O., & Passos, P. J. (2021). The role of variability in the control of the basketball dribble under different perceptual setups. *European Journal of Sport Science*, 21(4), 521–530. https:// doi.org/10.1080/17461391.2020.1759695
- Santuz, A., Ekizos, A., Eckardt, N., Kibele, A., & Arampatzis, A. (2018). Challenging human locomotion: stability and modular organisation in unsteady conditions. *Science Report*, 8(1). https://doi.org/10.1038/ s41598-018-21018-4

- Schall, M. C., Fethke, N. B., Chen, H., Oyama, S., & Douphrate, D. I. (2016). Accuracy and repeatability of an inertial measurement unit system for field-based occupational studies. *Ergonomics*, 59(4), 591–602. https://doi.org/10.1080/00140139.2015.1079335
- Sejdić, E., Findlay, B., Merey, C., & Chau, T. (2013). The effects of listening to music or viewing television on human gait. *Computers* in Biology and Medicine, 43(10), 1497–501. https://doi.org/10.1016/j. compbiomed.2013.07.019
- Slegers, N., Lee, D., & Wong, G. (2021). The relationship of intra-individual release variability with distance and shooting performance in basketball. *Journal of Sports Science and Medicine*, 20(3), 508–15. https://doi.org/10. 52082/jssm.2021.508
- Smith, B. A., Stergiou, N., & Ulrich, B. D. (2010). Lyapunov exponent and surrogation analysis of patterns of variability: Profiles in new walkers with and without down syndrome. *Motor Control*, 14(1), 126–42. https://doi.org/10.1123/mcj.14.1.126
- Stenum, J., Bruijn, S. M., & Jensen, B. R. (2014). The effect of walking speed on local dynamic stability is sensitive to calculation methods. *Journal of Biomechanics*, 47(15), 3776–79. https://doi.org/10.1016/j.jbiomech.2014.09.020
- Stergiou, N., Buzzi, U. H., Kurz, M. J., & Heidel, J. (2004). Nonlinear tools in human movement. In N. Stergiou (Ed.), *Innovative analyses of human movement* (pp. 63–87). Human Kinetics.
- Stergiou, N., & Decker, L. M. (2011). Human movement variability, nonlinear dynamics, and pathology: Is there a connection? *Human Movement Science*, 30(5), 869–88. https://doi.org/10.1016/j.humov. 2011.06.002
- Straussfogel, D., & von Schilling, C. (2009). Systems theory. In R. Kitchin & N. Thrift (Eds.), *International encyclopedia of human geography* (pp. 151–158). Elsevier.
- Strongman, C., & Morrison, A. (2021). Evaluating dynamic similarity of fixed, self-selected and anatomically scaled speeds in non-linear analysis of gait during treadmill running. *Human Movement Science*, 76, 102768. https://doi.org/10.1016/j.humov.2021.102768

- Terrier, P., & Reynard, F. (2015). Effect of age on the variability and stability of gait: A cross-sectional treadmill study in healthy individuals between 20 and 69 years of age. *Gait & Posture*, 41(1), 170–4. https://doi.org/10.1016/j.gaitpost.2014.09.024
- Toebes, M. J. P., Hoozemans, M. J. M., Furrer, R., Dekker, J., & van Dieen, J. H. (2012). Local dynamic stability and variability of gait are associated with fall history in elderly subjects. *Gait & Posture*, 36(3), 527–31. https://doi.org/10.1016/j.gaitpost.2012.05.016
- Van Emmerik, R. E. A., Rosenstein, M. T., McDermott, W. J., & Hamill, J. (2004). A nonlinear dynamics approach to human movement. *Journal of Applied Biomechanics*, 20(4), 396–420. https://doi.org/10.1123/jab.20.4.396
- van Schooten, K. S., Rispens, S. M., Pijnappels, M., Daffertshofer, A., & van Dieen, J. H. (2013). Assessing gait stability: The influence of state space reconstruction on inter- and intra-day reliability of local dynamic stability during over-ground walking. *Journal of Biomechanics*, 46(1), 137–41. https://doi.org/10.1016/j.jbiomech.2012. 10.032
- Walsh, G. S. (2021). Dynamics of Modular Neuromotor Control of Walking and Running during Single and Dual Task Conditions. *Neuroscience*, 465, 1–10. https://doi.org/10.1016/j.neuroscience.2021.04.004
- Wilkinson, R. D., & Lichtwark, G. A. (2021). Evaluation of an inertial measurement unit-based approach for determining centre-of-mass movement during non-seated cycling. *Journal of Biomechanics*, 126(110441), 1–7. https://doi.org/10.1016/j.jbiomech.2021.110441
- Wurdeman, S. R. (2018a). Lyapunov Exponent. In N. Stergiou (Ed.), Nonlinear analysis for human movement variability (1st ed., pp. 83–108). CRC Press.
- Wurdeman, S. R. (2018b). State space reconstruction. In N. Stergiou (Ed.), Nonlinear analysis for human movement variability (1st ed., pp. 55–82). CRC Press.
- Yang, F., & King, G. A. (2016). Dynamic gait stability of treadmill versus overground walking in young adults. *Journal of Electromyography and Kinesiology*, 31, 81–87. https://doi.org/10.1016/j.jelekin.2016.09.004

## Appendix 1: Methodological quality assessment

Ref No.	Study	1.	2.	3.	4.	5.	6.	7.	8.	9.	Total score (18 points max)	Quality (%)
1	(Abbasi, Zamanian and Svoboda 2019)	2	0	2	2	2	2	1	2	0	13	72.2
2	(Arshi, Mehdizadeh and Davids 2015)	2	0	2	2	2	1	2	2	0	13	72.2
3	(Ataabadi et al. 2021)	2	2	2	2	1	2	2	2	1	16	88.9
4	(Cignetti, Schena and Rouard 2009)	2	0	1	2	0	2	2	2	1	12	66.7
5	(Ekizos, Santuz and Arampatzis 2018)	2	0	2	2	0	2	2	2	0	12	66.7
6	(Ekizos et al. 2018)	2	0	2	2	2	2	2	2	2	16	88.9
7	(Ekizos, Santuz and Arampatzis 2017)	2	0	2	2	2	2	2	2	0	14	77.8
8	(Fohrmann et al. 2022)	2	0	2	2	1	2	2	2	2	15	83.3
9	(Frank, Prentice and Callaghan 2019)	2	0	2	2	0	1	2	1	0	10	55.6
10	(Hamacher et al. 2018)	2	N/A	0	2	N/A	2	2	2	0	10	71.4
11	(Hoenig et al. 2019)	2	0	2	2	0	2	2	2	0	12	66.7
12	(Hollander, Hamacher and Zech 2021)	2	0	2	2	0	2	2	2	1	13	72.2
13	(Jordan et al. 2009)	2	0	1	0	2	2	2	2	0	11	61.1
14	(Larson et al. 2021)	2	0	2	1	0	1	2	2	0	10	55.6
15	(Look et al. 2013)	1	0	0	1	0	2	2	2	0	8	44.4
16	(Mahaki et al. 2020)	1	0	2	2	2	2	2	2	2	15	83.3
17	(Mehdizadeh, Arshi and Davids 2014a)	2	0	2	2	1	2	2	2	0	13	72.2
18	(Mehdizadeh, Arshi and Davids 2014b)	2	0	2	1	1	2	2	2	0	12	66.7
19	(Mehdizadeh, Arshi and Davids 2016)	2	0	2	2	1	2	2	2	0	13	72.2
20	(Nessler et al. 2015)	2	0	2	2	0	2	2	2	0	14	77.8
21	(Ogaya et al. 2021)	2	0	2	2	0	2	2	2	2	14	77.8
22	(Padulo et al. 2023)	2	0	2	2	0	2	2	2	2	14	77.8
23	(Promsri 2022)	2	2	2	1	0	2	2	1	0	12	66.7
24	(Quintana-Duque and Saupe 2014)	0	N/A	0	0	0	0	1	2	1	4	25
25	(Raffalt et al. 2020)	2	0	2	2	0	2	2	2	0	12	55.6
26	(Raffalt et al. 2019)	2	0	2	2	0	1	2	2	0	11	61.1
27	(Rahatabad et al. 2021)	1	N/A	0	N/A	0	2	0	0	0	3	21.4
28	(Robalo et al. 2021)	2	2	1	2	0	2	0	2	0	11	61.1
29	(Santuz et al. 2018)	2	0	2	2	2	2	2	2	1	15	83.3
30	(Strongman and Morrison 2021)	2	0	2	2	1	2	2	2	0	13	72.2
31	(Walsh 2021)	2	0	2	2	2	2	2	2	0	14	77.8
Mean		1.84	0.21	1.65	1.73	0.73	1.81	1.81	1.87	0.48	12.1	67.6

Ref = reference; no. = number; max = maximum; 0, clearly no; 1, maybe or inadequate information; 2, clearly yes.

Ref No.	Study	Running	No. of Groups/Conditions	Segment/Joint	LyE Range
-	(Abbasi, Zamanian and Svoboda 2019)	Cycling	3 Without stretching (WS), Static stretching (SS), Dynamic stretching (DS)	Ankle, Knee, Hip (joint angles) Soleus (SL), Rectus Femoris (RF), Biceps Femoris (BF), Tiblias Anterior (TA), Gastrocnemius (G), Vastus Medialis (VM) (EMG)	Hip – 0.96 – 1.00
					Knee – 0.53 – 0.86
					Ankle – 0.97 – 1.25
					SL - 0.68 - 0.89
					RF – 0.73 – 1.09
					BF – 0.62 – 1.10
					TA – 0.54 – 0.99
					G – 0.90 – 1.18
					VM - 0.63 - 1.09
7	(Arshi, Mehdizadeh and Davids 2015)	Running	2 – Fw + La	Trunk	Fw- AP 1.57, ML 1.05
					La AP 1.15, ML 2.01
ε	(Ataabadi et al. 2021)	Running	3 conditions SSR, PSR, FSR	Ankle, Knee, Hip	Ankle – SSR 1.244 – 1.404, PSR 1.293 – 1.383, FSR 1.379 – 1.471
					Knee – SSR 0.916 – 1.269, PSR 0.945 – 1.242, FSR 1.045 – 1.321
					Hip – SSR 1.252 – 1.426, PSR 1.68 – 1.426, FSR 1.286 – 1.479
4	(Cignetti, Schena and Rouard 2009)	Skiing	surrogation vs. regular LyE's B + E	Leg, Arm	Arm Regular B – 0.18, E – 0.24 Surrocated B – 0.34 E 0.38
					Leg Regular B – 0.22, E – 0.28 Surrogated B – 0.35, E 0.38
Ŝ	(Ekizos, Santuz and Arampatzis 2018)	Running	3 – LT, ST, CG Pre vs. Post	Trunk	ST – 1.81 – 1.86
					LT – 1.78 – 1.79
					CG – 1.80 – 1.75
9	(Ekizos et al. 2018)	Running	6 marker sets	Trunk	1.67 – 1.97
7	(Ekizos, Santuz and Arampatzis 2017)	Running	Shod vs. Barefoot	Trunk	Shod – 1.62 – 1.95
					Barefoot - 1.68 - 2.08
œ	(Fohrmann et al. 2022)	Running	3 conditions – T, OO, Ol	Sternum, Sacrum, Tibia, Foot	Sternum – T 1.77 – 1.91, OO – 1.84 – 2.00, OI 1.85 – 2.00
					Sacrum – T 4.28 – 4.39, OO 4.44 – 4.50, Ol 4.45 – 4.53
					(Continued)

Appendix 2: LyE Extraction Values

(Contir	ued).				
Ref No.	Study	Running	No. of Groups/Conditions	Segment/Joint	LyE Range
					Tibia – T 4.43 – 4.46, OO – 4.89 – 4.96, OI 4.90 – 4.92 Foot – T 2.34 – 2.45, OO 2.56 – 2.67, OI 2.57 – 2.62
6	(Frank, Prentice and Callaghan 2019)	Running	4 shoe types – novice vs. trained	Ankle, Knee, Hip	Ankle Novice 1.41 – 1.49 Trained 1.19 – 1.23 Knee
					Novice 1.21 – 1.25 Trained 0.72 – 0.76 Hip Novice 1.13 – 1.21 Elite 0.61 – 0.64
10	(Hamacher et al. 2018)	Kayaking		Trunk, Left and Right UA, left and right hand (H)	Trunk – 0.001 Left UA – 0.658
					kight UA – 0.634 Left Hand – 0.254 Right Hand – 0.165
11	(Hoenig et al. 2019)	Running	2 groups – C vs. R B, M, E	Thorax, Pelvis, Foot	Thorax C – 0.91 – 0.93 R – 0.90 – 1.01
					Pelvis C – 2,42 – 2.54 R – 2.64 – 2.84 Ecot
					C - 6.98 - 7.04 C - 6.98 - 7.04 R - 7.23 - 7.33
12	(Hollander, Hamacher and Zech 2021)	Running	Shod vs. Barefoot – B, M, E	Shank	BShod 47.25 – 48.46Barefoot 49.03 – 50.74
					M Shod 47.71 – 48.14 Barefoot 48.99 – 50.12 -
					E Shod 48.14 – 48.71 Barefoot 49.55 – 51.04
13	(Jordan et al. 2009)	Running	5 speeds	Head, Ankle	Ankle – 0.080 - 0.092 Head – 0.063 – 0.070
14	(Larson et al. 2021)	Jumping	2 groups PU vs. PI	Ankle, Knee, Hip	Ankle PU – 0.04 – 0.17 Pl 0.07 – 0.17
					Knee PU - 0.04 – 0.16 Pl 0.04 – 0.17
					Hip PU 0.04 – 0.17 PI 0.06 – 0.17
15	(Look et al. 2013)	Running	2 Groups – amputation (A) vs. no amputation (NA) Different sneeds randing from 3 – 9 m/s	Hip, Knee, Sacrum	NA – knee – 0.095 – 0.137 m/ s
					Ă 0.090 – 0.138

<sup>(</sup>Continued)

(Continu	.(pər					
Ref No.	Study	Running	No. of Groups/Conditions		Segment/Joint	LyE Range
						Hip – 0.098 – 0.119 NA 0.075 – 0.131 A Sacrum ML A 0.006 – 0.16 Vertical (vert) A 0.09 – 0.20 NA ML 0.02 – 0.24
16	(Mahaki et al. 2020)	Running	Pre – Post (recorded every min, 3 minutes	Frunk		NA vert 0.19 – 0.25 Sagittal (pre) 0.50 – 1.41; (2004) 0.40 – 1.60
						(post) 0.46 - 1.56 Frontal (pre) 1.33 - 2.14; (post) 0.96 - 2.73
						Horizontal (pre) 0.37 – 3.02, (post) 0.33 – 2.75
17	(Mehdizadeh, Arshi and Davids 2014a)	Running	2 conditions – forward (Fw) and backward (B) running; 2 groups (skilled (S) vs. unskilled (US))	Ankle, Knee, Hip, Sacrum		Ankle Fw S 2.27 – 3.06, B S 2.20 – 3.28, Fw US 3.89 – 5.33, B US 3.62 – 4.83
						Knee Fw S 2.56 – 2.69, B S 2.35 – 2.65, Fw US 4.36 – 4.98, B US 3.89 – 4.71
						Hip Fw S 1.70 – 2.47, B S 1.93 – 2.64, Fw US 3.05 – 3.92, B US 3.10 – 3.98
						Sacrum Fw S 1.32 – 2.10, B S 1.69 – 2.25, Fw US 1.74 – 2.55, B US 1.45 – 2.19
18	(Mehdizadeh, Arshi and Davids 2014b)	Running	2 running speeds (80, 100, 120% preferred running speed) – long term (LT) vs. short term (ST) LyE B vs. Fw running		Trunk	Fw ST - 80% 1.15 - 1.45, 100% 1.14 - 1.55, 120% 1.25 - 1.53
						B ST – 80% 1.50 – 2.16, 100% 1.69 – 2.41, 120% 1.78 – 2.38
						Fw LT - 80% 0.01 - 0.08 100% 0.02 - 0.09, 120% 0.01 - 0.09
						B LT - 80% 0.03 - 0.08, 100% 0.03 - 0.11, 120% 0.03 - 0.10
19	(Mehdizadeh, Arshi and Davids 2016)	Running	3 running conditions – Fw, B, Lateral (La) 3 Speeds		Ankle	80% - Fw ML 1.03 AP 1.18; B ML 1.43, AP 1.76; La ML 1.55, AP 1.27
						100% - Fw ML 1.02, AP 1.18; B ML 1.41, AP 1.79; La ML 1.55, AP 1.32
						120% - Fw ML 1.02, AP 1.22; B ML 1.40, AP 1.74; La ML 1.55, AP 1.32
20	(Nessler et al. 2015)	Paddling	Wetsuit vs. Non-wetsuit		Wrist	Wetsuit – 1.49
						Non-weisunt – 1.38 (Continued)

	LyE Range	Ankle FL/Ex 0.85 vs. 0.90 Inv/Ev 0.65 vs. 0.71	Knee FL/Ex 1.04 vs. 1.04	Hip FL/Ex 1.00 vs.1.01 Abd/Add 0.74 vs. 0.78	INT/EXT 0.52 vs. 0.55	Lumbar Spine FL/EX 0.38 vs. 0.50 La bend 0.46 vs. 0.57	Rotation 1.19 vs. 1.19	0CON - 0.007	2CON 0.020	2IES 0.009	M 2.2 5.7	F 2.3 – 7.5	Knee Angle • 140 N 100 rpm 0.22 • 120 N 100 rpm 0.19 • 140N 80rpm 0.21	<ul> <li>120N 80rpm 0.32</li> </ul>	Knee x-coordinate • 140 N 100 rpm 0.18 • 120 N 100 rpm 0.24 • 140N 80rpm 0.20 • 120N 80rpm 0.26	Hip 0.27 – 0.58	Knee 0.32 – 0.43	Ankle 0.60 – 0.82	COM displacement AP 1 79 - 0 48 - 0 70	ML 0.72 - 1.04	Vert 1./9 - 0.6/ - 0./6 (Continued)
	Segment/Joint	Ankle, Knee, Hip, Lumbar Spine						N/A – Stride Frequency			Whole Body		Knee			Hip, Knee, Ankle, Pelvis					
	No. of Groups/Conditions	Pre vs. Post						3 conditions – 0% gradient (OCON), 2% gradient fixed speed (2CON), 2% gradient at isoefficiency speed (2IES)			5 principal movements in running – males (M) and females (F)		4			8 Speeds					
	Running	Running						Running			Running		Cycling			Running					
ed).	Study	(Ogaya et al. 2021)						(Padulo et al. 2023)			(Promsri 2022)		(Quintana-Duque and Saupe 2014)			(Raffalt et al. 2020)					
(Continu	Ref No.	21						22			23		24			25					

LyE Range	Rosenstein N1 1.79 - 0.86 - 1.79 2.46 - 0.85 - 1.67 N2 1.79 - 0.83 - 1.64 2.46 - 0.82 - 1.64 N3 1.79 - 0.92 - 1.72 N4 1.79 - 0.93 - 1.61 2.46 - 0.95 - 1.61 N1 1.79 - 0.95 - 1.61 2.46 - 1.55 - 2.28 2.46 - 1.55 - 2.28 2.46 - 1.57 - 2.28 N3 1.79 - 1.86 - 2.19 2.46 - 1.67 - 2.19 2.46 - 1.67 - 2.19 2.46 - 1.57 - 2.18 N3 1.79 - 1.26 - 1.88 N3 1.79 - 1.27 - 1.88 N4 1.79 - 1.27 - 1.88 N4 N3 N3 N3 N3 N3 N3 N3 N3 N3 N3 N3 N3 N3	2.46 - 1.15 - 1.77 1.12 - 1.40 Amateurs Wrist • NO 2.90 - 3.20 • VO 2.61 - 2.97 • AO 3.11 - 3.42 BO 2.80 - 3.07 Elbow • NO 3.10 - 3.51 • VO 2.81 - 3.61 • AO 3.10 - 3.51 • VO 2.81 - 3.61 • O 3.31 - 3.42 • Solutioner • O 1.36 - 3.35 • AO 2.34 - 3.81 • O 1.37 - 2.19 (Continued)
Segment/Joint	Sacrum	Brain
		Wrist, Elbow, Should
No. of Groups/Conditions	4 normalisation (N) procedures 2 algorithms at 2 speeds (1.79, 2.46)	5 Sensory Motor Areas Amateur vs. Professional 4 conditions NO – no occlusion VO – visual occlusion AO – auditory occlusion BO – both occlusion
Running	Running	Weighted Raise Basketball Dribbling
Study	(Raffalt et al. 2019)	(Robalo et al. 2021) (Robalo et al. 2021)
Ref No.	70	28

2012 😉 L. WINTER ET AL.

(Continued).

(Contii	ued).				
Ref No.	Study	Running	No. of Groups/Conditions	Segment/Joint	LyE Range
					Professionals Wrist • NO - 2.17 - 3.00 • VO 2.47 - 3.83 • AO 2.40 - 3.75 BO 2.20 - 3.87 Elbow • NO - 2.84 - 3.83 • AO - 2.84 - 3.83 AO - 2.48 - 3.87 Shoulder • NO - 2.19 - 2.68 • VO - 2.52 - 2.77 • AO - 2.52 - 2.77 • AO - 2.52 - 2.77
29	(Santuz et al. 2018)	Running	Unstable vs. Even	Trunk	<ul> <li>BO 2.28 - 2.71</li> <li>Unstable - 2.031</li> <li>Evan - 1 936</li> </ul>
30	(Strongman and Morrison 2021)	Running	3 Speeds Froude Fixed Self-selected	Hip, Knee	Hip Froude 0.84 – 2.16 Fixed 0.85 – 2.03 Self-selected – 0.58 – 2.37 Knee Froude 0.78 – 1.84 Fixed 0.95 – 2.18 Self-selected – 0.92 – 2.18
31	(Walsh 2021)	Running	Single Task vs. Dual Task	Trunk (JA); Muscle synergy	LS LS ST PRS - 1.80 - 5.35 ST PRS 120 - 1.89 - 4.85 DT PRS - 2.19 - 5.31 DT PRS 120 - 1.94 - 5.31 Muscle Synergy ST PRS - 0.26 - 0.30 ST PRS - 0.26 - 0.30 DT PRS - 0.26 - 0.29 DT PRS 120 - 0.29
PI = 1	forward, La = la end, LT = Long-term, ST = previously injured.	teral, AP = ant = Short-term (ST), CG	terior-posterior, La = lateral, SSR = Slow spee 3 = Control. T = treadmill, OO = overground outdoor, OI = o	d running, PSR = preferred-speed running, FSR = 1 verground indoor, UA = upper arm, C = competitive, R = recreation	st-speed running, B = beginning, al, M = middle, PU = previously un-injured,