

## Appendix D “Included studies”

1. Andersen E, Bang-Kittilsen G, Bigseth TT, et al. Effect of high-intensity interval training on cardiorespiratory fitness, physical activity and body composition in people with schizophrenia: a randomized controlled trial. *BMC Psychiatry*. 2020;20(1):425. doi:10.1186/s12888-020-02827-2
2. Armstrong HF, Bartels MN, Paslavski O, et al. The impact of aerobic exercise training on cardiopulmonary functioning in individuals with schizophrenia. *Schizophrenia Research*. 2016;173(1-2):116-117. doi:10.1016/j.schres.2016.03.009
3. Bhatia T, Mazumdar S, Wood J, et al. A randomised controlled trial of adjunctive yoga and adjunctive physical exercise training for cognitive dysfunction in schizophrenia. *Acta Neuropsychiatr*. 2017;29(2):102-114. doi:10.1017/neu.2016.42
4. Battaglia G, Alesi M, Inguglia M, et al. Soccer practice as an add-on treatment in the management of individuals with a diagnosis of schizophrenia. *Neuropsychiatr Dis Treat*. 2013;9:595-603. doi:10.2147/NDT.S44066
5. Beebe LH, Tian L, Morris N, Goodwin A, Allen SS, Kuldau J. Effects of exercise on mental and physical health parameters of persons with schizophrenia. *Issues Ment Health Nurs*. 2005;26(6):661-676. doi:10.1080/01612840590959551
6. Bredin SSD, Warburton DER, Lang DJ. The health benefits and challenges of exercise training in persons living with schizophrenia: a pilot study. *Brain Sci*. 2013;3(2):821-848. doi:10.3390/brainsci3020821
7. Caponnetto P, Auditore R, Maglia M, Pipitone S, Inguscio L. Psychological wellness, yoga and quality of life in patients affected by schizophrenia spectrum disorders: A pilot study. *Ment Illn*. 2019;11(1):8003. doi:10.4081/mi.2019.8003
8. Curcic D, Stojmenovic T, Djukic-Dejanovic S, et al. Positive impact of prescribed physical activity on symptoms of schizophrenia: randomized clinical trial. *Psychiatr Danub*. 2017;29(4):459-465. doi:10.24869/psyd.2017.459
9. Andrade e Silva B, Cassilhas RC, Attux C, et al. A 20-week program of resistance or concurrent

- exercise improves symptoms of schizophrenia: results of a blind, randomized controlled trial. *Braz J Psychiatry*. 2015;37(4):271-279. doi:10.1590/1516-4446-2014-1595
10. Heggelund J, Nilsberg GE, Hoff J, Morken G, Helgerud J. Effects of high aerobic intensity training in patients with schizophrenia: a controlled trial. *Nord J Psychiatry*. 2011;65(4):269-275. doi:10.3109/08039488.2011.560278
  11. Heggelund J, Morken G, Helgerud J, Nilsberg GE, Hoff J. Therapeutic effects of maximal strength training on walking efficiency in patients with schizophrenia - a pilot study. *BMC Res Notes*. 2012;5(1):344. doi:10.1186/1756-0500-5-344
  12. Ho SS, Dhaliwal SS, Hills AP, Pal S. The effect of 12 weeks of aerobic, resistance or combination exercise training on cardiovascular risk factors in the overweight and obese in a randomized trial. *BMC Public Health*. 2012;12:704. doi:10.1186/1471-2458-12-704
  13. Hsu C-C, Liang C-S, Tai Y-M, Cheng S-L. Incongruent changes in heart rate variability and body weight after discontinuing aerobic exercise in patients with schizophrenia. *Int J Psychophysiol*. 2016;109:132-137. doi:10.1016/j.ijpsycho.2016.08.011
  14. Ikai S, Uchida H, Suzuki T, Tsunoda K, Mimura M, Fujii Y. Effects of yoga therapy on postural stability in patients with schizophrenia-spectrum disorders: a single-blind randomized controlled trial. *J Psychiatr Res*. 2013;47(11):1744-1750. doi:10.1016/j.jpsychires.2013.07.017
  15. Kaltsatou ACH, Kouidi EI, Anifanti MA, Douka SI, Deligiannis AP. Functional and psychosocial effects of either a traditional dancing or a formal exercising training program in patients with chronic heart failure: a comparative randomized controlled study. *Clin Rehabil*. 2014;28(2):128-138. [https://journals.sagepub.com/doi/abs/10.1177/0269215513492988?casa\\_token=F24xGAANjgYAAA:VDC4Ggl2HbIBkPxXXeC3drGIZZ0GAwhQlvHLdkmX\\_gbkn\\_fKtO5JwavfvYpNDUdsp2Zo-11KNynjoA](https://journals.sagepub.com/doi/abs/10.1177/0269215513492988?casa_token=F24xGAANjgYAAA:VDC4Ggl2HbIBkPxXXeC3drGIZZ0GAwhQlvHLdkmX_gbkn_fKtO5JwavfvYpNDUdsp2Zo-11KNynjoA)
  16. Kim H-J, Song B-K, So B, Lee O, Song W, Kim Y. Increase of circulating BDNF levels and its relation to improvement of physical fitness following 12 weeks of combined exercise in chronic patients with schizophrenia: a pilot study. *Psychiatry Res*. 2014;220(3):792-796. doi:10.1016/j.psychres.2014.09.020

17. Kimhy D, Vakhrusheva J, Bartels MN, et al. The Impact of Aerobic Exercise on Brain-Derived Neurotrophic Factor and Neurocognition in Individuals With Schizophrenia: A Single-Blind, Randomized Clinical Trial. *Schizophr Bull.* 2015;41(4):859-868. doi:10.1093/schbul/sbv022
18. Loh SY, Abdullah A, Abu Bakar AK, Thambu M, Nik Jaafar NR. Structured Walking and Chronic Institutionalized Schizophrenia Inmates: A pilot RCT Study on Quality of Life. *Glob J Health Sci.* 2015;8(1):238-248. doi:10.5539/gjhs.v8n1p238
19. Marzolini S, Jensen B, Melville P. Feasibility and effects of a group-based resistance and aerobic exercise program for individuals with severe schizophrenia: A multidisciplinary approach. *Mental Health and Physical Activity.* 2009;2(1):29-36. doi:10.1016/j.mhpa.2008.11.001
20. Pajonk F-G, Wobrock T, Gruber O, et al. Hippocampal plasticity in response to exercise in schizophrenia. *Arch Gen Psychiatry.* 2010;67(2):133-143. doi:10.1001/archgenpsychiatry.2009.193
21. Röhricht F, Priebe S. Effect of body-oriented psychological therapy on negative symptoms in schizophrenia: a randomized controlled trial. *Psychol Med.* 2006;36(5):669-678. doi:10.1017/S0033291706007161
22. Ryu J, Jung JH, Kim J, et al. Outdoor cycling improves clinical symptoms, cognition and objectively measured physical activity in patients with schizophrenia: A randomized controlled trial. *J Psychiatr Res.* 2020;120:144-153. doi:10.1016/j.jpsychires.2019.10.015
23. Scheewe TW. Effects of exercise therapy on cardiovascular fitness and the metabolic syndrome in schizophrenia: a randomized clinical trial. *Schizophrenia Research.* 2012;136:S47. doi:10.1016/s0920-9964(12)70172-x
24. Scheewe TW, van Haren NEM, Sarkisyan G, et al. Exercise therapy, cardiorespiratory fitness and their effect on brain volumes: a randomised controlled trial in patients with schizophrenia and healthy controls. *Eur Neuropsychopharmacol.* 2013;23(7):675-685. doi:10.1016/j.euroneuro.2012.08.008
25. Shimada T, Ito S, Makabe A, Yamanushi A, Takenaka A, Kobayashi M. Aerobic exercise and cognitive functioning in schizophrenia: A pilot randomized controlled trial. *Psychiatry Res.* 2019;282:112638. doi:10.1016/j.psychres.2019.112638

26. Shimizu N, Umemura T, Matsunaga M, Hirai T. An interactive sports video game as an intervention for rehabilitation of community-living patients with schizophrenia: A controlled, single-blind, crossover study. *PLoS One*. 2017;12(11):e0187480. doi:10.1371/journal.pone.0187480
27. Svatkova A, Mandl RCW, Scheewe TW, Cahn W, Kahn RS, Hulshoff Pol HE. Physical Exercise Keeps the Brain Connected: Biking Increases White Matter Integrity in Patients With Schizophrenia and Healthy Controls. *Schizophr Bull*. 2015;41(4):869-878. doi:10.1093/schbul/sbv033
28. Varambally S, Gangadhar BN, Thirthalli J, et al. Therapeutic efficacy of add-on yogasana intervention in stabilized outpatient schizophrenia: Randomized controlled comparison with exercise and waitlist. *Indian J Psychiatry*. 2012;54(3):227-232. doi:10.4103/0019-5545.102414