

Running title: Overweight, Obesity and Intellectual Disabilities

Prevalence of Overweight and Obesity among Children and Adolescents with Intellectual Disabilities: A Systematic Review and Meta-Analysis

Christophe Maïano<sup>1,2,3</sup>, Olivier Hue<sup>4</sup>, Alexandre J. S. Morin<sup>2</sup>, Grégory Moullec<sup>3,5</sup>

<sup>1</sup> Cyberpsychology Laboratory, Department of Psychoeducation and Psychology, Université du Québec en Outaouais (UQO), Canada.

<sup>2</sup> Institute for Positive Psychology and Education, Australian Catholic University, Australia.

<sup>3</sup> Department of Psychoeducation and Psychology, Université du Québec en Outaouais (UQO), Canada.

<sup>4</sup> Department of Physical Activity Sciences, Université du Québec à Trois-Rivières, Canada

<sup>5</sup> Research Center, Hôpital du Sacré-Coeur de Montréal, affiliated Hospital of Université de Montréal, Canada

**Acknowledgements:** The preparation of this meta-analysis was supported by grants from the Social Sciences and Humanities Research Council of Canada (430-2012-0091, 435-2014-0909) awarded to the first three authors and the Australian Research Council (DP140101559) awarded to the first and third authors.

**Keywords.** Age; Additional diagnosis; Geographic regions; ID levels; Recruitment setting; Sex.

**Corresponding author:** Christophe Maïano, Université du Québec en Outaouais, Campus de Saint-Jérôme, Département de Psychoéducation et de Psychologie, 5 rue Saint-Joseph, Saint-Jérôme, Québec J7Z 0B7, email: christophe.maiano@uqo.ca

**Potential conflict of interest.** The authors declare no conflict of interest.

This is the prepublication version of the following manuscript:  
Maïano, C., Hue, O., Morin, A. J. S., & Moullec, G. (2016). Prevalence of overweight and obesity among children and adolescents with intellectual disabilities: A systematic review and meta-analysis. *Obesity Reviews*, 17(7), 599-611  
© 2016. This paper is not the copy of record and may not exactly replicate the authoritative document published in *Obesity Reviews*.

### Abstract

Although there have been numerous studies examining the prevalence of overweight and obesity among children and adolescents with intellectual disabilities, they have not yet been systematically integrated and synthesized through a systematic quantitative review process. The purpose of this systematic review and meta-analysis was to determine: (a) the prevalence of overweight/obesity among children and adolescents with intellectual disabilities; (b) the sources of heterogeneity in studies reporting the prevalence of overweight/obesity in this population; and (c) the risk of overweight/obesity in this population compared with their typically developing peers. A systematic literature search was performed and 16 studies, published between 1985 and 2015, met the inclusion criteria. The resulting pooled prevalence estimates for overweight, overweight-obesity, and obesity were respectively (a) 15%, 30%, and 13% in children; and (b) 18%, 33%, and 15% in adolescents. Subgroup analyses showed significant variations in the pooled prevalence estimates as a function of geographic region, recruitment setting, additional diagnoses, and norms used to define overweight or obesity. The findings also showed adolescents with intellectual disabilities to be respectively 1.54 and 1.80 times more at risk of overweight-obesity and obesity than typically developing adolescents. Unfortunately, no such comparison is available for children.

**Abbreviations used in the text:** B-M: Begg and Mazumdar; DD: developmental disabilities; ID: intellectual disabilities; IOTF: International Obesity Task Force; PRISMA: Preferred Reporting Items for Systematic reviews and Meta-Analyses; STROBE: STrengthening the Reporting of OBservational studies in Epidemiology statement; TD: typically developing; USA: United States of America.

### Introduction

Over the past three decades, a considerable amount of research regarding the prevalence of overweight and obesity among youth presenting various disabilities or special needs has been conducted and synthesized (1-6). However, the prevalence of overweight/obesity among children and adolescents with intellectual disabilities (ID; i.e., characterized by a deficit in intellectual functioning accompanied by a deficit in adaptive functioning and an onset during the developmental period) (7) has not been so closely examined. The first narrative review on this topic, published in 2011 (8), reported a prevalence of overweight excluding obesity ranging from 11% to 25%, and a prevalence of obesity from 7% to 36% in children and adolescents with ID.

Since then, five additional reviews have summarized the research examining the prevalence of overweight and/or obesity among children and/or adolescents with ID (10) or developmental disabilities (DD). The DD categorisation is broader than ID and encompasses Fragile X syndrome, Down syndrome, pervasive developmental disorders, fetal alcohol spectrum disorders, cerebral palsy, and ID (9,11-13). However, only a limited number of studies focusing specifically on children and/or adolescents with ID were included in these previous reviews.

Moreover, these reviews suffer from important limitations. First, five of these reviews (8-11,13) were non-systematic, and the sole systematic review (12) was restricted to the relationship between parental/parenting factors and obesity among children and/or adolescents with DD. Therefore, numerous studies on children and/or adolescents with ID could have been missed. Second, only two reviews have included studies comparing the prevalence of overweight or obesity among children and/or adolescents with ID and their typically developing (TD) peers (8,11). Furthermore, both of these reviews included very few such comparative studies, and most of them overlapped between the two reviews. Finally, no reviews have yet quantitatively examined whether the heterogeneity observed in the overweight/obesity prevalence of children and/or adolescents with ID could be attributed to the participants' characteristics (e.g., age, sex, ID severity, genetic syndromes, geographic region, or recruitment setting) or to the assessment methods (e.g., norms used to define overweight and obesity, measurement of height and weight).

Consequently, the prevalence of overweight and obesity among children and adolescents with an ID remains an underexplored area and there still remain gaps and inconsistencies in the knowledge. Still unknown is: (a) the extent to which children and adolescents with ID are at high risk for overweight, overweight-obesity or obesity; (b) whether the observed prevalence estimates reported in the literature varied when different sources of heterogeneity were considered; and (c) whether children and adolescents with ID are at greater risk for overweight, overweight-obesity or obesity than their TD peers.

In this context, a systematic review and meta-analysis of the prevalence rates of overweight/obesity among children and adolescents with ID appears to be of significant importance. Indeed, a better estimation of this critical public health problem in children and adolescents with ID would encourage scholars, practitioners and policy makers to further develop lifestyle intervention programs (i.e., healthy diet, physical activity, health promotion-education and behavioral modification) designed for tackling or managing weight problems in this population. Therefore, following the Meta-Analysis of Observational Studies in Epidemiology Statement (14), the aims of the current review were to determine, among children and adolescents with ID: (a) the prevalence of overweight, overweight-obesity, and obesity; (b) the sources of heterogeneity in studies reporting the prevalence of overweight, overweight-obesity and obesity; and (c) the risk of overweight, overweight-obesity and obesity, compared with their TD peers.

## Method

### Sources of Information and Search Strategy

A systematic electronic search was conducted in nine databases without imposing any year restriction [Academic Search Complete (1887–2015), Medline (1946–2015), PsycARTICLES (1904–2015), Psychology and Behavioral Sciences Collection (1965–2015), Scopus (1996–2015), CINAHL (1981–2015), Education Sources (1900–2015), ERIC (1966–2015) and SocINDEX (1908–2015)]. Studies were identified using all possible combinations of the following three groups of search terms: (a) intellectual\* disab\* OR learning disab\* OR learning difficult\* OR mental\* retard\* OR developmental dis\* OR developmental del\* OR cogniti\* dis\* OR mental dis\*; AND (b) obes\* OR overweight\* OR fat\* OR weight\* OR body mass index OR nutritional status OR adiposit\*; AND (c) child\* OR adolescen\*OR student\*OR youth\* OR paediatric\* OR pediatric\*. In addition, a hand search was carried out in reference lists of relevant articles and previous literature reviews on children and adolescents with disabilities (1-6), with ID (8,10), or with DD (11-13). Finally, an additional search was also performed in content pages of specific peer-reviewed journals devoted to ID or DD (e.g., *American Journal on Intellectual and Developmental Disabilities*, *Journal of Intellectual Disabilities Research*, *Intellectual and Developmental Disabilities*, *Journal of Applied Research in Intellectual Disabilities*, *Journal of Intellectual and Developmental Disability*, *Research in Developmental Disabilities*). This literature was last updated by hand-search on 5 December 2015.

### Inclusion Criteria

Only studies meeting four specific inclusion criteria were considered eligible for this review. First, study participants had to present an ID. Studies based on mixed samples of participants presenting multiple disabilities were also considered eligible if specific data regarding the prevalence of overweight, overweight-obesity, or obesity were available for children and/or adolescents with ID.

Second, study participants had to be composed of children (age range  $\geq 4 - 11$  years) and/or adolescents (age range  $\geq 11 - 18$  years). Studies including mixed samples of children and adolescents were considered eligible if specific data on the relevant outcomes were available for children and/or adolescents subgroups separately. Studies or samples in a study were not included when the age range of the participants overlapped multiple age categories: infants and children (e.g., 2-10 years), children and adolescents (e.g., 7-12 years, 10-13 years), adolescents and adults (e.g., 12-19 years, 17-20 years), etc.

Third, studies were retained if the prevalence estimates of overweight, overweight-obesity, or obesity were the primary outcome of the study and if they were assessed by means of height/weight measurement indicators, such as the body mass-index (BMI; Cole, 15) and the weight-for-length index (WLI; DuRant and Linder, 16). Here, children/adolescents with ID are categorized as overweight (excluding obesity) when their value of BMI or WLI is  $>$  cut-offs values for normal weight and  $<$  cut-offs values for obesity. Additionally, children/adolescents with ID are categorized as obese when their value of BMI or WLI is  $\geq$  cut-offs values for obesity. Finally, children/adolescents with ID are categorized as overweight-obese (a combined category including overweight and obesity) when their value of BMI or WLI is  $\geq$  the cut-offs values for overweight. This last combined category was considered to permit the consideration of studies in which no distinction was made between overweight and obesity.

The indicators based on height and weight measurement were preferred to other measures of adiposity (e.g., skinfold thickness, waist circumference, bioelectrical impedance analysis) for two reasons. First, previous reviews (8-13) showed that they were the most largely used in overweight/obesity prevalence studies among children and adolescents with ID. Second, a recent

scoping review (17) showed that the validity, reliability and/or sensitivity of other methods (i.e., skinfold thickness, bioelectrical impedance analysis) measuring adiposity in individuals with ID were limited and questionable. Additionally, the use of alternative fatness measurements can be limited or unfeasible in certain occasions (e.g., heterogeneous samples including various subtypes of ID, individuals with additional diagnoses, lack of national norms), and they can introduce a high level of noncompliance in this population (e.g., reluctance to undress). Consequently, we considered that the use and comparison of other body composition measures with the BMI or WLI is premature. When some participants' characteristics (e.g., sex ratio, age) and key information on ID subgroups used to calculate the overweight/obesity rate (e.g., sample size) were not reported in the manuscript, authors were contacted directly to provide the information. Finally, when the same dataset was used in various studies, only one study was included.

Fourth, only original cohort, cross-sectional and case-control studies were included. Reviews, reports, theoretical papers, or single-case studies were excluded. However, a hand search was carried out in reference lists of all previous published reviews on the topic. Finally, studies were retained if they were written in English and published in a peer-reviewed journal.

### **Study Selection and Data Extraction**

The studies were selected following the Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) Statement (18). Two authors examined the eligibility of relevant studies separately, based on the consecutive examination of the titles, abstracts, and full texts. The results were then discussed in committee, and disagreements were resolved by discussion. The following information was extracted from the selected studies: country, geographic region, design, recruitment setting, ID characteristics (i.e., sample size, sex ratio, age groups [children and/or adolescents], ID levels), TD comparison sample (i.e., yes-no, sample size), height and body weight measurement method (i.e., direct [by the research team or collected via measures taken by clinicians or teachers], or indirect), type of norm used to define overweight and obesity (e.g., International Obesity Task Force [IOTF], national norms, non-national norms) (see Table 1).

### **Quality Assessment of the Reviewed Studies**

The quality of the reviewed studies was rated using criteria developed based on the recommendations of the STrengthening the Reporting of OBservational studies in Epidemiology statement (STROBE; Vandenbroucke et al., 19). These criteria covered four potential sources of bias in each study. The first criterion assessed the study population biases and more specifically whether the reviewed study: (a) was population-based; (b) reported the main demographic characteristics of the children and adolescents with ID (i.e., age, sex, and ID level); and (c) reported additional diagnoses of the children and adolescents with ID (e.g., Down syndrome, autism). The second criterion assessed outcome biases, more specifically whether the outcomes (weight and height) were measured directly (i.e., by the research team, clinicians or teachers). The third criterion assessed analysis biases, more specifically whether the reviewed studies reported subgroup (e.g., age, sex, ID levels, additional diagnoses) and/or interaction analyses. Finally, the last criterion assessed data presentation biases, more precisely whether the reviewed studies reported the frequency and prevalence estimates of overweight/obesity.

### **Statistical Analysis**

Analyses were performed with the Comprehensive Meta-Analysis (20) software (version 2.2.064). Pooled estimates of overweight, overweight-obesity and obesity prevalence were generated using the random effects model, since the studies differed greatly regarding the participants' characteristics and the assessment methods. Forest plots for prevalence were generated using spreadsheets developed by Neyeloff, Fuchs, and Moreira (21). To compare the

risk of being overweight, overweight-obese or obese between children and adolescents with ID and their TD peers, odds ratios (OR) and 95% confidence intervals were calculated.

Potential sources of heterogeneity in prevalence estimates of overweight, overweight-obesity, and obesity were examined by performing a series of pre-specified subgroup (using a mixed effect model) analyses for the following variables: (a) sex; (b) geographic regions as defined by the World Health Organization (e.g., Europe, North America, South America, Western Pacific), except for the worldwide study of Lloyd et al. (22) in which geographic regions of specific subsamples were already defined; (c) recruitment settings (e.g., regular school, special school, Special Olympics); (d) ID levels (e.g., mild, mild to moderate, mild to profound, moderate to severe-profound); (e) type of norms used to define overweight or obesity (IOTF, national, non-national); (f) method used for measuring body height and weight (direct versus indirect); and (g) additional diagnoses (e.g., Down syndrome). No moderation analysis was performed when only one study was available in a pre-specified subgroup.

Heterogeneity of prevalence estimates within and between subgroups was assessed using the  $Q$  test (23) and the  $I^2$  statistic (24). Finally, potential publication bias was assessed by examining the funnel plots [including results from the Duval and Tweedie's (25) "trim and fill" test], Begg and Mazumdar's (B-M) rank correlation test (26) and Egger's test of intercept (27).

## Results

### Study Selection

The search identified a total of 1 727 possibly eligible articles (see Figure 1). This number fell to 769 after duplicates were removed. Based on titles and abstracts, 711 studies were excluded for reasons detailed in Figure 1. The full texts of the remaining 58 articles were screened, and 16 studies (22,28-42) published between 1985 and 2015 met the inclusion criteria and were included in this meta-analysis (see Table 1).

### Study Characteristics

Participant characteristics, study design and recruitment settings for the 16 retained studies are reported in Table 1. Studies were conducted mainly in Western Pacific ( $n = 4$ ), Europe ( $n = 4$ ), and North America ( $n = 4$ ). Two of the 16 studies (13%) included a TD sample that could be used for comparative analyses. Overall, a total of 36 345 participants with ID were involved in these studies ( $M = 2\ 272$ ; range = 25 to 20 031) and nearly half of the studies recruited their participants in regular and/or special schools (10/16 studies, 63%). Additionally, 10 of the 16 studies (63%) focused on children ( $\geq 4 - 11$  years) and 14 of the 16 (88%) on adolescents ( $\geq 11 - 18$  years). Additionally, participants were mostly boys ( $M = 61\%$ ,  $SD = 5\%$ , range = 53% to 71%).

In the vast majority of the study, height and body weight were measured directly either by the research team or collected via measures taken by clinicians (e.g., nurses, doctors, dieticians) or teachers ( $N = 12$ ; 75%). In the other studies, height and weight were reported by parents, career or schools without information on the method of measurement. Overweight, overweight-obesity and obesity status were determined using BMI in all studies, except for two (33, 41) using the WLI. The vast majority of the norms used in the reviewed studies were national ( $N = 8$ ; 50%) or from the IOTF ( $N = 5$ ; 31%). Only one used non-national norms and two do not mention the norms used.

In the reviewed studies using national and non-national norms, participants were considered as being (a) overweight if their age- and sex-specific BMI was  $\geq 85^{\text{th}}$  percentile and  $< 95^{\text{th}}$  percentile or if their WLI was  $\geq 110\%$  and  $\leq 119$ ; (b) overweight-obese if their age- and sex-specific BMI was  $\geq 85^{\text{th}}$  percentile or if their WLI was  $\geq 110\%$ ; and (c) obese if their age- and sex-specific BMI was  $\geq 95^{\text{th}}$  percentile or if their WLI was  $\geq 120\%$ .

In the studies using the IOTF norms (43), participants were considered as being (a) overweight if their age- and sex-specific BMI was equivalent to an adult BMI  $\geq 25$  kg/m<sup>2</sup> and  $< 30$  kg/m<sup>2</sup>; (b) overweight-obese if their age- and sex-specific BMI was equivalent to an adult BMI  $\geq 25$  kg/m<sup>2</sup>; and (c) obese if their age- and sex-specific BMI was equivalent to an adult BMI  $\geq 30$  kg/m<sup>2</sup>.

### Prevalence Estimates of Overweight, Overweight-Obesity, and Obesity

**Children.** *Overweight* (Figure 2a) prevalence estimates of were reported in six studies from several countries (Brazil, France, South Korea, United States of America [USA], Worldwide). The pooled prevalence estimate was 15% (95%CI = 10%–20%), with a high level of heterogeneity ( $Q_{(5)} = 42, p < .001; I^2 = 88$ ). The highest estimate was observed in the USA (25%) by Foley et al. (32), and the lowest in the USA (7%) and South Korea (8%) by Fox et al. (33) and Choi et al. (31), respectively. Finally, no evidence of publication bias was noted (Figure S1a, online supplements; B-M's test,  $p = .50$ ; Egger's test,  $p = .36$ ).

Prevalence estimates of *overweight-obesity* (Figure 3a) were reported in seven studies from several countries (Brazil, France, South Korea, USA, Worldwide). The pooled prevalence estimate was 30% (95%CI = 22%–39%), with a high level of heterogeneity ( $Q_{(6)} = 82, p < .001; I^2 = 93$ ). The highest estimate was observed in the USA (45%) by Foley et al. (32) and the lowest in South Korea (14%) by Choi et al. (31). Finally, no evidence of publication bias was noted by the B-M rank correlation test ( $p = .50$ ) and Egger's test of intercept ( $p = .46$ ). The Duvall and Tweedie's trim and fill revealed that two studies were missing on the left of the funnel plot (Figure S1b, online supplements). When these two studies are imputed to obtain a symmetrical funnel plot, the pooled prevalence estimate becomes 27% (95%CI = 20%–35%).

*Obesity* prevalence estimates (Figure 4a) were reported in nine studies from several countries (Egypt, Brazil, France, Japan, South Korea, Taiwan, USA, Worldwide). The pooled prevalence estimate was 13% (95%CI = 10%–16%), with a high level of heterogeneity ( $Q_{(8)} = 61, p < .001; I^2 = 87$ ). The highest estimate was observed in the USA (21%) by Foley et al. (32) and Fox et al. (33), and the lowest in South Korea (6%) by Choi et al. (31). Finally, no evidence of publication bias was noted by the B-M rank correlation test ( $p = .38$ ) and Egger's test of intercept ( $p = .30$ ). The Duvall and Tweedie's trim and fill revealed that two studies were missing on the left of the funnel plot (Figure S1c, online supplements). When these two studies are imputed to obtain a symmetrical funnel plot, the pooled prevalence estimate becomes 11% (95%CI = 9%–14%).

**Adolescents.** *Overweight* (Figure 2b) prevalence estimates were reported in eight studies from several countries (Australia, France, South Korea, USA, Worldwide). The pooled prevalence estimate was 18% (95%CI = 16%–21%), with a high level of heterogeneity ( $Q_{(7)} = 55, p < .001; I^2 = 87$ ). The highest prevalence estimate was observed in Australia (24%) and the USA (23%) by Krause et al. (34), and Foley et al. (32), respectively. Conversely, the lowest prevalence estimates were observed in France (13%) and South Korea (13%) by Mikulovic et al. (36) and Choi et al. (31), respectively. Finally, no evidence of publication bias was noted (Figure S2a, online supplements; B-M's test,  $p = .19$ ; Egger's test,  $p = .47$ ).

Prevalence estimates of *overweight-obesity* (Figure 3b) were reported in nine studies (Australia, France, South Korea, USA, Worldwide). The pooled prevalence estimate was 33% (95%CI = 27%–39%), with a high level of heterogeneity ( $Q_{(8)} = 235, p < .001; I^2 = 97$ ). The highest prevalence estimate was observed in the USA (51%) by Foley et al. (32) and the lowest in France (15%) by Mikulovic et al. (36). Finally, no evidence of publication bias was noted (Figure S2b, online supplements; B-M's test,  $p = .46$ ; Egger's test,  $p = .44$ ).

*Obesity* prevalence estimates (Figure 4b) were reported in 13 studies from several countries (Egypt, France, Indonesia, Japan, South Korea, Taiwan, USA, Worldwide). The pooled prevalence estimate was 15% (95%CI = 13%–18%) with a high level of heterogeneity ( $Q_{(12)}=230, p < .001; I^2 = 95$ ). The highest prevalence estimate was observed in Turkey (28%) and in the USA (28%) by Nogay (37) and Foley et al. (32), respectively. Inversely, the lowest prevalence estimate was observed in France (2%) by Mikulovic et al. (36). Finally, no evidence of publication bias was noted by the B-M rank correlation test ( $p = .21$ ) and Egger's test of intercept ( $p = .23$ ). The Duvall and Tweedie's trim and fill revealed that three studies were missing on the left of the funnel plot (Figure S2c, online supplements). When these three studies are imputed to obtain a symmetrical funnel plot, the pooled prevalence estimate was 13% (95%CI = 11%–16%).

### **Moderation Analyses**

Results from the moderation analyses, as well as the references of the studies used for these analyses are detailed in Tables S1-S6 in the online supplements.

**Children.** Findings showed significant disparities in overweight-obesity and obesity pooled prevalence estimates by geographic region (Tables S4-S6). Pairwise comparisons (available upon request from the first author) showed that the risk of being overweight-obese was greater in children with ID from North America (39%) than in those living in South America (25%). North American children with ID were also more likely to be obese (23%) than those from Europe (9%), South America (11%), and Western Pacific (9%). Results also showed significant variations in the pooled prevalence of obesity by recruitment setting (Tables S5-S6). The pooled prevalence of obesity were significantly higher in children recruited via Special Olympics (17%) than via special schools (9%). No significant variations were found as a function of sex, ID level, additional diagnoses, body-height and body-weight measurement method or the norms used to define overweight, overweight-obesity or obesity (Tables S1-S6, online supplements).

**Adolescents.** Findings showed significant disparities in overweight, overweight-obesity and obesity pooled prevalence estimates by geographic region (Tables S1-S6). Adolescents with ID from North America (22% and 48%) were more likely to be overweight and overweight-obese than those from Europe (17% and 23%), and more likely to be obese (27%) than those living in Europe (8%), South East Asia (10%), and Western Pacific (16%). Results (Tables S4-S6) also showed that pooled prevalence of overweight-obesity and obesity were significantly higher in adolescents with Down syndrome (61% and 33%) than without Down syndrome (35% and 17%). Findings also revealed higher pooled prevalence of obesity among adolescents with ID in studies using national norms (19%) rather IOTF norms (10%). No significant variations were found as a function of sex, ID level, or body-height and body-weight measurement method (Tables S1-S6, online supplements).

### **Comparison of Risk with Typically Developing Peers**

**Children.** None of the studies assessed the risk of being overweight, overweight-obese or obese in children with ID compared with their TD peers.

**Adolescents.** Only two studies provided data that can be used to compare the risk of overweight/obesity in adolescents with ID and their TD peers (34,39). Random effects models showed a statistically higher risk for overweight-obesity and obesity in adolescents with ID than in their TD peers, with a pooled OR of 1.54 (95%CI = 1.12–2.12,  $p = 0.008$ ) and of 1.80 (95%CI = 1.30–2.49,  $p < 0.001$ ); with a moderate level of heterogeneity ( $Q_{(1)} = 1.3, p = 0.26; I^2 = 39$ ) and no observed heterogeneity ( $Q_{(1)} = 0.11, p = 0.74; I^2 = 0$ ), respectively. Nevertheless, no significant differences were found in the risk of being overweight (OR = 1.15; 95%CI = 0.80–1.63;  $p = 0.46$ ), with a very low level of heterogeneity ( $Q_{(1)} = 1.1, p = 0.31; I^2 = 5$ ).



### Quality Rating of the Reviewed Studies

Table 2 provides the quality ratings of the reviewed studies based on STROBE's criteria (19). The fact that only three studies (19%) used a population-based sampling raises concerns about the representativeness of the other samples and, consequently, about the generalizability of their results (Table 2), thus supporting the importance of the quantitative review process conducted here. Additionally, information regarding sample characteristics was often lacking, especially concerning sex ratios (5/16 studies, 31%), ID levels (12/16 studies, 75%), and additional diagnoses (7/16 studies, 44%). The majority of the studies (75%) have measured height and weight directly. Only one study did not report subgroup analyses, but only four (25%) performed interaction analyses. Finally, only three studies did not report both frequency and percentage of overweight or obese children and adolescents with ID (Table 2).

### Discussion

#### Prevalence Estimates of Overweight, Overweight-Obesity, and Obesity

The first objective of this meta-analysis was to determine the pooled prevalence estimates of overweight/obesity among children and adolescents with ID. Among children, the results revealed pooled prevalence estimates of overweight, overweight-obesity, and obesity of 15% (7%–25%), 30% (14%–45%), and 13% (6%–21%), respectively. These pooled prevalence estimates are higher than those found in previous studies of TD children (44,45). In addition, these findings show that the highest and lowest prevalence estimates of overweight/obesity were observed in studies from the USA (32) and South Korea (31), respectively. Interestingly, findings from a French study (30) show that children with ID were two times more likely to be overweight (22%) than obese (10%), whereas in an American study (33) the children with ID were three times more likely to be obese (21%) than overweight (7%). However, in Brazil (29) and South Korea (31) prevalence estimates were nearly similar for overweight and obesity.

Among adolescents, the results revealed pooled prevalence estimates of overweight, overweight-obesity, and obesity of 18% (13%–24%), 33% (15%–51%), and 15% (2%–28%), respectively. These pooled prevalence estimates are higher than those found among TD adolescents (46). Additionally, these findings show that (a) the highest prevalence estimates of overweight were observed in studies from Australia (34) and the USA (32), and of obesity in studies from the USA (32) and Turkey (37); (b) the lowest prevalence estimates of overweight were observed in studies from France (36) and South Korea (31), and of obesity in studies from France (30,36). The highest prevalence estimates of overweight-obesity and obesity (> 20%) were found in three studies from the USA (32,38,39). Finally, two French studies (30,36) show that adolescents with ID were nearly three (19% vs. 7%) to seven (13% vs. 2%) times more likely to be overweight than obese. Inversely, studies from the USA (32,38,39) show that adolescents with ID were slightly more likely to be obese (20%–28%) than overweight (16%–23%). However, similar rates of overweight and obesity were observed in Australia (34) and South Korea (31).

#### Moderators

The second objective of this meta-analysis was to examine whether the heterogeneity of overweight/obesity prevalence estimates across studies could be attributed to various moderators, including the participants' characteristics (i.e., sex, geographic region, recruitment setting, ID level, and additional diagnoses) or the assessment method (i.e., type of norms used to define overweight or obesity, body-height and body-weight measurement method).

**Participants' characteristics.** The findings suggest that the risk of overweight was higher in adolescents with ID living in North America than in Europe. Additionally, findings show that children and adolescents with ID living in North America were significantly more overweight-obese than those living in South America and in Europe. Finally, children and

adolescents with ID living in North America were significantly more obese than those living in Europe, South America and in the Western Pacific. This finding suggests that the North-American environment may expose youth with ID to be at greater risk for weight gain when compared to other geographic areas. These results are consistent with those from recent reviews (44-46) and cross-national studies (47,48) conducted among TD youth. However, our findings should be interpreted with caution. Indeed, even within regions, most of the participants were from the same country (e.g., USA, France). Consequently, future research examining the prevalence of overweight/obesity among youth with ID should focus on more regions, as well as countries within regions.

Surprisingly, subsequent analyses suggest that children with ID recruited via Special Olympics were significantly more obese than those recruited in special schools. Higher prevalence estimates of obesity were also observed in adolescents recruited via Special Olympics compared to special schools, or regular and special schools, but the difference was non-significant. This result might be related to the fact that athletes with ID attending Special Olympics may have been misclassified as obese by reporting increased muscular mass, associated to a higher BMI. Further studies are required to determine the actual reasons of these counter-intuitive results.

Additional findings suggest that adolescents with Down syndrome were significantly more overweight-obese and obese than those without Down syndrome. This result is consistent with findings from previous reviews (4,10,11,13), and may be explained by Must et al.'s (13) observation that adolescents with Down syndrome have "lower fat-free mass (González-Agüero et al., 2011) and lower resting metabolic rates than typically developing children (Hill et al. 2013; Luke et al., 1996)" (p. 158). Additionally, according to Reinehr et al. (4, p. 270), adolescents with Down syndrome "have a predisposition to overeat, since the cerebral regions that are responsible for weight regulation (hypothalamus) may be damaged (van Mil et al., 2001; Luke et al., 1996)." Nevertheless, since only two studies (34,39) were included in this subgroup analysis, these differences should be interpreted with caution and require additional scientific attention. Finally, no significant variations were found in the pooled prevalence estimates for other characteristics of participants with ID, including sex and ID level.

**Assessment methods.** These findings show that the norms used to define obesity significantly influenced prevalence estimates among adolescents with ID. Indeed, findings show that studies relying on national norms provided significantly higher pooled prevalence estimates of obesity than those using the IOTF norms, which is consistent with data obtained in TD youth (49). It is thus important that future studies systematically provided prevalence estimates of overweight/obesity among youth with ID using both norms (national and IOTF) in order to enable international and national comparisons. Finally, no significant variations were found in the pooled prevalence estimates for assessment methods, including body-height and body-weight measurement method.

### **Comparison with TD peers**

The last objective of this meta-analysis was to examine whether children and adolescents with ID were at greater risk of overweight/obesity than their TD peers. The findings show that the risk of being overweight was nearly identical between adolescents with ID and their TD counterparts. However, additional findings indicate that adolescents with ID were nearly two times more at risk of being overweight-obese and obese than their TD peers. This result is consistent with findings from previous reviews (11,13). It shows that obesity represents an important health threat for this population that deserves more attention from practitioners and policy makers. Nevertheless, since only two studies (38,39) from the USA were included in these

analyses, these differences should be interpreted with caution and require future research.

### **Limitations and Directions for Future Studies**

Although informative, the findings from this systematic review and meta-analysis should be interpreted with caution given the limitations of the reviewed studies. First, the reviewed studies were conducted mostly in Europe, North America, and Western Pacific regions. Moreover, only a few considered participants' characteristics (e.g., sex, age, ID level, living arrangements) and additional diagnoses (e.g., autism spectrum disorder, Down syndrome, Fragile X, Prader-Willi syndrome, physical disabilities) in subgroup analyses. Therefore, the moderating role of these variables should be more thoroughly examined in future studies.

Second, as already illustrated in previous reviews (8-11,13), most of the risk factors commonly associated with overweight/obesity in the general population (e.g., dietary intake, physical activity, sedentary behavior, socioeconomic status) were insufficiently controlled in the reviewed studies. Consequently, the role of these factors could not be examined in the present meta-analysis and should be examined in future studies.

Third, only a few of the reviewed studies compared the risk of overweight/obesity between children and adolescents with ID and their TD peers, and more specifically while controlling for key participant characteristics, such as sex and age. Consequently, it is unknown whether boys and girls or early and late children or adolescents with ID are at greater risk of overweight/obesity than their TD peers. Clearly, this issue should be examined in future studies.

Fourth, all the reviewed studies (except 28,34,36, 37) recruited participants in a single setting, and all used only one criterion (i.e., national norms or IOTF) to define overweight or obesity. Consequently, the role of these potential moderators in the prevalence estimates of overweight and obesity should be more thoroughly investigated in future studies.

Fifth, none of the reviewed studies (except 32) relied on a longitudinal design, precluding examination of longitudinal trajectories of overweight or obesity prevalence among youth with ID. Consequently, it is still unknown (a) whether overweight or obesity prevalence is plateauing or increasing with time, and (b) whether trajectories of overweight or obesity prevalence could differ according to the participants' characteristics (e.g., sex, age, ID level, additional diagnoses). A longitudinal design study is needed to provide clear answers to these questions.

Finally, the reviewed studies were generally poorly described and lacking in details on key variables, such as sex ratio, age or ID level. Last but not least, as regards descriptive epidemiology, only three studies were population-based, designed using a national survey database and/or a random-sampling method (34,35,38). This raises serious concerns about the representativeness of most of the samples, and thus the value of the reported prevalence rates.

### **Conclusion**

This meta-analysis highlights that a large proportion of children and adolescents with ID are overweight and obese, and that they are significantly more obese than their TD peers. Unfortunately, many potentially important determinants of overweight and obesity (e.g., sex, age, ID level, living arrangements, additional diagnoses, dietary intake, physical activity) are clearly understudied in this population and deserve further investigation. Additionally, such a high prevalence among children and adolescents with ID is worrisome and problematic given their known higher risk of developing secondary health problems (52). A key policy priority should thus be to develop and test specific age subgroups lifestyle intervention for tackling or managing this serious public health issue in this vulnerable population (for systematic reviews see 53,54).

### **References**

1. Allison DB, Packer-Munter W, Pietrobelli A, Alfonso VC, Faith MS. Obesity and developmental disabilities: Pathogenesis and treatment. *J Dev Phys Disabil* 1998; **10/3**: 215-255.

2. Ells LJ, Lang R, Shield JPH *et al.* Obesity and disability—a short review. *Obes Rev* 2006; **7/4**: 341-345.
3. Holcomb MJ, Puffaff LA, McIntosh DE. Obesity rates in special populations of children and potential interventions. *Psychol Schools* 2009; **46/8**: 797-804.
4. Reinehr T, Dobe M, Winkel K, Schaefer A, Hoffmann D. Obesity in Disabled Children and Adolescents: An overlooked group of patients. *Dtsch Arztebl Int* 2010; **107/15**: 268.
5. Minihan PM, Fitch SN, Must A. What does the epidemic of childhood obesity mean for children with special health care needs? *J Law Med Ethics* 2007; **35/1**: 61-77.
6. Rimmer JH, Rowland JL, Yamaki K. Obesity and secondary conditions in adolescents with disabilities: Addressing the needs of an underserved population. *J Adol Health* 2007; **41**: 224–229.
7. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 5<sup>th</sup> ed. Arlington, VA: American Psychiatric Association, 2013.
8. Mañano C. Prevalence and risk factors of overweight and obesity among children and adolescents with intellectual disabilities. *Obes Rev* 2011; **12**: 189-197.
9. Bandini L, Danielson M, Esposito LE *et al.* Obesity in children with developmental and/or physical disabilities. *Disabil Health J* 2015; **8/3**: 309-316.
10. Casey AF, Rasmussen R. Obesity. In: Matson JL, Matson ML (eds.). *Comorbid conditions in individuals with intellectual disabilities* [Autism and child psychopathology series]. Cham, Switzerland: Springer, 2015, pp. 275-298.
11. Grondhuis SN, Aman MG. Overweight and obesity in youth with developmental disabilities: A call to action. *J Intellect Disabil Res* 2013; **58/9**: 787-799.
12. McGillivray J, McVilly K, Skouteris H, Boganin C. Parental factors associated with obesity in children with disability: A systematic review. *Obes Rev* 2013; **14/7**: 541-554.
13. Must A, Curtin C, Hubbard K, Sikich L, Bedford J, Bandini L. Obesity prevention for children with developmental disabilities. *Curr Obes Rep* 2014; **3/2**: 156-170.
14. Stroup DF, Berlin JA, Morton SC *et al.* Meta-analysis of observational studies in epidemiology: A proposal for reporting. *J Am Med Assoc* 2000; **283/15**: 2008-2012.
15. Cole TJ. A method for assessing age-standardized weight-for-height in children seen cross-sectionally. *Ann Hum Biol* 1979; **6**: 249-68.
16. DuRant RH, Linder CW. An evaluation of five indexes of relative body weight for use with children. *J Am Diet Assoc* 1981; **78/1**: 35-41.
17. Casey AF. Measuring body composition in individuals with intellectual disability: A scoping review. *J Obes*; 2013.
18. Liberati A, Altman DG, Tetzlaff J *et al.* The PRISMA Statement for reporting systematic reviews and meta-analysis of studies that evaluate health care interventions: Explanation and elaboration. *PLoS Med* 2009; **6**: 1-6.
19. Vandembroucke JP, von Elm EV, Altman DG *et al.* Strengthening the reporting of observational studies in epidemiology (STROBE): Explanation and elaboration. *PLoS Med* 2007; **4/10**: e297.
20. Borenstein M, Hedges L, Higgins J, Rothstein H (eds) *Comprehensive meta-analysis version 2*. Englewood, NJ: Biostat, 2005.
21. Neyeloff JL, Fuchs SC, Moreira LB. Meta-analyses and Forest plots using a microsoft excel spreadsheet: Step-by-step guide focusing on descriptive data analysis. *BMC Res Notes* 2012; **5/1**: 52.
22. Lloyd M, Temple VA, Foley JT. International BMI comparison of children and youth with intellectual disabilities participating in Special Olympics. *Res Dev Disabil* 2012; **33**: 1708-1714.

23. Cochran WG. The comparison of percentages in matched samples. *Biometrika* 1950; **37**: 256-66.
24. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *Br Med J* 2003; **327/7414**: 557.
25. Duval S, Tweedie R. Trim and fill: A simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics* 2000; **56/2**: 455-463.
26. Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. *Biometrics* 1994; **50/4**: 1088-1101.
27. Egger M, Smith GD, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *Br Med J* 1997; **315/7109**: 629-634.
28. AbdAllah AM, El-Sherbeny SS, Khairy, S. Nutritional status of mentally disabled children in Egypt. *Egypt J of Hosp Med* 2007; **29**: 604-615.
29. Batista LRV, Moreira EAM, Rauen MS, Corso ACT, Fiates GMR. Oral health and nutritional status of semi-institutionalized persons with mental retardation in Brazil. *Res Dev Disabil* 2009; **30/5**: 839-846.
30. Bégarie J, Maïano C, Leconte P, Ninot G. The prevalence and determinants of overweight and obesity among French youths and adults with intellectual disabilities attending special education schools. *Res Dev Disabil* 2013; **34/5**: 1417-1425.
31. Choi E, Park H, Ha Y, Hwang WJ. Prevalence of overweight and obesity in children with intellectual disabilities in Korea. *J Appl Res Intellect Disabil* 2012; **25/5**: 476-483.
32. Foley JT, Lloyd M, Vogl D, Temple VA. Obesity trends of 8–18 year old Special Olympians: 2005–2010. *Res Dev Disabil* 2014; **35/3**: 705-710.
33. Fox R, Hartney CW, Rotatori AF, Kurpiers EM. Incidence of obesity among retarded children. *Educ Train Ment Retard* 1985; **20/3**: 175-181.
34. Krause S, Ware R, McPherson L, Lennox N, O’Callaghan M. Obesity in adolescents with intellectual disability: Prevalence and associated characteristics. *Obes Res Clin Pract* 2015. Advance online publication. doi: 10.1016/j.orcp.2015.10.006.
35. Lin JD, Yen CF, Li CW, Wu JL. Patterns of obesity among children and adolescents with intellectual disabilities in Taiwan. *J Appl Res Intellect Disabil* 2005; **18/2**: 123-129.
36. Mikulovic J, Marcellini A, Compte R *et al.* Prevalence of overweight in adolescents with intellectual deficiency. Differences in socio-educative context, physical activity and dietary habits. *Appetite* 2011; **56**: 403–407.
37. Nogay NH. Nutritional status in mentally disabled children and adolescents: A study from Western Turkey. *Pak J Med Sci* 2013; **29/2**: 614-617.
38. Phillips KL, Schieve LA, Visser S *et al.* Prevalence and impact of unhealthy weight in a national sample of US adolescents with autism and other learning and behavioral disabilities. *Matern Child Health J* 2014; **18/8**: 1964-1975.
39. Rimmer JH, Yamaki K, Lowry BMD, Wang E, Vogel LC. Obesity and obesity-related secondary conditions in adolescents with intellectual/developmental disabilities. *J Intellect Disabil Res* 2010; **54**: 787-794.
40. Salaun L, Berthouze-Aranda S. Obesity in school children with intellectual disabilities in France. *J Appl Res Intellect Disabil* 2011; **24**: 333-340.
41. Takeuchi E. Incidence of obesity among school children with mental retardation in Japan. *Am J Ment Retard* 1994; **99/3**: 283-288.
42. Tamin TZ, Idris FH, Mansyur M, Syarif DR. Prevalence and determinants of obesity in students with intellectual disability in Jakarta. *Med J Indones* 2014; **23/2**: 106-111.

43. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: International survey. *Br Med J* 2000; 320:1240-1243.
44. Wang Y, Lobstein TIM. Worldwide trends in childhood overweight and obesity. *Int J Pediatr Obes* 2006; **1/1** : 11-25.
45. Wang Y, Lim H. The global childhood obesity epidemic and the association between socio-economic status and childhood obesity. *Int Rev Psychiatry* 2012; **24/3**: 176-188.
46. Bibiloni MDM, Pons A, Tur JA. Prevalence of overweight and obesity in adolescents: A systematic review. *Int Sch Res Notices Obes* 2013; 392747.
47. Haug E, Rasmussen M, Samdal O *et al*. Overweight in school-aged children and its relationship with demographic and lifestyle factors: results from the WHO-Collaborative Health Behaviour in School-aged Children (HBSC) study. *Int J Public Health* 2009; **54/2**: 167-179.
48. Lissau I, Overpeck MD, Ruan WJ, Due P, Holstein BE, Hediger ML. Body mass index and overweight in adolescents in 13 European countries, Israel, and the United States. *Arch Pediatr Adolesc Med* 2004; **158/1**: 27-33.
49. Reilly JJ, Kelly J, Wilson DC. Accuracy of simple clinical and epidemiological definitions of childhood obesity: systematic review and evidence appraisal. *Obes Rev* 2010; **11/9**: 645-655.
50. De Onis M, Lobstein T. Defining obesity risk status in the general childhood population: which cut-offs should we use?. *Int J Pediatr Obes* 2010; **5/6**: 458-460.
51. Rolland-Cachera MF. Childhood obesity: Current definitions and recommendations for their use. *Int J Pediatr Obes* 2011; **6/5-6**: 325-331.
52. van Schrojenstein Lantman-de HM, Walsh PN. Managing health problems in people with intellectual disabilities. *Br Med J* 2008; **337**: a2507.
53. Casey AF, Rasmussen R. Reduction measures and percent body fat in individuals with intellectual disabilities: A scoping review. *Disabil Health J* 2013; **6**: 2-7.
54. Maïano C, Normand CL, Aimé A, Bégarie J. Lifestyle interventions targeting changes in body weight and composition among youth with an intellectual disability: A systematic review. *Res Dev Disabil* 2014; **35/8**: 1914-1926.





Figure 1. Results of Search Based on the PRISMA Statement (18)

(a)



(b)





Figure 2. Forest Plot of Random-Effects Pooled Prevalence Estimates of *Overweight* among (a) Children and (b) Adolescents



(a)



(b)





Figure 3. Forest Plot of Random-Effects Pooled Prevalence Estimates of *Overweight-Obesity* among (a) Children and (b) Adolescents

(a)



(b)




Figure 4. Forest Plot of Random-Effects Pooled Prevalence Estimates of Obesity among (a) Children and (b) Adolescents

Table 1

*Main Characteristics of the Studies Included in the Meta-Analysis*

Study	Country	Geographic Region	Design	Recruitment Setting	ID				TD		Measurement method	Norms used to define OW or OB	Outcome
					Sample size (N)	Sex (% of boys)	Age group (age range in years)	ID level	Yes-No	Sample size (N)			
Abdallah et al. (28)	Egypt	Eastern Mediterranean	CS	Regular and special schools	574	NA	Children (9-10) & Adolescents (11-14)	NA	O	Direct	NA	OB	
Batista et al. (29)	Brazil	South America	CS	ID association	54	NA	Children (5-11)	NA	O	Direct	Non-national	OW, OW-OB, OB	
Bégarie et al. (30)	France	Europe	CS	Special school	865	NA	Children (5-10) & Adolescents (11-18)	Mild to severe	O	Direct	IOTF	OW, OW-OB, OB	
Choi et al. (31)	South Korea	Western Pacific	CS	Special school	2404	71	Children (7-10) & Adolescents (11-18)	NA	O	Direct	National	OW, OW-OB, OB	
Foley et al. (32)	USA	North America	Cohort	Special Olympics	1122	61	Children (8-11) & Adolescents (12-18)	NA	O	Direct	National	OW, OW-OB, OB	
Fox et al. (33)	USA	North America	CS	Regular school	118	53	Children (5.5-10.4)	Mild to profound	O	Direct	National	OW, OW-OB, OB	
Krause et al. (34)	Australia	Western Pacific	CS	Regular and special schools	261	56	Adolescents (13-18)	NA	O	Direct	IOTF	OW, OW-OB, OB	
Lin et al. (35)	Taiwan	Western Pacific	CS	National survey	187	64	Children (4-6) & Adolescents (13-18)	Mild to profound	O	Indirect	National	OB	
Lloyd et al. (22)	Worldwide	North America, Africa-Eastern Mediterranean, South America, Europe, South-East Asia, Asia-Western Pacific	CS	Special Olympics	9678	63	Children (8-11) & Adolescents (12-18)	NA	O	Direct	IOTF	OW, OW-OB, OB	
Mikulovic et al. (36)	France	Europe	CS	Regular and special schools	183	NA	Adolescents (11-14)	NA	O	Direct	IOTF	OW, OW-OB, OB	
Nogay (37)	Turkey	Europe	CS	Regular and special schools	25	60	Adolescents (14-18)	Mild to severe	O	Direct	National	OB	
Phillips et al. (38)	USA	North America	CC	National survey	60	59	Adolescents (12-17)	NA	●	8141	Indirect	National	OW, OW-OB, OB
Rimmer et al. (39)	USA	North America	CC	ID association	163	68	Adolescents (12-18)	NA	●	12973	Indirect	National	OW, OW-OB, OB
Salaun & Berthouze-Aranda (40)	France	Europe	CS	Special school	192	56	Children (6-11) & Adolescents (12-18)	Mild	O	Direct	IOTF	OW-OB	
Takeuchi (41)	Japan	Western Pacific	CS	Special school	20031	65	Children (6-11) & Adolescents (12-17)	NA	O	Indirect	National	OB	
Tamin et al. (42)	Indonesia	South-East Asia	CS	Special school	428	NA	Adolescents (14-16)	NA	O	Direct	NA	OB	

Note. CC= case control; CS= cross-sectional; ID= intellectual disability; OB= obesity; OW= overweight; OW-OB= overweight including obesity; USA= United States of America; IOTF= International Obesity Task Force; ●= Yes; ○= No; NA= not available; TD= typically developing.

*Quality Assessment of Studies*

Study	Population bias			Outcome bias	Analysis bias		Data presentation bias
	Population-based	Main demographic characteristics reported	Additional diagnosis reported	Direct measurement of height and weight	Subgroup analyses reported	Interaction analyses reported	Frequency and prevalence of outcomes reported
Abdallah et al. (28)	○	○ (ID level)	○	●	● (age, SES, sex)	○	●
Batista et al. (29)	○	○ (sex, ID level)	○	●	● (age)	○	●
Bégarie et al. (30)	○	●	●	●	● (age, sex, comorbidity, ID level, etc.)	●	●
Choi et al. (31)	○	○ (ID level)	●	●	● (age, sex)	●	●
Foley et al. (32)	○	○ (sex, ID level)	○	●	● (age, sex)	○	○
Fox et al. (33)	○	●	●	●	● (age, sex, ID level, comorbidity, etc.)	●	●
Krause et al. (34)	●	○ (ID level)	●	●	● (age, sex, comorbidity, behaviour problems, mobility, medication, etc.)	●	●
Lin et al. (35)	●	○ (ID level)	●	○	● (age, sex)	○	●
Lloyd et al. (22)	○	○ (ID level)	○	●	● (age, sex)	○	○
Mikulovic et al. (36)	○	○ (ID level)	○	●	● (age, sex, school type)	○	○
Nogay (37)	○	●	○	●	● (age, sex)	○	●
Phillips et al. (38)	●	○ (sex, ID level)	●	○	○	○	●
Rimmer et al. (39)	○	○ (age, sex, ID level)	●	○	● (diagnosis)	○	●
Salaun & Berthouze-Aranda (40)	○	●	●	●	● (age, sex)	○	●
Takeuchi (41)	○	○ (ID level)	●	○	● (age, sex)	○	●
Tamin et al. (42)	○	○ (sex, ID level)	○	●	● (age)	○	●

Note. ● (reported characteristics) = Yes; ○ (missing characteristics) = No; ID = intellectual disability; SES = socioeconomic status

**Online Supplemental Materials for:  
Prevalence of Overweight and Obesity Among Children and Adolescents with Intellectual  
Disabilities: A Systematic Review and Meta-Analysis**

This online supplement comprises nine sections including:

S1. References for Full-Text Articles Assessed for Eligibility but Excluded from the Meta-Analysis

Table S1. Subgroup Analyses of Prevalence Estimates of *Overweight* for Children and Adolescents

Table S2. Studies Included in the Subgroup Analyses of Prevalence Estimates of *Overweight* for Children and Adolescents

Table S3. Subgroup Analyses of Prevalence Estimates of *Overweight-Obesity* for Children and Adolescents

Table S4. Studies Included in the Subgroup Analyses of Prevalence Estimates of *Overweight-Obesity* for Children and Adolescents

Table S5. Subgroup Analyses of Prevalence Estimates of *Obesity* for Children and Adolescents

Table S6. Studies Included in the Subgroup Analyses of the Prevalence Estimates of *Obesity* for Children and Adolescents

Figure S1. Funnel Plot of Standard Error by Logit Event Rate of (a) Overweight, (b) Overweight-Obesity, and (c) Obesity among *Children*

Figure S2. Funnel Plot of Standard Error by Logit Event Rate of (a) Overweight, (b) Overweight-Obesity, and (c) Obesity among *Adolescents*

**Section 1. References for Full-Text Articles Assessed for Eligibility but Excluded from the Meta-Analysis**

- Abass AO, 'Dele O. Body composition characteristics of primary school deaf and mentally retarded pupils in Nigeria. *Eur J Scient Res* 2010; **40/3**: 423-427.
- Aldridge J, Pullen J, Whelan C. A survey of the nutritional status of adults and children using the learning disability services of a Health Trust. *J Intellect Disabil* 2000; **4/2**: 141-152.
- Ayoob KT, Kaminer RK, Zawel J. Chronic overeating without obesity in children with developmental disabilities: Description of a new syndrome. *Ment Retard* 1994; **32/3**: 194-199.
- Baer MT, Herman DR, Kozlowski B, Taylor M, Blyle, E, Trabms C. Weight-for-height provides a good estimate of body composition regardless of neuromotor status in children with developmental delays. *FASEB J* 1998; **12/5**: A838.
- Bandini LG, Curtin C, Hamad C, Tybor DJ, Must A. Prevalence of overweight in children with developmental disorders in the continuous National Health and Nutrition Examination Survey (NHANES) 1999-2002. *J Pediatr* 2005; **146/6**: 738-743.
- Calis EA, Veugelers R, Rieken R, Tibboel D, Evenhuis HM, Penning C. Energy intake does not correlate with nutritional state in children with severe generalized cerebral palsy and intellectual disability. *Clin Nutr* 2010; **29/5**: 617-621.
- Chowdhury SD, Ghosh T. Nutritional and socioeconomic status in cognitive development of Santal children of Purulia district, India. *Ann Hum Biol* 2011; **38/2**: 188-193.
- Culley WJ, Jolly DH, Mertz ET. Heights and weights of mentally retarded children. *Am J Ment Defic* 1963; **68**: 203-210.
- De S, Small J, Baur LA. Overweight and obesity among children with developmental disabilities. *J Intellect Dev Disabil* 2008; **33/1**: 43-47.
- Ells LJ, Shield JP, Lidstone JS *et al*. Teesside Schools Health Study: Body mass index surveillance in special needs and mainstream school children. *Public health* 2008; **122/3**: 251-254.
- Emerson E, Robertson J. Obesity in young children with intellectual disabilities or borderline intellectual functioning. *Int J Pediatr Obes* 2010; **5**: 320-326.
- Emerson E. Overweight and obesity in 3-and 5-year-old children with and without developmental delay. *Public Health* 2009; **123/2**: 130-133.
- Frey GC, Chow B. Relationship between BMI, physical fitness, and motor skills in youth with mild intellectual disabilities. *Int J Obes* 2006; **30/5**: 861-867.
- Garn SM, Weir HF. Assessing the nutritional status of the mentally retarded. *Am J Clin Nutr* 1971; **24/7**: 853-854.
- George VA, Shacter SD, Johnson, PM. BMI and attitudes and beliefs about physical activity and nutrition of parents of adolescents with intellectual disabilities. *J Intellect Disabil Res* 2011; **55/11**: 1054-1063.
- Gibson JC, Temple VA, Anholt JP, Gaul CA. Nutrition needs of young Special Olympics participants. *J Intellect Dev Disabil* 2011; **36**: 268-272.
- Ha Y, Vann JC, Choi E. Prevalence of overweight and mothers' perception of weight status of their children with intellectual disabilities in South Korea. *J Sch Nurs* 2010; **26**: 212-222.
- Harris N, Rosenberg A, Jangda S, O'Brien K, Gallagher ML. Prevalence of obesity in International Special Olympic athletes as determined by body mass index. *J Am Diet Assoc* 2003; **103**: 235-237.

- Lionti T, Reid SM, Reddihough D, Sabin MA. Monitoring height and weight: Findings from a developmental paediatric service. *J Paediatr Child Health* 2013; **49/12**: 1063-1068.
- Lloyd M, Foley JT, Temple VA. Body mass index of children and youth with an intellectual disability by country economic status. *Prev Med* 2014; **69**: 197-201.
- Maaskant MA, Knijff-Raeven V, Ankie GM, Van Schrojenstein Lantman-de Valk H, Veenstra, MY. Weight status of persons with intellectual disabilities. *J Appl Res Intellect Disabil* 2009; **22/5**: 426-432.
- Marshall D, McConkey R, Moore G. Obesity in people with intellectual disabilities: The impact of nurse-led health screenings and health promotion activities. *J Adv Nurs* 2003; **41/2**: 147-153.
- Mathur M, Bhargava R, Benipal R *et al*. Dietary habits and nutritional status in mentally retarded children and adolescents: A study from North Western India. *J Indian Assoc Child Adolesc Ment Health* 2007; **3/2**: 22-29.
- Murphy CM, Allison DB, Babbitt RL, Patterson HL. Adiposity in children: Is mental retardation a critical variable? *Int J Obes* 1992; **16/9**: 633-638.
- Neter JE, Schokker DF, de Jong E, Renders CM, Seidell JC, Visscher TL. The prevalence of overweight and obesity and its determinants in children with and without disabilities. *J Pediatr* 2011; **158/5**: 735-739.
- Pesce KA, Wodarski LA, Wang M. Nutritional status of institutionalized children and adolescents with developmental disabilities. *Res Dev Disabil* 1989; **10/1**: 33-52.
- Rimmer JH, Yamaki K, Davis BM, Wang E, Vogel LC. Obesity and overweight prevalence among adolescents with disabilities. *Prev Chronic Dis* 2011; **8**: 1-6.
- Salaun L, Berthouze-Aranda SE. Physical fitness and fatness in adolescents with intellectual disabilities. *J Appl Res Intellect Disabil* 2012; **25/3**: 231-239.
- Sánchez-Lastres J, Eirís-Puñal J, Otero-Cepeda JL, Pavón-Belinchón P, Castro-Gago M. Nutritional status of mentally retarded children in north-west Spain. I. Anthropometric indicators. *Acta Paediatr* 2003; **92/6**: 747-753.
- Sari HY, Bahceci B. Nutritional status of children with an intellectual disability. *Int J Disabil Hum Dev* 2012; **11/1**: 17-21.
- Shabayek MM. Assessment of the nutritional status of children with special needs in Alexandria. Part II: Anthropometric measures. *J Egypt Public Health Assoc* 2003; **79/5-6**: 363-382.
- Slevin E, Truesdale-Kennedy M, McConkey R, Livingstone B, Fleming P. Obesity and overweight in intellectual and non-intellectually disabled children. *J Intellect Disabil Res* 2014; **58/3**: 211-220.
- Souza NC, Mendonca JN, Portari GV, Jordao JA, Marchini JS, Chiarello PG. Intestinal permeability and nutritional status in developmental disorders. *Alt Ther Health Med* 2011; **18/2**: 19-24.
- Stedman KV, Leland Jr LS. Obesity and intellectual disability in New Zealand. *J Intellect Dev Disabil* 2010; **35/2**: 112-115.
- Stewart L, Van de Ven L, Katsarou V *et al*. High prevalence of obesity in ambulatory children and adolescents with intellectual disability. *J Intellect Disabil Res* 2009; **53/10**: 882-886.
- Suzuki M, Saitoh S, Tasaki Y, Shimomura Y, Makishima R, Hosoya N. Nutritional status and daily physical activity of handicapped students in Tokyo metropolitan schools for deaf, blind, mentally retarded, and physically handicapped individuals. *Am J Clin Nutr* 1991; **54/6**: 1101-1111.

- Vélez JC, Fitzpatrick AL, Barbosa CI, Díaz M, Urzua M, Andrade, AH. Nutritional status and obesity in children and young adults with disabilities in Punta Arenas, Patagonia, Chile. *Int J Rehabil Res* 2008; **31/4**: 305-313.
- Verstraelen CJF, Maaskant MA, Knijff-Raeven V, Curfs LMG, van Schrojenstein Lantman-de Valk HMJ. Weighting the weights: Agreement among anthropometric indicators identifying the weight status of people with intellectual disabilities. *J Appl Res Intellect Disabil* 2009; **22/3**: 307-313.
- Wallén EF, Müllersdorf M, Christensson K, Malm G, Ekblom Ö, Marcus C. High prevalence of cardio-metabolic risk factors among adolescents with intellectual disability. *Acta Paediatr* 2009; **98/5**: 853-859.
- Webb Y. Feeding and nutrition problems of physically and mentally handicapped children in Britain: a report. *Int J Food Sci Nutr* 1980; **34/4**: 281-285.
- Yamaki K, Rimmer JH, Lowry BD, Vogel LC. Prevalence of obesity-related chronic health conditions in overweight adolescents with disabilities. *Res Dev Disabil* 2011; **32/1**: 280-288.
- Yokohama Y. The incidence of obesity in the mentally retarded children. *Jpn J Spec Educ* 1983; **21/1**: 27-36.



**Table S1. Subgroup Analyses of Prevalence Estimates of *Overweight* for Children and Adolescents**

Moderators	Samples	Subgroups	N studies	Random effects models		Tests for heterogeneity				Subgroup analyses - mixed effects		
				Prevalence	95%CI	<i>Q</i>	df	<i>p</i>	<i>I</i> <sup>2</sup> (%)	<i>Q</i>	df	<i>p</i>
Sex	Children	Boys	4	14%	(8% to 21%)	25	3	< 0.001	88	0.003	1	0.95
		Girls	4	14%	(8% to 22%)	15	3	0.002	79			
	Adolescents	Boys	4	18%	(14% to 22%)	34	3	< 0.001	91			
		Girls	4	21%	(17% to 25%)	16	3	< 0.001	82			
Geographic region	Children	Europe	2	22%	(15% to 30%)	0	1	0.99	0	4.6	2	0.10
		North America	3	16%	(10% to 26%)	15	2	< 0.001	86			
		South America	2	14%	(12% to 17%)	0	1	0.94	0			
	Adolescents	Europe	3	17%	(15% to 20%)	4	2	0.16	45			
		North America	4	22%	(21% to 24%)	3	3	0.45	0			
Western Pacific	2	18%	(10% to 31%)	21	1	< 0.001	95					
Recruitment setting	Children	Special Olympics	2	20%	(13% to 29%)	10	1	< 0.001	90	0.6	1	0.45
		Special schools	2	13%	(5% to 31%)	11	1	< 0.001	91			
	Adolescents	Regular and special schools	2	18%	(10% to 32%)	8	1	< 0.001	87			
		Special Olympics	2	21%	(17% to 25%)	10	1	< 0.001	90			
		Special schools	2	16%	(11% to 22%)	15	1	< 0.001	93			
ID Level	Children	-	-	-	-	-	-	-	-	-	-	
	Adolescents	-	-	-	-	-	-	-	-	-	-	
Norms used to define OW	Children	IOTF	2	17%	(14% to 21%)	1.5	1	0.22	32	0.7	1	0.40
		National	3	12%	(5% to 26%)	37	2	< 0.001	95			
	Adolescents	IOTF	4	19%	(17% to 22%)	9	3	0.03	66			
		National	4	18%	(12% to 25%)	42	3	< 0.001	93			
Measurement of height and weight	Children	-	-	-	-	-	-	-	-	-	-	
	Adolescents	Direct	6	18%	(15% to 22%)	55	5	< 0.001	91	0.0004	1	0.98
		Indirect	2	18%	(14% to 24%)	0.4	1	0.54	0			
Children	-	-	-	-	-	-	-	-	-			
Additional diagnosis	Adolescents	Down syndrome	2	27%	(19% to 37%)	1.3	1	0.26	21	1.9	1	0.17
		ID without Down syndrome	2	19%	(13% to 28%)	2	1	0.15	52			

Note. ID = intellectual disability; IOTF = International Obesity Task Force; OW = overweight.

**Table S2. Studies Included in the Subgroup Analyses of Prevalence Estimates of *Overweight* for Children and Adolescents**

<b>Moderators</b>	<b>Samples</b>	<b>Subgroups</b>	<b>Studies</b>
<b>Sex</b>	Children	Boys vs. Girls (n = 4)	Choi et al. (31); Foley et al. (32); Fox et al. (33); Lloyd et al. (22)
	Adolescents	Boys vs. Girls (n = 4)	Choi et al. (31); Foley et al. (32); Krause et al. (34); Lloyd et al. (22)
<b>Geographic region</b>	Children	Europe (n = 2)	Bégarie et al. (30); Lloyd et al. (22)
		North America (n = 3)	Foley et al. (32); Fox et al. (33); Lloyd et al. (22)
		South America (n = 2)	Batista et al. (29); Lloyd et al. (22)
	Adolescents	Europe (n = 3)	Bégarie et al. (30); Lloyd et al. (22); Mikulovic et al. (36)
		North America (n = 4)	Foley et al. (32); Lloyd et al. (22); Phillips et al. (38); Rimmer et al. (39)
		Western Pacific (n = 2)	Choi et al. (31); Krause et al. (34)
<b>Recruitment setting</b>	Children	Special Olympics (n = 2)	Foley et al. (32); Lloyd et al. (22)
		Special schools (n = 2)	Bégarie et al. (30); Choi et al. (31)
	Adolescents	Regular and special schools (n = 2)	Krause et al. (34); Mikulovic et al. (36)
		Special Olympics (n = 2)	Foley et al. (32); Lloyd et al. (22)
<b>ID level</b>	Children	Not examined	Not examined
	Adolescents	Not examined	Not examined
<b>Norms used to define OW</b>	Children	IOTF (n = 2)	Bégarie et al. (30); Lloyd et al. (22)
		National (n = 3)	Choi et al. (31); Foley et al. (32); Fox et al. (33)
	Adolescents	IOTF (n = 4)	Bégarie et al. (30); Krause et al. (34); Lloyd et al. (22); Mikulovic et al. (36)
		National (n = 4)	Choi et al. (31); Foley et al. (32); Phillips et al. (38); Rimmer et al. (39)
<b>Measurement of height and weight</b>	Children	Not examined	Not examined
	Adolescents	Direct (n = 6)	Bégarie et al. (30); Choi et al. (31); Foley et al. (32); Krause et al. (34); Lloyd et al. (22); Mikulovic et al. (36)
		Indirect (n = 2)	Phillips et al. (38); Rimmer et al. (39)
<b>Additional diagnosis</b>	Children	No studies	No studies
	Adolescents	Down Syndrome (n = 2)	Krause et al. (34); Rimmer et al. (39)
		ID without Down syndrome (n = 2)	Krause et al. (34); Rimmer et al. (39)

Note. ID = intellectual disabilities; IOTF = International Obesity Task Force; OW = overweight

**Table S3. Subgroup Analyses of Prevalence Estimates of *Overweight-Obesity* for Children and Adolescents**

Moderators	Samples	Subgroups	N studies	Random effects models		Tests for heterogeneity				Subgroup analyses - mixed effects		
				Prevalence	95%CI	Q	df	p	I <sup>2</sup> (%)	Q	df	p
Sex	Children	Boys	5	29%	(19% to 41%)	55	4	< 0.001	93	0.005	1	0.95
		Girls	5	29%	(20% to 42%)	28	4	< 0.001	85			
	Adolescents	Boys	5	34%	(26% to 44%)	137	4	< 0.001	97	0.7	1	0.42
		Girls	5	39%	(32% to 47%)	51	4	< 0.001	92			
Geographic region	Children	Europe	3	32%	(25% to 40%)	0.3	2	0.87	0	13.7	2	0.001
		North America	3	39%	(31% to 48%)	10	2	0.01	81			
		South America	2	25%	(22% to 27%)	0.7	1	0.39	0			
	Adolescents	Europe	4	23%	(20% to 27%)	10	3	0.02	71	57.9	2	< 0.001
		North America	4	48%	(43% to 53%)	13	3	0.01	77			
		Western Pacific	2	36%	(19% to 58%)	44	1	< 0.001	98			
Recruitment setting	Children	Special Olympics	2	38%	(25% to 53%)	19	1	< 0.001	95	1.2	1	0.27
		Special schools	3	25%	(13% to 44%)	18	2	< 0.001	89			
	Adolescents	Regular and special schools	2	28%	(7% to 66%)	46	1	< 0.001	98	2.8	2	0.25
		Special Olympics	2	41%	(24% to 60%)	124	1	< 0.001	99			
		Special schools	3	27%	(25% to 28%)	0.4	2	0.80	0			
ID Level	Children	-	-	-	-	-	-	-	-	-	-	
	Adolescents	-	-	-	-	-	-	-	-	-	-	
Norms used to define OW-OB	Children	IOTF	3	31%	(29% to 33%)	0.3	2	0.86	0	0.1	1	0.76
		National	3	27%	(12% to 52%)	75	2	< 0.001	97			
	Adolescents	IOTF	5	28%	(22% to 36%)	65	4	< 0.001	94	1.4	1	0.24
		National	4	38%	(24% to 54%)	153	3	< 0.001	98			
Measurement of height and weight	Children	-	-	-	-	-	-	-	-	-	-	
	Adolescents	Direct	7	31%	(25% to 38%)	229	6	< 0.001	97	3.1	1	0.08
		Indirect	2	40%	(33% to 46%)	0.6	1	0.46	0			
Additional diagnosis	Children	-	-	-	-	-	-	-	-	-	-	
	Adolescents	Down syndrome	2	61%	(47% to 74%)	2	1	0.13	55	5.3	1	0.02
		ID without Down syndrome	2	35%	(22% to 52%)	6	1	0.01	84			

Note . ID = intellectual disability; IOTF = International Obesity Task Force; OW-OB = overweight-obesity.

**Table S4. Studies Included in the Subgroup Analyses of Prevalence Estimates of *Overweight-Obesity* for Children and Adolescents**

<b>Moderators</b>	<b>Samples</b>	<b>Subgroups</b>	<b>Studies</b>
<b>Sex</b>	Children	Boys vs. Girls (n = 5)	Choi et al. (31); Foley et al. (32); Fox et al. (33); Lloyd et al. (22); Salaun & Berthouze-Aranda (40)
	Adolescents	Boys vs. Girls (n = 5)	Choi et al. (31); Foley et al. (32); Krause et al. (34); Lloyd et al. (22); Salaun & Berthouze-Aranda (40)
<b>Geographic region</b>	Children	Europe (n = 3)	Bégarie et al. (30); Lloyd et al. (22); Salaun & Berthouze-Aranda (40)
		North America (n = 3)	Foley et al. (32); Fox et al. (33); Lloyd et al. (22)
		South America (n = 2)	Batista et al. (29); Lloyd et al. (22)
	Adolescents	Europe (n = 4)	Bégarie et al. (30); Lloyd et al. (22); Mikulovic et al. (36); Salaun & Berthouze-Aranda (40)
		North America (n = 4)	Foley et al. (32); Lloyd et al. (22); Phillips et al. (38); Rimmer et al. (39)
		Western Pacific (n = 2)	Choi et al. (31); Krause et al. (34)
<b>Recruitment setting</b>	Children	Special Olympics (n = 2)	Foley et al. (32); Lloyd et al. (22)
		Special schools (n = 3)	Bégarie et al. (30); Choi et al. (31); Salaun & Berthouze-Aranda (40)
	Adolescents	Regular and special schools (n = 2)	Krause et al. (34); Mikulovic et al. (36)
		Special Olympics (n = 2)	Foley et al. (32); Lloyd et al. (22)
		Special schools (n = 3)	Bégarie et al. (30); Choi et al. (31); Salaun & Berthouze-Aranda (40)
<b>ID level</b>	Children	Not examined	Not examined
	Adolescents	Not examined	Not examined
<b>Norms used to define OW-OB</b>	Children	IOTF (n = 3)	Bégarie et al. (30); Lloyd et al. (22); Salaun & Berthouze-Aranda (40)
		National (n = 3)	Choi et al. (31); Foley et al. (32); Fox et al. (33)
	Adolescents	IOTF (n = 5)	Bégarie et al. (30); Krause et al. (34); Lloyd et al. (22); Mikulovic et al. (36); Salaun & Berthouze-Aranda (40)
		National (n = 4)	Choi et al. (31); Foley et al. (32); Phillips et al. (38); Rimmer et al. (39)
<b>Measurement of height and weight</b>	Children	Not examined	Not examined
	Adolescents	Direct (n = 7)	Bégarie et al. (30); Choi et al. (31); Foley et al. (32); Krause et al. (34); Lloyd et al. (22); Mikulovic et al. (36); Salaun & Berthouze-Aranda (40)
		Indirect (n = 2)	Phillips et al. (38); Rimmer et al. (39)
<b>Additional diagnosis</b>	Children	No studies	No studies
	Adolescents	Down Syndrome (n = 2)	Krause et al. (34); Rimmer et al. (39)
		ID without Down syndrome (n = 2)	Krause et al. (34); Rimmer et al. (39)

Note. ID = intellectual disabilities; IOTF = International Obesity Task Force; OW = overweight

**Table S5. Subgroup Analyses of Prevalence Estimates of *Obesity* for Children and Adolescents**

Moderators	Samples	Subgroups	N studies	Random effects models		Tests for heterogeneity				Subgroup analyses - mixed effects		
				Prevalence	95%CI	Q	df	p	I <sup>2</sup> (%)	Q	df	p
Sex	Children	Boys	6	13%	(10% to 18%)	47	5	<0.001	89	0.1	1	0.78
		Girls	6	14%	(11% to 18%)	17	5	<0.001	71			
	Adolescents	Boys	6	16%	(12% to 21%)	131	5	<0.001	96			
		Girls	6	19%	(16% to 23%)	48	5	<0.001	90			
Geographic region	Children	Europe	2	9%	(5% to 16%)	0.1	1	0.73	0	40.6	3	<0.001
		North America	3	23%	(20% to 26%)	1.5	2	0.48	0			
		South America	2	11%	(8% to 14%)	1.3	1	0.26	23			
		Western Pacific	3	9%	(5% to 15%)	13	2	<0.001	84			
	Adolescents	Europe	4	8%	(5% to 13%)	20	3	<0.001	85			
		North America	4	27%	(24% to 30%)	7	3	0.08	55			
		South East Asia	2	10%	(5% to 21%)	27	1	<0.001	96			
		Western Pacific	4	16%	(13% to 19%)	24	3	<0.001	87			
Recruitment setting	Children	Special Olympics	2	17%	(12% to 24%)	6	1	0.02	83	5.3	1	0.02
		Special schools	3	9%	(5% to 14%)	13	2	0.002	84			
	Adolescents	Regular and special schools	4	13%	(6% to 25%)	33	3	<0.001	91			
		National survey	2	17%	(13% to 23%)	0.4	1	0.55	0			
		Special Olympics	2	20%	(9% to 37%)	126	1	<0.001	99			
		Special schools	4	12%	(10% to 14%)	25	3	<0.001	88			
ID Level	Children	-	-	-	-	-	-	-	-	-	-	
	Adolescents	-	-	-	-	-	-	-	-	-	-	
Norms used to define OB	Children	IOTF	2	14%	(12% to 17%)	1.1	1	0.30	6	0.1	1	0.73
		National	5	13%	(9% to 20%)	45	4	<0.001	91			
	Adolescents	IOTF	4	10%	(6% to 17%)	60	3	<0.001	95			
		National	7	19%	(14% to 25%)	165	6	<0.001	96			
Measurement of height and weight	Children	Direct	7	13%	(10% to 18%)	42	6	<0.001	86	1.3	1	0.26
		Indirect	2	11%	(10% to 12%)	0.2	1	0.70	0			
	Adolescents	Direct	9	14%	(11% to 19%)	200	8	<0.001	96			
		Indirect	4	17%	(12% to 23%)	15	3	<0.001	80			
Additional diagnosis	Children	-	-	-	-	-	-	-	-	-	-	
	Adolescents	Down syndrome	2	33%	(25% to 42%)	0.3	1	0.61	0	6.1	1	0.01
		ID without Down syndrome	2	17%	(10% to 27%)	3	1	0.11	61			


Note. ID = intellectual disability; IOTF = International Obesity Task Force; OB = obesity.

**Table S6. Studies Included in the Subgroup Analyses of the Prevalence Estimates of *Obesity* for Children and Adolescents**


<b>Moderators</b>	<b>Samples</b>	<b>Subgroups</b>	<b>Studies</b>
<b>Sex</b>	Children	Boys vs. Girls (n = 6)	Choi et al. (31); Foley et al. (32); Fox et al. (33); Lin et al. (35); Lloyd et al. (22); Takeuchi (41)
	Adolescents	Boys vs. Girls (n = 6)	Choi et al. (31); Foley et al. (32); Krause et al. (34); Lin et al. (35); Lloyd et al. (22); Takeuchi (41)
<b>Geographic region</b>	Children	Europe (n = 2)	Bégarie et al. (30); Lloyd et al. (22)
		North America (n = 3)	Foley et al. (32); Fox et al. (33); Lloyd et al. (22)
		South America (n = 2)	Batista et al. (29); Lloyd et al. (22)
		Western Pacific (n = 3)	Choi et al. (31); Lin et al. (35); Takeuchi (41)
Adolescents	Europe (n = 4)	Bégarie et al. (30); Lloyd et al. (22); Mikulovic et al. (36); Nogay (37)	
	North America (n = 4)	Foley et al. (32); Lloyd et al. (22); Phillips et al. (38); Rimmer et al. (39)	
	South East Asia (n = 2)	Lloyd et al. (22); Tamin et al. (42)	
	Western Pacific (n = 4)	Choi et al. (31); Krause et al. (34); Lin et al. (35); Takeuchi (41)	
<b>Recruitment setting</b>	Children	Special Olympics (n = 2)	Foley et al. (32); Lloyd et al. (22)
		Special schools (n = 3)	Bégarie et al. (30); Choi et al. (31); Takeuchi (41)
	Adolescents	Regular and special schools (n = 4)	Abdallah et al. (28); Krause et al. (34); Mikulovic et al. (36); Nogay (37)
		National survey (n = 2)	Lin et al. (35); Phillips et al. (38)
		Special Olympics (n = 2)	Foley et al. (32); Lloyd et al. (22)
Special schools (n = 4)	Bégarie et al. (30); Choi et al. (31); Takeuchi (41); Tamin et al. (42)		
<b>ID level</b>	Children	Not examined	Not examined
	Adolescents	Not examined	Not examined
<b>Norms used to define OB</b>	Children	IOTF (n = 2)	Bégarie et al. (30); Lloyd et al. (22)
		National (n = 5)	Choi et al. (31); Foley et al. (32); Fox et al. (33); Lin et al. (35); Takeuchi (41)
	Adolescents	IOTF (n = 4)	Bégarie et al. (30); Krause et al. (34); Lloyd et al. (22); Mikulovic et al. (36)
		National (n = 7)	Choi et al. (31); Foley et al. (32); Lin et al. (35); Nogay (37); Phillips et al. (38); Rimmer et al. (39); Takeuchi (41)
<b>Measurement of height and weight</b>	Children	Direct (n = 7)	Abdallah et al. (28); Batista et al. (29); Bégarie et al. (30); Choi et al. (31); Foley et al. (32); Fox et al. (33); Lloyd et al. (22)
		Indirect (n = 2)	Lin et al. (35); Takeuchi (41)
	Adolescents	Direct (n = 9)	Abdallah et al. (28); Bégarie et al. (30); Choi et al. (31); Foley et al. (32); Krause et al. (34); Lloyd et al. (22); Mikulovic et al. (36); Nogay (37); Tamin et al. (42)
		Indirect (n = 4)	Lin et al. (35); Phillips et al. (38); Rimmer et al. (39); Takeuchi (41)
<b>Additional diagnosis</b>	Children	Not examined	Not examined
	Adolescents	Down Syndrome (n = 2)	Krause et al. (34); Rimmer et al. (39)
ID without Down syndrome (n = 2)		Krause et al. (34); Rimmer et al. (39)	

*Note.* ID = intellectual disabilities; IOTF = International Obesity Task Force; OW = overweight

(a)



(b)



(c)






Figure S1. Funnel Plot of Standard Error by Logit Event Rate of (a) Overweight, (b) Overweight-Obesity, and (c) Obesity among *Children*


(a)



(b)



(c)



**Figure S2. Funnel Plot of Standard Error by Logit Event Rate of (a) Overweight, (b) Overweight-Obesity, and (c) Obesity among *Adolescents***