

Research Bank

Journal article

Muscle glycogen utilization during an Australian Rules Football game

Routledge, Harry E., Leckey, Jill J., Lee, Matt J., Garnham, Andrew, Graham, Stuart, Burgess, Darren, Burke, Louise M., Erskine, Robert M., Close, Graeme L. and Morton, James P.

Accepted author manuscript version reprinted, by permission, from International Journal of Sports Physiology and Performance, 2019, 14 (1): 122-124,

<https://doi.org/10.1123/ijsp.2018-0106>. © 2019 Human Kinetics, Inc.

Note. This article will be published in a forthcoming issue of the *International Journal of Sports Physiology and Performance*. The article appears here in its accepted, peer-reviewed form, as it was provided by the submitting author. It has not been copyedited, proofread, or formatted by the publisher.

Section: Brief Review

Article Title: Muscle Glycogen Utilisation during an Australian Rules Football Game

Authors: Harry E Routledge^{1,2}, Jill J Leckey³, Matt J Lee⁴, Andrew Garnham³, Stuart Graham², Darren Burgess², Louise M Burke³, Robert M Erskine^{1,5}, Graeme L Close¹ and James P Morton¹

Affiliations: ¹Research Institute for Sport and Exercise Sciences, Liverpool John Moores University, Liverpool, UK. ²Port Adelaide Football Club, Alberton, SA, Australia. ³Exercise and Nutrition Research Program, Mary MacKillop Institute for Health Research, Australian Catholic University, Melbourne, Victoria, Australia. ⁴College of Sport and Exercise Science, Victoria University, Melbourne, Victoria, Australia. ⁵Institute of Sport, Exercise and Health, University College London, London, UK.

Journal: *International Journal of Sports Physiology and Performance*

Acceptance Date: May 21, 2018

©2018 Human Kinetics, Inc.

DOI: <https://doi.org/10.1123/ijsp.2018-0106>

Muscle Glycogen Utilisation during an Australian Rules Football Game

Submission Type: Case-Study

Harry E Routledge^{1,2}, Jill J Leckey³, Matt J Lee⁴, Andrew Garnham³, Stuart Graham², Darren Burgess², Louise M Burke³, Robert M Erskine^{1,5}, Graeme L Close¹ and James P Morton¹

¹Research Institute for Sport and Exercise Sciences
Liverpool John Moores University
Byrom Street Campus
Liverpool
L3 3AF
UK

²Port Adelaide Football Club
Allan Scott Headquarters
Alberton Oval
Brougham Place
Alberton
SA 5014
Australia

³Exercise and Nutrition Research Program
Mary MacKillop Institute for Health Research
Australian Catholic University
Melbourne
Victoria, Australia

⁴ College of Sport and Exercise Science
Victoria University
Melbourne
Victoria
Australia

⁵Institute of Sport, Exercise and Health,
University College London,
London,
United Kingdom

Address for correspondence:

Dr James Morton
Research Institute for Sport and Exercise Sciences
Liverpool John Moores University
Tom Reilly Building
Byrom St Campus
Liverpool
L3 3AF
United Kingdom
Email: J.P.Morton@ljmu.ac.uk
Tel: +44 151 904 6233

Running Head: Glycogen use in AF match play

Abstract Word Count: 248

Text Only Word Count: 1492

Number of Figures and Tables: 1 Table

Abstract

Purpose: To better understand the carbohydrate (CHO) requirement of Australian Football (AF) match play by quantifying muscle glycogen utilisation during an in-season AF match.

Methods: After a 24 h CHO loading protocol of 8 g/kg and 2 g/kg in the pre-match meal, two elite male forward players had biopsies sampled from *m. vastus lateralis* before and after participation in a South Australian Football League game. Player A (87.2kg) consumed water only during match play whereas player B (87.6kg) consumed 88 g CHO via CHO gels. External load was quantified using global positioning system technology. **Results:** Player A completed more minutes on the ground (115 vs. 98 min) and covered greater total distance (12.2 vs. 11.2 km) than Player B, though with similar high-speed running (837 vs. 1070 m) and sprinting (135 vs. 138 m), respectively. Muscle glycogen decreased by 66% in Player A (Pre-: 656, Post-: 223 mmol·kg⁻¹ dw) and 24% in Player B (Pre-: 544, Post-: 416 mmol·kg⁻¹ dw), respectively. **Conclusion:** Pre-match CHO loading elevated muscle glycogen concentrations (i.e. >500 mmol·kg⁻¹ dw), the magnitude of which appears sufficient to meet the metabolic demands of elite AF match play. The glycogen cost of AF match play may be greater than soccer and rugby and CHO feeding may also spare muscle glycogen use. Further studies using larger sample sizes are now required to quantify the inter-individual variability of glycogen cost of match play (including muscle and fibre-type specific responses) as well examine potential metabolic and ergogenic effects of CHO feeding.

Keywords: carbohydrate loading, vastus lateralis, AF, high speed running

Introduction

Invasive team sports such as soccer^{1,2}, rugby league² and Australian Football (AF)^{2,3} are characterised by high-intensity (>19.8km.h⁻¹) intermittent activity profiles. Given the duration of activity (i.e. 80-120 minutes) and high-intensity intermittent profiles, muscle glycogen is considered as the predominant energy substrate to support the metabolic demands of match play^{4,5}. In relation to soccer, it has been reported that match-play in elite Danish soccer players⁵ depletes muscle glycogen concentration by a magnitude of 50% and requires a glycogen cost of approximately 200 mmol·kg⁻¹ dw. We also observed similar absolute glycogen utilisation and relative depletion rates in English professional rugby league players during competitive match play⁶. Nonetheless, in the absence of a controlled carbohydrate (CHO) loading protocol, it is noteworthy that pre-match muscle glycogen concentrations in team sport athletes may range from 300-600 mmol·kg⁻¹ dw⁷ and that approximately 50% of muscle fibres are classified as empty or partially empty after soccer match play, thus having potential implications for high-intensity physical performance late in the match. Indeed, in relation to physical performance, it was recognised as early as the 1970s⁸ that commencing match play with reduced pre-exercise muscle concentration (i.e. < 200 mmol·kg⁻¹ dw) reduces total distance covered by 25% when compared with higher pre-match muscle glycogen availability (i.e. > 400 mmol·kg⁻¹ dw).

Given the longer duration and greater proportion of time spent at high-intensity workloads in AF match play^{2,3} versus both soccer^{1,2} and rugby², it could be suggested that the CHO requirements for AF players are increased accordingly. Indeed, high velocity running was reported to be significantly greater in AF (1322m) versus rugby league (327m) and soccer (517m)². To the authors' knowledge, however, no researchers have yet quantified the muscle glycogen cost of AF match play in elite players. The aim of the present case-study was to

quantify glycogen use in *m. vastus lateralis* from two elite male AF players after participation in a South Australian Football League game.

Methods

Subjects

Two male forward players from a South Australian Football League (SANFL) list (see Table 1) volunteered to take part in the study. The study was conducted according to the Declaration of Helsinki and was approved by the Human Research Ethics Committee of Australian Catholic University Melbourne.

Design

In a case-study design, muscle biopsies were obtained from *m. vastus lateralis* before and after participation in a competitive SANFL league game undertaken in August 2017. Quantification of external loading during match play was assessed via global positioning system (GPS) technology.

Dietary Controls. In the day prior to match play, each player consumed a prescribed CHO loading diet providing CHO, protein and fat intakes corresponding to 8 g/kg, 2 g/kg and 1 g/kg body mass, respectively, based on contemporary guidelines of Thomas and colleagues⁹. At 4 hours prior to match play, each player also consumed a pre-match meal providing 2 g/kg CHO, 0.2 g/kg protein and 0.3 g/kg fat. During match play, players were able to consume fluids and sports foods at opportunities that were both predictable (between quarters) and random (rotation on the bench, access to trainer on the ground when the ball is not in play). On such occasions; Player A consumed water only whereas Player B also consumed an additional 88 g of CHO in the form of four isotonic CHO gels (SiS GO Gels, UK) consumed at the quarter time breaks (~20-minute intervals during the game).

Muscle Biopsies. Muscle biopsies (100 mg) were sampled from *m. vastus lateralis* at 60 minutes prior to the beginning of the game (i.e. prior to the warm-up period) and within 10

minutes of completion of match play participation. Biopsies were obtained under local anesthesia (1% Xylocaine) using a Bergstrom biopsy needle and immediately frozen in liquid nitrogen for later analysis. Approximately 3 mg of freeze-dried muscle tissue was powdered and dissected free of all visible non-muscle tissue. Powdered muscle tissue was then mixed with 250 μl of 2 M HCl, incubated at 95°C for 2 hours (agitated gently every 20 min), and then neutralized with 750 μl of 0.66 M NaOH. Glycogen concentration was subsequently assayed in triplicate via enzymatic analysis with fluorometric detection. Muscle glycogen values were expressed as millimoles per kilograms dry weight ($\text{mmol}\cdot\text{kg}^{-1}\text{ dw}$).

GPS Analysis. Both players wore a portable micro-technology device (Optimeye S5, Catapult Innovations, Australia) which recorded activity profile data. The portable device was worn inside a purpose built elastic vest, positioned across the upper back between the scapula. Each device was activated 30-minutes prior to the start of the game to allow acquisition of satellite signal and lock (>8 satellites). Satellite data sampled at 10 Hz provided measures of duration, total distance, average speed and distance covered within four specific velocity bands corresponding to: jogging (2-3.9 $\text{m}\cdot\text{s}^{-2}$), running (4-5.4 $\text{m}\cdot\text{s}^{-2}$), high-speed running (5.5-6.9 $\text{m}\cdot\text{s}^{-2}$) and sprinting (7 $\text{m}\cdot\text{s}^{-2}$). These speed zones are similar to those previously reported in soccer¹ and allow for direct comparisons between football codes. At the conclusion of the match, data were downloaded and analyzed using (Openfield version 11.12, Catapult Innovations).

Results

Subject characteristics, external workload, muscle glycogen utilisation and CHO intake are presented in Table 1. Individual data are presented for both Player A and B.

Discussion

The aim of the present case-study was to quantify muscle glycogen utilisation during AF match play. We studied two elite male AF players during a competitive match from the

South Australian Football League. Our data demonstrate that a standardised one day dietary CHO loading protocol of 8 g/kg and 2 g/kg in the pre-match meal elevates muscle glycogen concentration (i.e. $>500 \text{ mmol.kg}^{-1} \text{ dw}$) to a magnitude that is sufficient to fuel the metabolic demands of AF match play.

In relation to the external loads reported here, we observed similar loading profiles to that previously reported in elite AF match play where larger sample sizes (e.g. 39 players) have been studied³. This is the case for parameters such as total distance, average speed, high-intensity running and sprinting. The between player differences in such parameters are also similar to those previously reported³. Despite only studying two forward players, we are confident that the external loads reported here are therefore representative of the customary loads experienced in AF match play from a wider sample of teams and players, though we acknowledge that load differences between playing positions are to be expected³.

In accordance with a higher workload, our data suggest that the glycogen cost of AF match play (e.g. Player A utilised $433 \text{ mmol.kg}^{-1} \text{ dw}$) may be greater than that reported in soccer^{5,7} and rugby match play⁶ where approximately $200 \text{ mmol.kg}^{-1} \text{ dw}$ was utilised in both instances. Such differences in absolute glycogen cost are likely due to greater duration of activity and time spent in higher intensity threshold zones. In contrast to Player A, Player B experienced less total glycogen use ($138 \text{ mmol.kg}^{-1} \text{ dw}$). While such inter-individual variation in glycogen use may be due to differences in total distance covered, duration, pre-match muscle glycogen concentrations¹¹, training status and muscle oxidative capacity¹², it is noteworthy that Player B also consumed an additional 88 g of CHO during match play. As such, differences in glycogen use between players may also be due, in part, to a potential muscle glycogen sparing effect of CHO feeding, an effect that has been reported previously in *m. vastus lateralis* during running¹³. Indeed, the game characteristics of AF support a more aggressive approach to CHO intake during play than is reported by elite soccer players², with opportunities for fuel

replacement during scheduled breaks between quarters and time spent on the interchange bench. Nonetheless, we acknowledge that randomised control trials incorporating larger sample sizes are now required to verify any potential metabolic or ergogenic effect of CHO feeding during competitive match play. Additionally, glycogen use in specific muscles (e.g. gastrocnemius versus vastus lateralis), muscle fibre types and intra-cellular storage pools could also be quantified using transmission electron microscopy¹⁴. We also acknowledge that the glycogen utilisation observed here is also reflective of those activities undertaken during the warm-up period. As such, future studies could also sample biopsies in the minutes prior to match play and at the end of each quarter to further characterise both the total absolute use and rates of glycogen use as the match progresses.

Practical Applications

Pre-match muscle glycogen concentration $> 500 \text{ mmol.kg}^{-1} \text{ dw}$ (as achieved via CHO loading of 8 g/kg) is sufficient to fuel the physical demands of elite forward AF match play. Given the apparently greater muscle glycogen cost of AF match-play compared to soccer and rugby, sport physiologists and nutritionists should ensure that AF players consume sufficient dietary CHO intake (likely $> 6 \text{ g/kg}$) in the 24 hours prior to participation in match play.

Conclusion

We provide novel data by reporting muscle glycogen utilisation in *m. vastus lateralis* from two elite male AF forward players during competitive match play. Our data suggest that total glycogen use is greater than that reported in elite players from other invasive team sports, such as soccer and rugby. Additionally, these data suggest that CHO loading with 8 g/kg body mass is sufficient to meet the metabolic demands of AF match play and that previous suggestions of 10-12 g/kg body (for athletes involved in intermittent exercise > 90 minutes) mass may not be necessary for this population. Further studies are now required to quantify the

“Muscle Glycogen Utilisation during an Australian Rules Football Game” by Routledge HE et al.
International Journal of Sports Physiology and Performance
© 2018 Human Kinetics, Inc.

inter-individual variability of glycogen use as well as examine any potential metabolic and ergogenic effects of CHO feedings during match play.

References

1. Anderson L, Orme P, Di Michele R, Close GL, Morgans R, Drust B, Morton JP. Quantification of training load during one-, two- and three-game week schedules in professional soccer players from the English Premier League: implications for carbohydrate periodisation. *J Sports Sci.*2016;34(13):1250-1259.
2. Varley MC, Gabbett T, Aughey RJ. Activity profiles of professional soccer, rugby league and Australian football match play. *J Sports Sci.*2014;32(20):1858-1866.
3. Coutts AJ, Kempton T, Sullivan C, Bilsborough J, Cordy J, Rampinini E. Metabolic power and energetic costs of professional Australian Football match-play. *J Sci Med Sport.*2015;18(2):219-224.
4. Mujika I, Burke LM. Nutrition in team sports. *Ann Nutr Metab.*2010;57(2):26-35.
5. Gunnarsson TP, Bendiksen M, Bischoff R, Christensen PM, Lesvig B, Madsen K, Stephens F, Greenhaff P, Krstrup P, Bangsbo J. Effect of whey protein- and carbohydrate-enriched diet on glycogen resynthesis during the first 48 h after a soccer game. *Scand J Med Sci Sports.*2013;23(4):508-515.
6. Bradley WJ, Morehen JC, Haigh J, Clarke J, Donovan TF, Twist C, Cotton C, Shepard, S, Cocks M, Sharma A, Impey SG, Cooper RG, MaClaren DP, Morton JP, Close GL. Muscle glycogen utilisation during Rugby match play: Effects of pre-game carbohydrate. *J Sci Med Sport.* 2016;19(12):1033-1038.
7. Krstrup, P, Mohr M, Steensberg A, Bencke, J, Kjaer M, Bangsbo J. Muscle and blood metabolites during a soccer game: implications for sprint performance. *Med Sci Sports Exerc.*2006;38(6):1165-1174.
8. Saltin B. Metabolic fundamentals in exercise. *Med Sci Sports.*1973;5(3):137-146.
9. Thomas DT, Erdman KA, Burke LM. Position of the academy of nutrition and dietetics, dietitians of Canada, and the American college of sports medicine: nutrition and athletic performance. *J Acad Nutr Diet.*2016;116(3):501-28.
10. Bangsbo J, Norregaard L, Thorsoe F. The effect of carbohydrate diet on intermittent exercise performance. *Int J Sports Med.*1992;13(2):152-157.
11. Wisbey B, Montgomery PG, Pyne DB, Rattray B. Quantifying movement demands of AFL football using GPS tracking. *J Sci Med Sport.*2010;13:531-536.
12. Hargreaves M, McConell G, Proietto J. Influence of muscle glycogen on glycogenolysis and glucose uptake during exercise in humans. *J Appl Physiol.*1995;78(1):288-292.
13. Holloszy JO, Coyle EF. Adaptations of skeletal muscle to endurance exercise and their metabolic consequences. *J Appl Physiol Respir Environ Exerc Physiol.*1984;56(4):831-838.

14. Tsintzas OK, Williams C, Boobis L, Greenhaff P. Carbohydrate ingestion and glycogen utilization in different muscle fibre types in man. *J Physiol.*1995;489 :243-250.
15. Nielsen J, Krstrup P, Nybo L, Gunnarsson TP, Madsen K, Schroder HD, Bangsbo J, Ortenblad N. Skeletal muscle glycogen content and particle size of distinct subcellular localizations in the recovery period after a high-level soccer match. *Appl Physiol.*2012;112(10):3559-3367.

Table 1 – Subject characteristics, external workload, muscle glycogen utilisation and in-game CHO intake in Player A and B.

	Player A	Player B
Playing Position	Forward	Forward
Age (years)	22	27
Body Mass (kg)	87.2	87.6
Height (m)	1.88	1.82
Warm Up Duration (min)	12	12
Distance (m)	1478	1501
High Speed Running (m) 5.5-6.9 m.s²	156	159
Sprinting (m) >7 m.s²	23	31
Match Play Duration (min)	115	98
Number of Rotations	2	4
Average Speed (m/min)	106	114
Total Distance (m)	12229	11182
Walking (m) 0.1-1.9 m.s²	3801	2801
Jogging (m) 2-3.9 m.s²	5231	4718
Running (m) 4-5.4 m.s²	2180	2396
High Speed Running (m) 5.5-6.9 m.s²	837	1070
Sprinting (m) >7 m.s²	135	138
Pre-Match Glycogen (mmol.kg⁻¹ dw)	656	544
Post-Match Glycogen (mmol.kg⁻¹ dw)	223	416
Total Glycogen Utilisation (mmol.kg⁻¹ dw)	433	138
Total Exogenous CHO Consumed (g)	0	88
Exogenous CHO (g/h)	0	54
Exogenous CHO (g/min)	0	0.9