

# Television Viewing and Low Leisure-Time Physical Activity in Adolescence Independently Predict the Metabolic Syndrome in Mid-Adulthood

PATRIK WENNBERG, PHD<sup>1</sup>  
 PER E. GUSTAFSSON, PHD<sup>1</sup>  
 DAVID W. DUNSTAN, PHD<sup>2</sup>

MARIA WENNBERG, PHD<sup>1</sup>  
 ANNE HAMMARSTRÖM, PHD<sup>1</sup>

**OBJECTIVE**—We investigated whether television (TV) viewing and low leisure-time physical activity in adolescence predict the metabolic syndrome in mid-adulthood.

**RESEARCH DESIGN AND METHODS**—TV viewing habits and participation in leisure-time physical activity at age 16 years were assessed by self-administered questionnaires in a population-based cohort in Northern Sweden. The presence of the metabolic syndrome at age 43 years was ascertained in 888 participants (82% of the baseline sample) using the International Diabetes Federation criteria. Odds ratios (ORs) and CIs were calculated using logistic regression.

**RESULTS**—The overall prevalence of the metabolic syndrome at age 43 years was 26.9%. Adjusted OR for the metabolic syndrome at age 43 years was 2.14 (95% CI 1.24–3.71) for those who reported “watching several shows a day” versus “one show/week” or less and 2.31 (1.13–4.69) for leisure-time physical activity “several times/month” or less compared with “daily” leisure-time physical activity at age 16 years. TV viewing at age 16 years was associated with central obesity, low HDL cholesterol, and hypertension at age 43 years, whereas low leisure-time physical activity at age 16 years was associated with central obesity and triglycerides at age 43 years.

**CONCLUSIONS**—Both TV viewing and low leisure-time physical activity in adolescence independently predicted the metabolic syndrome and several of the metabolic syndrome components in mid-adulthood. These findings suggest that reduced TV viewing in adolescence, in addition to regular physical activity, may contribute to cardiometabolic health later in life.

*Diabetes Care* 36:2090–2097, 2013

Current public health guidelines recommend moderate-intensity aerobic physical activity for a minimum of 30 min on 5 days each week or vigorous-intensity aerobic physical activity for a minimum of 20 min on 3 days each week to promote and maintain health (1). Aside from this, sedentary behavior such as prolonged sitting during commuting and in the workplace and the domestic environment has recently emerged as a distinct contributor to adverse health effects, particularly for cardiometabolic outcomes (2). On the physical activity continuum, sedentary behaviors

do not simply exist at one end but, rather, are a class of behaviors that can coexist and compete with physical activity (3). Changes in television (TV)-viewing levels (a common leisure-time sedentary behavior) have been associated with concurrent changes in cardiometabolic biomarkers that are independent of the protective effects of regular moderate- to vigorous-intensity physical activity (4). Generally, evidence for a physical activity-independent link between TV viewing and cardiometabolic outcomes has been generated from cross-sectional analyses. Few longitudinal studies have been conducted, and a

recently published review concluded that there is insufficient evidence to conclude that a definitive longitudinal relationship exists between sedentary behavior, markers of cardiometabolic health, and metabolic conditions (2). Moreover, there is a lack of prospective studies that have examined the relationship between sedentary behaviors during childhood or adolescence and adult cardiometabolic risk. One of the few longitudinal studies with a life course perspective showed that TV viewing in childhood was associated with raised cholesterol and overweight in adulthood (5). However, the individuals were not followed longer than up to age 26 years. Thus, although interventions to reduce TV viewing in children have shown promising short-term results (6), uncertainty remains as to whether preventive actions during childhood and adolescence may have an impact on cardiometabolic risk in adulthood.

The aim of this study was to investigate whether TV viewing and low leisure-time physical activity in adolescence predict the metabolic syndrome in mid-adulthood. We also aimed to examine the extent to which leisure-time physical activity in mid-adulthood mediates the relationships of TV viewing and low leisure-time physical activity in adolescence with the metabolic syndrome in mid-adulthood.

## RESEARCH DESIGN AND METHODS

Participants were drawn from the Northern Swedish Cohort, a 27-year prospective cohort study previously described in detail (7). The sample is defined as all school leavers of the 9th (final) grade of the Swedish compulsory school in 1981 in the municipality of Luleå at participant age 16 years ( $n = 1,083$ ). Ethics approval was granted by the regional ethics review board in Umeå, Sweden. The cohort has been found to be representative of the corresponding age cohort of Sweden in various demographic comparisons (7). Follow-up data collections

From the <sup>1</sup>Department of Public Health and Clinical Medicine, Family Medicine, Umeå University, Umeå, Sweden; and the <sup>2</sup>Baker IDI Heart and Diabetes Institute, Melbourne, Victoria, Australia.

Corresponding author: Patrik Wennberg, patrik.wennberg@fammed.umu.se.

Received 24 September 2012 and accepted 11 December 2012.

DOI: 10.2337/dc12-1948

© 2013 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

were conducted in 1983 (age 18 years), 1986 (age 21 years), 1995 (age 30 years), and 2008 (age 43 years). At the 2008 follow-up,  $n = 1,010$  were still active in the cohort (94% of those of the baseline sample who were still alive [ $n = 1,071$ ]).

The present report is based on data from the age 16 and 43 years data collections. As a result of internal dropout, the analyses were conducted in 888 participants (464 men and 424 women): 82% of the baseline sample and 88% of those participating at the 2008 data collection. (See the statistical analysis, below, for details and analysis of the missing data.)

At ages 16 and 43 years, participants completed comprehensive self-administered questionnaires. The questionnaires covered health, behaviors, leisure activities, and work, family, school, and other social circumstances (7) and were used in the present report to operationalize lifestyle factors, socioeconomic status, and family history of diabetes. Information on weight and height at age 16 years was retrieved from school health records, as measured by school nurses as a part of a compulsory school health examination, and was used to calculate BMI. For the operationalization of the metabolic syndrome at age 43 years, the participants took part in a more comprehensive health examination performed by trained medical personnel at the participants' respective health care center (8). Blood pressure (two consecutive readings) was measured according to the World Health Organization Multinational MONItoring of trends and determinants in CARdiovascular disease (MONICA) manual (9). Waist circumference was measured to the nearest 0.5 cm in indoor clothing. Blood samples were drawn after an overnight fast and analyzed with respect to HDL cholesterol (HDL-C), triglycerides, and glucose. Coefficient of variance at high and low concentrations was 1.7 and 1.5% for triglycerides, 2.8 and 2.8% for HDL-C, and 1.5 and 1.2% for glucose, respectively.

### TV-viewing habits and participation in leisure-time physical activity at age 16 years

TV-viewing habits were operationalized through the question, "How often do you watch television?" with the following five response options: 1) several shows a day, 2) one show per day, 3) one show every other day, 4) one show per week, and 5) less than one show per week. Because of the low number in several categories (e.g.,

14 participants in the fifth category), we collapsed category 2 with category 3 and category 4 with category 5. Consequently, analyses were performed using three levels of TV viewing. Participants reported leisure-time physical activity (sport activities or exercise) during the last 12 months, with six response options: daily, several times per week, once a week, several times per month, once a month, and seldom. The last three categories were collapsed based on the low number in each of these categories. Additionally, the same question regarding leisure-time physical activity (coded in the same way) was used at age 43 years.

### Metabolic syndrome and its components at age 43 years

The metabolic syndrome and its components were operationalized in a manner similar to that in our previous reports (8,10,11), according to the guidelines by the International Diabetes Federation (12): waist circumference  $\geq 80$  cm for women and  $\geq 94$  cm for men and two or more of the following criteria: 1) increased triglycerides ( $\geq 1.7$  mmol/L) or specific treatment for that lipid abnormality, 2) reduced HDL-C ( $< 1.29$  mmol/L for women and  $< 1.03$  mmol/L for men) or specific treatment for that lipid abnormality, 3) increased blood pressure (systolic blood pressure  $\geq 130$  mmHg or diastolic blood pressure  $\geq 85$  mmHg) or treatment of hypertension, and 4) increased fasting glucose ( $\geq 5.6$  mmol/L) or diagnosed type 2 diabetes. Information on medication was based on self-report. Current type 2 diabetes was based on self-reported presence of diabetes at age 43 years, excluding those who also reported diabetes at age 30 years, who were regarded as having type 1 diabetes. Both the metabolic syndrome and its components were used as binary outcomes in the subsequent analyses.

### Covariates

Socioeconomic status in adolescence was operationalized in the same manner as in our previous reports (8,10,11). Parental occupation, as reported by participants at age 16 years, was coded into three social groups, corresponding to the manual workers (blue-collar) (III), lower white-collar employees (II), and upper white-collar employees and self-employed (I) classification scheme of Statistics Sweden (13). As only 8.5% of the fathers and 2.5% of the mothers belonged to group I, groups I and II were collapsed into a

favorable socioeconomic category, whereas the manual group was retained as a disadvantaged socioeconomic category. Socioeconomic disadvantage was defined as both parents belonging to manual occupation versus one or both parents belonging to a nonmanual or self-employed occupation. For a participant who only lived with one parent or who had one unknown or deceased parent, only the occupation of the parent with whom they were currently living was considered. For parents who were not currently employed, the last held occupation was used. Annual consumption of alcohol was estimated from self-reported typical frequency and quantity of alcoholic beverage consumption and categorized into sex-specific quintiles. Due to the high percentage of abstainers, quintiles 1 and 2 were collapsed, and the final alcohol consumption variable thus comprised four levels of alcohol consumption (10). Family history of diabetes and smoking habits were self-reported in the questionnaire and dichotomized. Dietary intake was assessed in a short simplified food-frequency questionnaire, and a calculation of nutrient intake was not possible. However, we included a single question on intake of sweets and pastries, which was dichotomized. BMI was calculated from measured height and weight as weight in kilograms divided by the square of height in meters.

Overweight was defined as BMI 25.0–29.9 kg/m<sup>2</sup> and obesity as BMI  $\geq 30$  kg/m<sup>2</sup>. Incident overweight or obesity was defined as BMI  $\geq 25$  kg/m<sup>2</sup> at follow-up (age 43 years) among participants with a sex-specific BMI  $< 90$ th percentile at baseline (age 16 years).

### Statistical analysis

Statistical analyses were conducted using SPSS statistics, version 19 (SPSS, Chicago, IL).  $\chi^2$  test was used for testing differences in proportions. Kruskal-Wallis test was used for testing group differences in ordinal continuous variables. Correlation between TV viewing and leisure-time physical activity was calculated using Spearman rank correlation. Logistic regression was used to estimate odds ratios (ORs) and 95% CIs. Multivariate models included TV viewing and leisure-time physical activity in the same model and a model with additional adjustment for sex, socioeconomic disadvantage, family history of diabetes, BMI, intake of sweets/pastries, alcohol consumption, and smoking at age 16 years. In order to

test whether the associations between TV viewing and low leisure-time physical activity with metabolic syndrome were mediated by leisure-time physical activity in adulthood, we also included leisure-time physical activity at age 43 years in the last multivariate model (model 3). We performed a test for interactions of TV viewing and leisure-time physical activity by including an interaction term in the multivariate-adjusted model (model 2). The same multivariate model was used in analyses of the metabolic syndrome components. A  $P$  value  $<0.05$  was considered significant. In the study, 107 individuals (10.7%) were excluded because data on the metabolic syndrome were missing and 6 individuals (0.6%) were excluded because data on TV viewing or leisure-time physical activity were missing. For examination of potential selection bias, those with incomplete measures were compared with those with complete measures. There were no differences in TV viewing, leisure-time physical activity, or any of the included covariates in this study. A total of 888 individuals remained after exclusions. The multivariate analyses were conducted as a complete case analysis in which 19 additional cases were lost as a result of missing data on covariates.

**RESULTS**—The overall prevalence of the metabolic syndrome at age 43 years among the 888 participants was 26.9%, and the incident overweight or obesity was 54.5%. Selected characteristics according to TV-viewing habits and participation in leisure-time physical activity at age 16 years are presented in Table 1. A higher frequency of TV viewing at age 16 years was related to male sex, socioeconomic disadvantage, lower alcohol consumption, and lower proportion of daily smoking at age 16 years and to prevalence of the metabolic syndrome and all metabolic syndrome components except for central obesity at age 43 years. Boys' and girls' participation in leisure-time physical activity at age 16 years was unevenly distributed with a higher proportion of girls compared with boys in one category: once per week. A lower frequency of leisure-time physical activity at age 16 years was related to socioeconomic disadvantage, higher alcohol consumption, and a higher proportion of daily smokers at age 16 years and to prevalence of the metabolic syndrome, central obesity, and raised triglycerides at age 43 years. TV

viewing and leisure-time physical activity at age 16 years were not closely related (Spearman correlation coefficient  $-0.032$ ,  $P = 0.343$ ).

Table 2 shows the ORs for the presence of the metabolic syndrome at age 43 years according to TV viewing and leisure-time physical activity at age 16 years. We found no significant interaction between TV viewing or low leisure-time physical activity and sex on presence of the metabolic syndrome at age 43 years ( $P = 0.704$  and  $P = 0.745$ , respectively). Thus, the analyses were not stratified by sex. TV viewing and low leisure-time physical activity were associated with the metabolic syndrome in crude analyses. These associations with the metabolic syndrome were only modestly attenuated when TV viewing and low leisure-time physical activity were included in the same model (model 1) and when analyses were further adjusted for the other covariates (model 2). To test whether the association between TV viewing and low leisure-time physical activity in adolescence with the metabolic syndrome was dependent on leisure-time physical activity in adulthood, we also added leisure-time physical activity at age 43 years to the multivariate-adjusted model (model 3). This adjustment attenuated the association for TV viewing somewhat, but the  $P$  for trend ( $P = 0.008$ ) across the TV-viewing categories as well as the OR for several shows per day versus one show per week or less remained significant (OR 1.96 [95% CI 1.13–3.42]), whereas the association for low leisure-time physical activity was more markedly attenuated and no longer significant.

Fig. 1 shows the relationship between TV viewing at age 16 years and the metabolic syndrome at age 43 years across two categories of physical activity. The interaction term between TV viewing and leisure-time physical activity on risk for the metabolic syndrome was not significant ( $P = 0.21$ ). Compared with those who watched one TV show per day or less and were physically active once a week or more, those who watched several TV shows per day had an increased risk for the metabolic syndrome irrespective of their level of leisure-time physical activity.

In crude analyses of the metabolic syndrome components, TV viewing at age 16 years was associated with all metabolic syndrome components except for central obesity at age 43 years, whereas low leisure-time physical activity at age 16

years was associated only with central obesity and raised triglycerides (Table 3). After multivariate adjustments, TV viewing at age 16 years was associated with central obesity, low HDL-C, and raised blood pressure, but the associations with raised triglycerides and fasting glucose were attenuated and no longer significant. After similar adjustments, low leisure-time physical activity at age 16 years remained associated with central obesity and raised triglycerides at age 43 years.

**CONCLUSIONS**—To our knowledge, this is the first prospective study examining the relationship between TV viewing and participation in leisure-time physical activity in adolescence with the metabolic syndrome and its components in mid-adulthood. We found that both TV viewing and low leisure-time physical activity in adolescence independently predicted the metabolic syndrome and several of the metabolic syndrome components in mid-adulthood. Our results suggested a dose-response relationship for both TV viewing and leisure-time physical activity with subsequent cardiometabolic risk. Furthermore, our analyses indicate that low leisure-time physical activity in adolescence is mediated by leisure-time physical activity in mid-adulthood to a considerable degree, whereas TV viewing in adolescence is not.

The finding of an association between TV viewing and cardiometabolic risk is consistent with the results from several previous cross-sectional studies on adults (14–17). Our results also expand previous longitudinal findings (4,5,18,19) by showing that the link between TV viewing and cardiometabolic risk stretches over a 27-year period from adolescence and into midlife. Consistent with previous studies (15,20), TV viewing and leisure-time physical activity were not closely related at age 16 years. Moreover, the relationship between TV viewing and the metabolic syndrome was not dependent on leisure-time physical activity in mid-adulthood, which may provide further support for the hypothesis that the association between sedentary behavior, such as TV viewing, and cardiometabolic risk is not mediated by physical activity habits (2). The association between low adolescent leisure-time physical activity and the metabolic syndrome, on the other hand, appeared to be mediated to a large extent by leisure-time physical activity in mid-adulthood. Sedentary behavior and

**Table 1—Selected characteristics according to TV-viewing habits and participation in leisure-time physical activity at age 16 years for the 888 participants from the Northern Swedish Cohort**

	n	TV viewing at age 16 years			P for difference	
		One show/week or less (n = 137)	One show/day or every other day (n = 492)	Several shows/day (n = 257)		
Assessed at age 16 years						
Sex (male)	888	28.8	48.6	71.2	<0.001	
Socioeconomic disadvantage	883	38.7	34.3	46.1	0.007	
Family history of diabetes	881	6.6	3.3	4.3	0.219	
BMI (kg/m <sup>2</sup> )	888	20.1 (2.60)	19.9 (2.59)	19.9 (2.87)	0.388	
Daily intake of sweets/pastries	886	16.1	15.6	17.6	0.787	
Highest quintile of alcohol consumption	882	27.9	19.8	16.1	0.020	
Daily smoking	888	35.0	25.7	21.0	0.010	
Assessed at age 43 years						
Metabolic syndrome	888	19.7	24.1	36.2	<0.001	
Metabolic syndrome components						
Central obesity	884	56.6	57.8	64.6	0.150	
Raised triglycerides	883	16.1	22.7	32.0	0.001	
Low HDL-C	883	20.4	29.0	32.8	0.035	
Raised blood pressure	888	31.4	38.9	50.2	0.001	
Raised fasting glucose	883	15.3	14.7	24.6	0.003	
Leisure-time physical activity at age 16 years						
	n	Daily (n = 80)	Several times/week (n = 348)	Once a week (n = 251)	Several times/month or less (n = 209)	P for difference
Assessed at age 16 years						
Sex (male)	888	61.3	55.7	40.2	57.4	<0.001
Socioeconomic disadvantage	883	32.5	37.8	32.8	48.5	0.004
Family history of diabetes	881	1.3	3.8	2.8	7.2	0.051
BMI (kg/m <sup>2</sup> )	888	19.7 (2.18)	20.0 (2.56)	19.9 (2.80)	19.8 (2.87)	0.622
Daily intake of sweets/pastries	886	11.3	14.7	19.2	17.3	0.274
Highest quintile of alcohol consumption	882	10.0	11.6	20.1	37.7	<0.001
Daily smoking	888	8.8	14.1	28.3	48.8	<0.001
Assessed at age 43 years						
Metabolic syndrome	888	16.3	24.1	26.3	36.4	0.001
Metabolic syndrome components						
Central obesity	884	53.8	55.2	57.0	72.2	<0.001
Raised triglycerides	883	15.0	23.7	23.2	30.4	0.042
Low HDL-C	883	23.8	26.6	30.4	32.4	0.332
Raised blood pressure	888	38.8	38.5	43.4	43.1	0.564
Raised fasting glucose	883	16.3	15.7	16.9	22.5	0.210

Data are % or means (SD) unless otherwise indicated.

physical activity have also been suggested to have different environmental determinants (21), which makes it likely that different strategies may need to be adopted within interventions targeting sedentary behavior compared with physical activity behavior.

We found that TV viewing in adolescence, in contrast to leisure-time physical activity, was associated with raised blood pressure and low HDL-C in mid-adulthood. These differences indicate

that the associations between sedentary behavior and low leisure-time physical activity with subsequent metabolic risk may be mediated by different cardiometabolic pathways. A longitudinal association between sedentary behavior and hypertension has previously been shown in adults (18,22), but we have found no previous study reporting a significant association between sedentary behavior in adolescence and hypertension in adulthood. Sedentary behavior has been

reported to predict lipid levels in several previous studies on different age-groups. Ki et al. (23) reported that TV viewing during early adulthood (at age 23 years) was inversely associated with HDL-C in mid-adulthood (at age 45 years). Hancox, Milne, and Poulton (5) found that TV viewing during childhood and adolescence was associated with total cholesterol in early adulthood (at age 26 years). The likely mechanisms by which sedentary behavior may affect lipid

Table 2—ORs for presence of the metabolic syndrome at age 43 years according to TV viewing and leisure-time physical activity at age 16 years in 869 participants

	n	Crude	Model 1 <sup>a</sup>	Model 2 <sup>b</sup>	Model 3 <sup>c</sup>
TV viewing					
One show/week or less	136	1.00	1.00	1.00	1.00
One show/day or every other day	483	1.31 (0.81–2.10)	1.39 (0.86–2.24)	1.39 (0.83–2.31)	1.30 (0.78–2.18)
Several shows/day	250	2.46 (1.50–4.06)	2.47 (1.50–4.09)	2.14 (1.24–3.71)	1.96 (1.13–3.42)
P for trend		<0.001	<0.001	0.003	0.008
Leisure-time physical activity					
Daily	78	1.00	1.00	1.00	1.00
Several times/week	342	1.58 (0.83–3.01)	1.51 (0.79–2.89)	1.44 (0.74–2.81)	1.32 (0.67–1.59)
Once a week	246	1.76 (0.91–3.40)	1.69 (0.87–3.28)	1.76 (0.88–3.52)	1.55 (0.77–3.12)
Several times/month or less	203	2.81 (1.45–5.44)	2.64 (1.35–5.15)	2.31 (1.13–4.69)	1.82 (0.88–3.77)
P for trend		<0.001	0.001	0.007	0.064

Data are OR (95% CI) unless otherwise indicated. <sup>a</sup>TV viewing and leisure-time physical activity in the same model. <sup>b</sup>Adjusted for model 1 variables plus sex, socioeconomic disadvantage, family history of diabetes, BMI, intake of sweets/pastries, alcohol consumption, smoking, and TV viewing/leisure-time physical activity at age 16 years. <sup>c</sup>Adjusted for model 2 variables plus physical activity at age 43 years.

metabolism are not clear, but animal studies have suggested that sedentary behavior suppresses the activity of the enzyme lipoprotein lipase in skeletal muscles (24). Lipoprotein lipase is the rate-limiting enzyme for hydrolysis of triglyceride-rich lipoproteins, which contributes to the formation of HDL-C. Whether a similar relationship exists in humans awaits the findings from experimental research currently underway.

Aside from the possible physiological mechanisms, it has been suggested that sedentary behavior may influence cardiometabolic risk via other behavioral mechanisms that are linked to sedentary behavior. This hypothesis is supported by results from previous research showing that students reporting medium or high TV viewership snacked more frequently while watching TV (25). In our study, we adjusted for general intake of sweets/

pastries but could unfortunately not control for snacking related to TV viewing. It has also been suggested that sedentary behavior increases cardiometabolic risk via a psychosocial pathway. For example, in a prospective study of university graduates by Sanchez-Villegas et al. (26), sedentary behavior was reported to be associated with increased risk for mental disorders (diagnosed depression, anxiety, or stress) at follow-up. This and other components of a psychosocial pathway need to be further investigated, and we have an excellent opportunity to continue such work within the Northern Swedish Cohort.

The finding of an association between low leisure-time physical activity at age 16 years and the metabolic syndrome at age 43 years is consistent with several previous reports (27–29). Furthermore, low leisure-time physical activity in adolescence was associated with central obesity and raised triglycerides in mid-adulthood. Interestingly, these prospective associations between TV viewing and low leisure-time physical activity with the metabolic syndrome were not dependent on BMI at age 16 years. This indicates that both sedentary behavior and physically inactive behavior may precede overweight in a potential causal chain from adolescent behavior to adult cardiometabolic risk. However, this will need to be further explored within well-controlled intervention studies.

The strengths of this study include the prospective design and the high proportion of the original cohort that was assessed for the metabolic syndrome according to standardized criteria. The low attrition to follow-up also enabled us to




Figure 1—ORs for presence of the metabolic syndrome (95% CI) at age 43 years according to leisure-time physical activity and TV viewing (adjusted for sex, socioeconomic disadvantage, family history of diabetes, BMI, intake of sweets/pastries, alcohol consumption, and smoking). TV viewing one show per day or less, ●; TV viewing several shows a day, ○.

examine the internal dropout, which did not indicate that the dropout was systematic with respect to key measures. The study was limited by the crude assessment of TV-viewing habits and participation in leisure-time physical activity. At 16 years of age, BMI and several lifestyle factors were collected, but the metabolic syndrome components were not assessed. If individuals watching a lot of TV in adolescence were on the projection to the metabolic syndrome, this could bias the results, since adjustment for BMI and several lifestyle factors may not fully compensate for missing information on metabolic syndrome components at baseline. Information was also missing for TV viewing at age 43 years. TV viewing at different ages has been shown to be moderately correlated (18,19,30), which indicates that viewing habits established in adolescence may persist into adulthood. However, several studies have found that TV viewing in childhood and adolescence is a stronger predictor of adverse health outcomes in adulthood than TV viewing in adulthood (5,31). It is therefore likely that the association between TV viewing and the metabolic syndrome in adulthood is modified rather than mediated by TV viewing in adulthood. Moreover, although TV viewing was the most important indicator of sedentary behavior during leisure-time in the early 1980s, before computers were commonly available (32), several other sedentary behaviors at school or the workplace and during commuting and leisure time may be linked to TV viewing and of relevance for cardiometabolic health. Leisure-time physical activity was also crudely assessed, and therefore we cannot rule out residual confounding from sedentary behaviors and leisure-time physical activity that were not assessed in this study.

In conclusion, this study supports previous findings that prolonged TV viewing is independently associated with subsequent metabolic risk and provides some new evidence that this association may stretch over a considerable proportion of the lifespan: from adolescence to mid-adulthood. Low leisure-time physical activity in adolescence was also independently associated with metabolic risk in mid-adulthood, and this study indicates that the associations between sedentary behavior and low leisure-time physical activity with subsequent metabolic risk may be mediated by different cardiometabolic pathways. These findings suggest that reduced TV viewing in

**Table 3—ORs for the presence of the metabolic syndrome components at age 43 years according to TV viewing and leisure-time physical activity at age 16 years in 869 participants**

	Central obesity		Raised triglycerides		Low HDL-C		Raised blood pressure		Raised fasting glucose	
	Crude OR (95% CI)	Multivariate OR (95% CI) <sup>a</sup>	Crude OR (95% CI)	Multivariate OR (95% CI) <sup>a</sup>	Crude OR (95% CI)	Multivariate OR (95% CI) <sup>a</sup>	Crude OR (95% CI)	Multivariate OR (95% CI) <sup>a</sup>	Crude OR (95% CI)	Multivariate OR (95% CI) <sup>a</sup>
TV viewing										
One show/week or less	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
One show/day or every other day	1.04 (0.78–1.53)	1.25 (0.83–1.90)	1.61 (0.97–2.69)	1.41 (0.81–2.45)	1.63 (1.03–2.60)	1.90 (1.17–3.06)	1.39 (0.92–2.09)	1.36 (0.88–2.10)	0.92 (0.54–1.57)	0.84 (0.48–1.48)
Several shows/day	1.43 (0.93–2.19)	1.72 (1.07–2.77)	2.64 (1.55–4.51)	1.73 (0.96–3.12)	1.98 (1.20–3.26)	2.23 (1.32–3.78)	2.20 (1.42–3.42)	1.78 (1.10–2.87)	1.86 (1.07–3.20)	1.34 (0.74–2.44)
<i>P</i> for trend	0.060	0.020	0.001	0.064	0.009	0.005	<0.001	0.016	0.003	0.124
Leisure-time physical activity										
Daily	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Several times/week	1.06 (0.65–1.74)	0.95 (0.57–1.59)	1.72 (1.89–3.34)	1.77 (0.89–3.52)	1.12 (0.63–1.98)	1.03 (0.58–1.83)	1.05 (0.63–1.74)	1.01 (0.59–1.71)	0.91 (0.47–1.76)	0.80 (0.40–1.59)
Once a week	1.12 (0.67–1.86)	0.99 (0.57–1.71)	1.67 (0.84–3.30)	2.14 (1.04–4.40)	1.37 (0.76–2.46)	1.28 (0.70–2.34)	1.26 (0.74–2.13)	1.38 (0.79–2.42)	1.04 (0.53–2.06)	1.01 (0.49–2.08)
Several times/month or less	2.14 (1.25–3.68)	1.90 (1.05–3.42)	2.40 (1.21–4.75)	2.32 (1.11–4.87)	1.42 (0.78–2.58)	1.17 (0.62–2.20)	1.24 (0.73–2.13)	1.02 (0.57–1.84)	1.47 (0.74–2.89)	1.03 (0.49–2.17)
<i>P</i> for trend	0.001	0.008	0.018	0.033	0.119	0.389	0.232	0.607	0.061	0.458

<sup>a</sup>Adjusted for sex, socioeconomic disadvantage, family history of diabetes, BMI, intake of sweets/pastries, alcohol consumption, smoking, and TV viewing/leisure-time physical activity at age 16 years.

adolescence, in addition to and independently of regular leisure-time physical activity in adolescence and adulthood, may contribute to cardiometabolic health later in life.

**Acknowledgments**—The study was supported by grants from the County Council of Västerbotten, the Swedish Research Council, and the Swedish Council for Working Life and Social Research. P.E.G. is supported by Umeå University. D.W.D. is supported by an Australian Research Council Future Fellowship.

No potential conflicts of interest relevant to this article were reported.

P.W. was responsible for the concept and design of the study, gave final approval of the version to be published, obtained funding, analyzed data, wrote early drafts, and wrote the final manuscript. P.E.G. was responsible for the concept and design of the study, gave final approval of the version to be published, obtained funding, made substantial contributions to analysis and interpretation of data, and revised the manuscript critically for important intellectual content. D.W.D. was responsible for the concept and design of the study, gave final approval of the version to be published, made substantial contributions to analysis and interpretation of data, and revised the manuscript critically for important intellectual content. M.W. was responsible for the concept and design of the study, gave final approval of the version to be published, obtained funding, made substantial contributions to analysis and interpretation of data, and revised the manuscript critically for important intellectual content. A.H. was the principal investigator, conceived the Northern Swedish Cohort, collected data, was responsible for the concept and design of the study, gave final approval of the version to be published, obtained funding, made substantial contributions to analysis and interpretation of data, and revised the manuscript critically for important intellectual content. P.W. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

## References

- Haskell WL, Lee IM, Pate RR, et al.; American College of Sports Medicine; American Heart Association. Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Circulation* 2007;116:1081–1093
- Thorp AA, Owen N, Neuhaus M, Dunstan DW. Sedentary behaviors and subsequent health outcomes in adults: a systematic review of longitudinal studies, 1996–2011. *Am J Prev Med* 2011;41:207–215
- Spanier PA, Marshall SJ, Faulkner GE. Tackling the obesity pandemic: a call for sedentary behaviour research. *Can J Public Health* 2006;97:255–257
- Wijndaele K, Healy GN, Dunstan DW, et al. Increased cardiometabolic risk is associated with increased TV viewing time. *Med Sci Sports Exerc* 2010;42:1511–1518
- Hancox RJ, Milne BJ, Poulton R. Association between child and adolescent television viewing and adult health: a longitudinal birth cohort study. *Lancet* 2004;364:257–262
- Robinson TN. Reducing children's television viewing to prevent obesity: a randomized controlled trial. *JAMA* 1999;282:1561–1567
- Hammarström A, Janlert U. Cohort profile: the northern Swedish cohort. *Int J Epidemiol* 2011;41:1545–1552
- Gustafsson PE, Persson M, Hammarström A. Life course origins of the metabolic syndrome in middle-aged women and men: the role of socioeconomic status and metabolic risk factors in adolescence and early adulthood. *Ann Epidemiol* 2011;21:103–110
- World Health Organization. Cardiovascular Diseases Unit. *MONICA Manual: WHO MONICA Project*. Geneva, World Health Org., 1990
- Gustafsson PE, Hammarström A. Socio-economic disadvantage in adolescent women and metabolic syndrome in mid-adulthood: an examination of pathways of embodiment in the Northern Swedish Cohort. *Soc Sci Med* 2012;74:1630–1638
- Gustafsson PE, Janlert U, Theorell T, Westerlund H, Hammarström A. Do peer relations in adolescence influence health in adulthood? Peer problems in the school setting and the metabolic syndrome in middle-age. *PLoS ONE* 2012;7:e39385
- Alberti KG, Zimmet P, Shaw J; IDF Epidemiology Task Force Consensus Group. The metabolic syndrome—a new worldwide definition. *Lancet* 2005;366:1059–1062
- Statistics Sweden. *Swedish Socioeconomic Classification. Reports on Statistical Co-ordination 1982:4*. Stockholm, Statistics Sweden, 1984
- Gustat J, Srinivasan SR, Elkasabany A, Berenson GS. Relation of self-rated measures of physical activity to multiple risk factors of insulin resistance syndrome in young adults: the Bogalusa Heart Study. *J Clin Epidemiol* 2002;55:997–1006
- Dunstan DW, Salmon J, Owen N, et al.; AusDiab Steering Committee. Associations of TV viewing and physical activity with the metabolic syndrome in Australian adults. *Diabetologia* 2005;48:2254–2261
- Chang PC, Li TC, Wu MT, et al. Association between television viewing and the risk of metabolic syndrome in a community-based population. *BMC Public Health* 2008;8:193
- Gardiner PA, Healy GN, Eakin EG, et al. Associations between television viewing time and overall sitting time with the metabolic syndrome in older men and women: the Australian Diabetes, Obesity and Lifestyle study. *J Am Geriatr Soc* 2011;59:788–796
- Stamatakis E, Hamer M, Mishra GD. Early adulthood television viewing and cardiometabolic risk profiles in early middle age: results from a population, prospective cohort study. *Diabetologia* 2012;55:311–320
- Grøntved A, Ried-Larsen M, Møller NC, et al. Youth screen-time behaviour is associated with cardiovascular risk in young adulthood: the European Youth Heart Study. *Eur J Prev Cardiol*. 5 July 2012 [Epub ahead of print]
- Hu FB, Li TY, Colditz GA, Willett WC, Manson JE. Television watching and other sedentary behaviors in relation to risk of obesity and type 2 diabetes mellitus in women. *JAMA* 2003;289:1785–1791
- Owen N, Leslie E, Salmon J, Fotheringham MJ. Environmental determinants of physical activity and sedentary behavior. *Exerc Sport Sci Rev* 2000;28:153–158
- Beunza JJ, Martínez-González MA, Ebrahim S, et al. Sedentary behaviors and the risk of incident hypertension: the SUN Cohort. *Am J Hypertens* 2007;20:1156–1162
- Ki M, Poulidou T, Li L, Power C. Physical (in)activity over 20 y in adulthood: associations with adult lipid levels in the 1958 British birth cohort. *Atherosclerosis* 2011;219:361–367
- Zderic TW, Hamilton MT. Physical inactivity amplifies the sensitivity of skeletal muscle to the lipid-induced downregulation of lipoprotein lipase activity. *J Appl Physiol* 2006;100:249–257
- Thomson M, Spence JC, Raine K, Laing L. The association of television viewing with snacking behavior and body weight of young adults. *Am J Health Promot* 2008;22:329–335
- Sanchez-Villegas A, Ara I, Guillén-Grima F, Bes-Rastrollo M, Varo-Cenarruzabeitia JJ, Martínez-González MA. Physical activity, sedentary index, and mental disorders in the SUN cohort study. *Med Sci Sports Exerc* 2008;40:827–834
- Laaksonen DE, Lakka HM, Salonen JT, Niskanen LK, Rauramaa R, Lakka TA. Low levels of leisure-time physical activity and cardiorespiratory fitness predict development of the metabolic syndrome. *Diabetes Care* 2002;25:1612–1618
- Wareham NJ, Hennings SJ, Byrne CD, Hales CN, Prentice AM, Day NE. A quantitative analysis of the relationship between habitual energy expenditure,

- fitness and the metabolic cardiovascular syndrome. *Br J Nutr* 1998;80:235–241
29. Rennie KL, McCarthy N, Yazdgerdi S, Marmot M, Brunner E. Association of the metabolic syndrome with both vigorous and moderate physical activity. *Int J Epidemiol* 2003;32:600–606
30. Parsons TJ, Manor O, Power C. Television viewing and obesity: a prospective study in the 1958 British birth cohort. *Eur J Clin Nutr* 2008;62:1355–1363
31. Erik Landhuis C, Poulton R, Welch D, Hancox RJ. Programming obesity and poor fitness: the long-term impact of childhood television. *Obesity* (Silver Spring) 2008;16:1457–1459
32. Clark BK, Sugiyama T, Healy GN, Salmon J, Dunstan DW, Owen N. Validity and reliability of measures of television viewing time and other non-occupational sedentary behaviour of adults: a review. *Obes Rev* 2009;10:7–16